

COMPLICIAL SETS

DOMINIC VERITY

ABSTRACT. The primary purpose of this work is to characterise strict ω -categories as simplicial sets with structure. We prove the Street-Roberts conjecture which states that they are exactly the “complicial sets” defined and named by John Roberts in his handwritten notes of that title [27].

CONTENTS

1. Introduction	3
1.1. Historical Background	3
1.2. Relationships to Other Work	5
1.3. Overview and Structure	7
2. Simplicial Operators and Simplicial Sets	10
2.1. Simplicial Operators	10
2.2. The Algebraist’s Δ and 2-Categories	12
2.3. The Algebraist’s Δ and Monoidal Categories	15
2.4. Simplicial Sets	18
2.5. Semi-Simplicial Sets	22
2.6. Analysing Products of Simplicial Sets - the Theory of Shuffles	23
3. Some Categorical Background	28
3.1. Reflective Full Sub-Categories	28
3.2. LFP Categories and LE-Theories	32
4. Double Categories, 2-Categories and n-Categories	39
4.1. Categories in the Small	39
4.2. Double Categories	42
4.3. 2-Categories and Double Categories with Connections	43
4.4. n -Categories and ω -Categories	48
5. An Introduction to the Decalage Construction	51
5.1. Nerves and Decalage	51
5.2. Comonad Transformations and Simplicial Reconstruction	53
6. Stratifications and Filterings of Simplicial Sets	59
6.1. Stratified Simplicial Sets	59
6.2. Superstructures and Filtered Semi-Simplicial Sets	64
7. Pre-Complicial Sets	68
7.1. Introducing Pre-Complicial Sets	68
7.2. Tensor Products of Pre-Complicial Sets	69
7.3. Pre-Tensors and Preservation of t-Extensions	73

Date: October 15, 2004.

2000 *Mathematics Subject Classification.* Primary 18D05, 55U10; Secondary 18D15, 18D20, 18D35, 18F99, 18G30.

7.4.	Some Other Preservation Properties	80
7.5.	A Monoidal Biclosed Structure on Pre-Complcial sets	82
7.6.	Superstructures of Pre-Complcial Sets	85
8.	Complcial Sets	86
8.1.	Introducing Complcial Sets	86
8.2.	Pasting Squares and Filling Lemmas	87
8.3.	Tensor Products and Complcial Sets	91
8.4.	Superstructures of Complcial Sets	102
9.	The Path Category Construction	104
9.1.	The Complcial Category of Prisms	104
9.2.	Path Categories and Superstructures	107
9.3.	A Complcial Double Category with Connections	109
10.	Decalage for Complcial Sets and Complicially Enriched Categories	113
10.1.	A Decalage Construction on Complcial Sets	113
10.2.	A Path Construction on Complicially Enriched Categories	118
10.3.	A Decalage Construction on Complicially Enriched Categories	122
10.4.	Semi-Simplicial Reconstruction	124
11.	Street's ω -Categorical Nerve Construction	129
11.1.	Parity Complexes	129
11.2.	Collapsers and Stratified Parity Complexes	142
11.3.	ω -Categorical Nerve Constructions	145
11.4.	Relating Parity Complex Products and the Tensor Product of Stratified Sets	150
11.5.	An Inductive Proof of the Street-Roberts Conjecture	157
	References	166

1. INTRODUCTION

1.1. Historical Background. This work presents a proof of a result that has sometimes been referred to as the Street-Roberts conjecture (after [31]). This postulates an equivalence between the category of *strict* ω -categories and a category of structures called *complicial sets* which are certain kinds of enhanced simplicial sets originally studied by Roberts [27].

The genesis of this work dates back to the mid-1970 and Roberts' work on non-abelian cohomology. His original interest in this topic grew from his conviction that (strict) ω -categories were the appropriate algebraic structures within which to value such theories [26]. This led him to define complicial sets to be simplicial sets with distinguished elements, which he originally referred to as "neutral" then later as "hollow" but for which we prefer the term *thin* (after [9]), satisfying some natural conditions related to those that characterise *Kan complexes* in the homotopy theory of simplicial sets [13].

In particular, his conditions include a certain kind of *unique thin horn filler* condition. These closely resemble the horn fillers of Kan, with the notable exceptions that only certain *admissible* horns (for which a specified class of faces are required to be thin) are assumed to have fillers, that those fillers are all themselves thin and that all such fillers are assumed to be unique.

Roberts was motivated in this definition by his observation that it should be possible to naturally generalise the classical nerve constructions of Algebraic Topology to provide a functor from the category of (strict) ω -categories to the category of simplicial sets which, in a suitable sense, encapsulated a natural notion of *higher non-abelian cocycle*. While the nerve construction on groups and partially ordered sets, and their common generalisation to categories, is well known and easily described to students of Algebraic Topology, the same cannot be said of its generalisation to ω -categories. Indeed its very definition poses substantial technical challenges which eluded Roberts at the time.

Even without an explicit construction of this nerve functor, Roberts set about studying those simplicial sets that would occur as the nerve of some ω -category. He observed that this functor would not provide a fully-faithful representation and suggested rectifying this failing by introducing thin elements into his study. He then set to characterising those augmented simplicial sets which would arise in the replete image of the postulated fully-faithful functor and thus complicial sets were born.

One might forgive Roberts if his identification of this category were to prove deficient in some way, after all he was working without a nerve functor and at a time when the theory of strict ω -categories had advanced little further than fundamental definitions. It is therefore a great tribute to Roberts' insight that our main theorem here (theorem 266 of section 11) establishes his sought-for equivalence under a definition of complicial set which only differs from his in manner of expression. He provided a precise conjecture in 1978, our only contribution has been to prove it as stated!

The next big advance in this story came with Street's papers on orientals [31] and parity complexes [33]. He was introduced to this study, during a visit Roberts made to Sydney in the (southern hemisphere) summer of 1977-8, and was captivated by it. While he was quickly able to establish the sought for equivalence at dimension 2, it became apparent to him that establishing the general equivalence would be quite a difficult problem. Consequently he decided to concentrate on providing a rigorous definition of the nerve of an ω -category.

After a couple of false starts to that end, he soon realised that the crux of the matter was to define the free ω -category $\mathcal{O}_n$ on an n -simplex. However, in order to make this insight

precise it was necessary to define exactly the kinds of combinatorial structures that might give rise to free ω -categories and even to explicate the sense in which freeness itself should be interpreted. Furthermore, once these concepts had been defined it would still be necessary to define exactly how an n -simplex might be considered to be such a structure.

The first of these questions elicited the introduction of certain kinds of inductively defined combinatorial structures called ω -computads, which provide the desired definitions but are often inconvenient to calculate with. To facilitate the work we are engaged in here, Street later introduced a restricted form of ω -computad, called a *parity complex*. These satisfy some very strong loop-freeness conditions designed to ensure that we may describe the cells of the associated free ω -category as (pairs of) subsets of the parity complex itself.

To understand how one might render an n -simplex as a parity complex, Street started by observing that $\mathcal{O}_n$ should play the part of some form of non-abelian n -cocycle in the sought after cohomology theory. This led him to conjecture that we could take insight from abelian cohomology and build a consistently oriented parity complex whose elements were the faces of a standard n -simplex, each of which should be oriented as a cell to “map from odd numbered faces to even numbered ones”. Under this definition, he was able to demonstrate that the resulting structure satisfied the strong loop-freeness conditions required of a parity complex and thereby provide a completely explicit description of the free ω -category $\mathcal{O}_n$.

Street completed the construction of his ω -categorical nerve by enriching his orientals to a functor from the *category of simplicial operators* Δ and applying Kan’s construction [17]. His paper [31] goes on to lift this functor to map ω -categories to simplicial sets with thin elements (which we call stratified sets in the sequel) and to formalise Roberts’ original conjecture in this context. He identifies a family of admissible horns which *strictly* contains the class introduced by Roberts and in [32] he demonstrates that within the nerve of an ω -category such horns do indeed have unique thin fillers. It is worth noting, however, that while Street’s complicial set definition appears, at first sight, to be stronger than Roberts’, our proof here demonstrates that they are in fact equipotent.

My own contribution to this story began in 1991 when I first read [31] and immediately became captivated by this problem. In particular, I had been searching for an approach to defining structures which we now call *weak ω -categories* and became convinced that Street’s parenthetical remarks regarding a potential simplicial definition held great merit. In ignorance of [32], I set about proving that in any nerve Street’s admissible horns had unique fillers and in a very short time, and to my great pleasure, I succeeded in constructing a decomposition of each oriental which established this result.

Spurred on by my initial success, I decided to turn my hand to proving Street’s version of Roberts’ original conjecture and soon succeeded in showing that Street’s nerve functor was fully faithful. While this result in itself was a clear contribution to the then extant state of knowledge in this area, I was not satisfied since the method used to do so had originally been conceived as a proof of the complete Street-Roberts conjecture. My initial reaction to this block was to cast around for a categorical abstraction which might strengthen my result, but to no avail. Somewhat disheartened by this, I wrote notes on my proof to date and circulated this handwritten manuscript to a few interested parties (including Street) before settling down to complete my PhD work on enriched category theory.

Returning to this problem in 1993, I realised that an argument based upon the decalage construction would close the gap in my proof. In 1994 I gave talks on the resulting proof to the Sydney (now Australian) Category Seminar, at the University College of North Wales

in Bangor, at the Mathematical Sciences Research Institute in Berkley and to a Peripatetic Seminar on Sheaves and Logic at the Newton Institute in Cambridge UK.

Unfortunately, however, career events overtook me before I had time to commit the final proof to paper. In early 1995 I followed up a series of consultancy engagements in the financial markets by accepting a full time role in investment banking. I spent the subsequent 5 years as a derivative securities analyst and trader manager before returning to academia in mid 2000.

On returning to this field, two things struck me immediately. Firstly, and most pleasingly, an area of study that had given so much joy to a small band of enthusiasts had grown into a dynamic area of wide debate and interest, driven by an influx of new ideas and approaches. Secondly, and much to my relief, nobody appeared to have provided a proof of the Street-Roberts result upon which I had devoted so much time prior to my sojourn into the business world.

The proof presented herein is fundamentally no different to the one that I spoke on in 1994. The primary innovation on that original work has been the adoption of a *(lax) Gray tensor product* of complicial sets as a unifying organisational and constructional tool.

I should like to dedicate this work to three groups of people. Firstly of course to my wife, Sally, and children, Lottie and Florrie, who have put up with a husband and father incessantly crouched over the word processor. They have never once questioned the importance to me of completing this work and have sustained me in body and soul over the 12 years it has been in gestation. Secondly I would like to thank Ross Street, who has been the most formative influence in my development as a mathematician. It was his friendship, support and inspiration which convinced me to return to academe and to work which I had long since convinced myself I would never commit to paper. In the end it was his quiet reminders, to the effect that I owed it to the community to write up my ideas, that spurred me to find the time in a busy schedule to write these scratchings. Finally I should like to thank the staff of the Postgraduate Professional Development Program of the Division of Information and Communication Sciences at Macquarie University, of which I am the academic director. Without their support and hard work, filling in for me while I wrote this work, I would never have been able to find the silence and space to organise these ideas.

1.2. Relationships to Other Work. While this work predominantly interests itself in a study of strict ω -categorical structures, it is nevertheless squarely motivated by a broader program to define and study weakened ω -categorical notions. In particular, its purpose from the very start has been to act as a first step toward providing a full account of the theory of Street’s simplicial weak ω -category notion, which first appeared in sketch form in his orientals paper [31] and upon which he later elaborated in [36].

Indeed, much of the work on (lax) Gray tensor product of complicial sets given here routinely generalises to Street’s category of weak complicial sets. In particular, using these generalisations we may show that the category of weak complicial sets supports biclosed structures generalising those on the category of bicategories and homomorphisms as discussed in [30]. These allow us to enrich the category of weak complicial sets over itself in natural ways and opens the possibility of providing a coherence result for Street’s weak ω -categories along the lines of that described for tricategories in [14].

Notice however that, just as in the bicategorical and tricategorical cases, the corresponding “monoidal structure” on weakly complicial sets is only *weakly* adjoint (in some suitable sense) to this biclosed structure. As a result it is only weakly coherent, making it somewhat

inconvenient to calculate with. One convenience provided by the weak complicial approach, however, is that we can always perform all required calculations in a bigger category of pre-complicial sets, on which the corresponding tensor is actually part of a genuine monoidal biclosed structure.

The reader will, of course, be aware that over the past ten years a wide variety of weak ω -categorical notions have been proposed by various authors. Notable amongst these are those of Joyal [16], Batanin [4] and Baez and Dolan [2], although more may be found in the literature or in Leinster’s survey of the current state of the definitional art [23]. Many of these drew their initial inspiration from Street’s parenthetical remarks in [23] although none of them, except for the summary given in Leinster [23] and Street’s own account [36], expand directly on the purely simplicial approach analysed here. This work starts from a point of view neatly summed up by Street’s comment in [36] that a simplicial formulation bears the distinct advantage that:

Simplicial sets are lovely objects about which algebraic topologists know a lot. If something is described as a simplicial set, it is ready to be absorbed into topology. Or, in other words, no matter which definition of weak ω -category eventually becomes dominant, it will be valuable to know its simplicial nerve.

In many ways, our work here also parallels work of Ronald Brown and his coworkers, which established analogous results for the somewhat simpler groupoidal case. While our work and theirs share at least one common motivation, to develop a coherent and complete theory of higher dimensional cohomology, their primary interest is not the development of an encompassing theory of weak ω -categories but rather the explication of a theory within which to make explicit calculations of homotopic invariants. To that end a study of groupoids rather than categories is adequate for most purposes.

The complicial sets discussed here are most closely related to their simplicial T-complexes, which were first discussed in Dakin’s thesis [9], where the term “thin” was first coined, and later developed and popularised in the work of Brown, Higgins and others. Since this work was groupoid oriented, the definitions involved are somewhat simpler than those discussed here and a proof of the equivalence between simplicial T-complexes and ω -groupoids is far more easily attained. Indeed, a proof of this equivalence in the rank 2 case dates to Dakin’s thesis in 1977.

While the simplicial approach is important to their work, it should be pointed out that in general their preference has been to work in a cubical context. This allows them to more easily build generalised van Kampen Theorems, for computing homotopical invariants, and to define monoidal closed structures for analysing homotopy classes of maps. A comprehensive and up-to-date account of this work may be found in Brown’s excellent survey article [5]. In this cubical program the closest result to the one presented here is that of Al-Agl, Brown and Steiner [1] which establishes an equivalence between the category of (strict) ω -categories and a category of multiple (cubical) categories with connection.

While this cubical approach is attractive for the reasons discussed above, it is our conviction that it does not provide quite such a convenient context within which to develop a theory of weak ω -categories. Nevertheless cubical calculations, and most particularly those involved in a theory of tensor products, remain vital even in the simplicial context. The theory discussed here, therefore, appears to combine the best of both worlds by allowing for a theory with all of the beauty, elegance and economy of the simplicial approach while at the same time providing a simple and highly explicit description of the cubical Gray tensor product.

1.3. Overview and Structure. This work is almost exclusively devoted to proving the Street-Roberts conjecture, in the form presented in [31]. On the way we substantially develop Roberts' theory of complicial sets [26] itself and make some contributions to Street's theory of parity complexes [33]. In particular, we study a new monoidal closed structure on the category of complicial sets which we show to be the appropriate generalisation of the (lax) Gray tensor product of 2-categories to this context. Under the equivalence conjectured by Street [31] and Roberts [26], which we prove here, this tensor product coincides with those of Crans [8], Steiner [28] and others.

From the outset, it has been designed to be as self contained as possible. While much of the material covered in its first five sections is classical in nature, it has been presented here in order to fix our notation for the sequel and to aggregate together a number of familiar (and not so familiar) pieces of Algebraic Topology and Category Theory from a diverse range of sources. Given the influx into this field of mathematicians from diverse backgrounds, it was felt prudent that no prerequisite assumptions be made. Most particularly, it was recognised that some readers might not be fully conversant with certain of the more abstract aspects of general (enriched and internal) category theory and higher category theory.

In particular, where categorical abstractions are concerned we assume little more than that the reader should have a general grounding in basic categorical concepts, such as functor, natural transformation, limit, colimit, comma category and 2-category etcetera, all of which may be gleaned from Mac Lane's book [22].

Congruent with this philosophy, sections 2 to 5 are all contextual in nature, providing fairly standard presentations of traditional material. Section 2 consists of a brief introduction to the theory of simplicial sets, up to and including the theory of shuffles. While this is **not** intended to provide an exhaustive treatment of the simplicial algebra necessary to read this work, it should provide most of the necessary background and adequate pointers to the available literature.

At some points in the sequel our arguments are substantially simplified by couching them in more abstract categorical terms, along the lines described in Kelly's book [19]. In particular, our construction of the monoidal biclosed Gray tensor structure on complicial sets relies on Day's reflection theorem for monoidal biclosed categories [10] and many of our later constructions and calculations are couched in terms of left exact theories and their coalgebras. While a thorough reading of Kelly's book would handsomely repay the effort involved, the results of greatest interest here are collected together in section 3; the reader should refer to the cited literature for detailed proofs.

Section 4 rehearses the basic definitions in the theory of (internal) categories, double categories and ω -categories. We also remind the reader of the relationship between double categories and 2-categories, by discussing the Spencer's recognition principle for those double categories that arise as the double categories of pasting squares in some 2-category (see Brown and Mosa [7]).

Section 5, the last of these contextual introductions, provides a account of the classical simplicial decalage construction and the method of simplicial reconstruction. We review those parts of the theory of (co)monads which were developed in order to provide a general way of constructing functors into categories of simplicial structures. In this context, it is a classical result that the category of simplicial sets supports a canonical comonad, called the decalage comonad, which is in some sense generic for this construction. Again, this material will be very familiar to Algebraic Topologists and Category Theorists but may be less familiar

to others and, indeed, the 2-categorical presentation we give here may be considered to be somewhat non-standard.

From section 6 we concentrate on developing the theory of complicial sets and for much of this work we study these as novel structures in their own right. Only much later, in section 11, do we “tie the knot” by relating our constructions back to the traditional theory of (strict) ω -categories. In order to do so we contribute to the theory of parity complexes and provide a deepened analysis of Street’s nerve functor.

One of the attractions of complicial sets as a foundation for ω -category theory, of both the strong and weak variety, is that they build upon the familiar theory of simplicial sets. However it is too much to hope that simplicial sets themselves are enough, especially since Street’s canonical nerve construction does not provide us with a full representation of ω -categories as simplicial sets. The issue here is that this nerve does not record enough information about the identities in our ω -categories, a deficiency we rectify by storing this missing data using a structure dubbed *hollowness* by Street but later renamed *stratification*. We examine the theory of such *stratified simplicial sets* in section 6.1 and later, in section 6.2, we discuss *filtered semi-simplicial sets* which may be used to provide an alternative foundation to our work and which we apply at a strategically important point in the sequel.

In section 7 we introduce a class of stratified sets which, for reasons that shall become apparent, we choose to call *pre-complicial sets*. Pre-complicial sets satisfy a relatively weak constraint, which allows us to make inferences about the thinness of simplices that are related by virtue of being faces on a suitable higher dimensional thin simplex. Our primary result in this section is that the full subcategory of pre-complicial sets admits a (non-symmetric) biclosed monoidal structure, which will become our main tool in much of what follows.

In section 8, we finally define Roberts’ *complicial sets* to be pre-complicial sets in which all *admissible horns* have unique thin fillers. Most of the effort in this section is devoted to extending the results of section 7 to the reflective full subcategory of complicial sets. In particular we show that we may reflect our biclosed monoidal structure on pre-complicial sets down onto the category of complicial sets.

In section 9, we use the biclosed monoidal structure on complicial sets to build internal and enriched categories from complicial sets. Specifically, we present a construction which allows us to derive a category of paths in a complicial set whose homsets are themselves complicial sets. Later we show that, in a suitable sense, we may iterate this construction to provide an explicit equivalence inverse to Street’s nerve functor.

In section 10 we build analogues of the traditional simplicial decalage construction on the categories of complicial sets and complicially enriched categories. Using an argument which deploys a certain complicial double category with connection derived in the previous section, we show that these constructions correspond to each other under the action of our path category functor. Applying simplicial reconstruction to these then establishes the fact that the path category construction provides us with a fully faithful representation of complicial sets as complicially enriched categories.

Finally, in section 11 we quickly review Street’s work on parity complexes [33], [34] and ω -categorical nerves [31], [32] and establish its connection to our work on the theory of complicial sets. In particular, we construct a canonical isomorphism between the free ω -category on the parity complex product of a pair of simplices and the ω -category obtained by reflecting the tensor product of corresponding standard simplices. This confirms our intuition that the complicial tensor generalises the (lax) Gray tensor product of 2-categories [15] and

that, under Street's nerve construction, it coincides with the ω -categorical tensor products of Crans [8] and Steiner [28].

Once we have done this we can partially free ourselves from the complicial world and discuss the all important relationship between Street's nerve construction and our own path category construction. Finally, we use this to provide a quick and easy proof that Street's nerve provides an **equivalence** between the categories of ω -categories and complicial sets, as originally conjectured in [31].

2. SIMPLICIAL OPERATORS AND SIMPLICIAL SETS

2.1. Simplicial Operators.

Definition 1 (The category Δ_+). Let Δ_+ denote the skeletal category of finite ordinals and order preserving functions. In other words, Δ_+ has:

- **Objects** ordered sets $[n] = \{0 < 1 < \dots < n\}$ one for each $n \geq -1$,
- **Maps** $\alpha: [n] \longrightarrow [m]$ which are order preserving functions from the ordered set $[n]$ to the ordered set $[m]$,
- **Composition** simply that of functions, which is well defined since the composite of order preserving maps is again order preserving.

We use the notation $\text{id}_{[n]}$ to denote the identity function on $[n]$. The maps of Δ_+ are often referred to as *simplicial operators*, for reasons which will become clear.

Notation 2 (The Topologist's Δ). The category Δ_+ is known as the *Algebraist's Δ* since, as recalled later on, this category “classifies” the algebraic theory of *monoids*. However, for much of this work we will be interested in studying simplicial operators from the topological perspective. To this end, we will often restrict our attention to Δ the full subcategory of Δ_+ whose objects are the **non-zero** ordinals $[n]$ for $n \geq 0$. Following the usual tradition, we will usually refer to this category as the *Topologist's Δ* .

Under the topological interpretation of Δ , the object $[n]$ is considered to be a combinatorial rendering of the standard n -dimensional simplex. This explains our rather peculiar, but nonetheless entirely standard, use of the notation $[n]$ to denote the $(n+1)^{\text{th}}$ ordinal.

Notation 3 (faces and degeneracies). The following standard notation and nomenclature will be used throughout:

- The injective maps in Δ_+ are referred to as *face operators*.
- For each $n \in \mathbb{N}$ and $j \in [n]$ define the simplicial operator $\delta_j^n: [n-1] \longrightarrow [n]$ by

$$\delta_j^n(i) = \begin{cases} i & \text{if } i < j, \\ i+1 & \text{otherwise.} \end{cases}$$

This is called the j^{th} *elementary face operator of $[n]$* .

- The surjective maps in Δ_+ are referred to as *degeneracy operators*.
- For each $n \in \mathbb{N}$ and $j \in [n]$ define the simplicial operator $\sigma_j^n: [n+1] \longrightarrow [n]$ by

$$\sigma_j^n(i) = \begin{cases} i & \text{if } i \leq j, \\ i-1 & \text{otherwise.} \end{cases}$$

This is called the j^{th} *elementary degeneracy operator of $[n]$* .

- For each $n \in \mathbb{N}$ and $i \in [n]$ the operator $\varepsilon_i^n: [0] \longrightarrow [n]$ given by $\varepsilon_i^n(0) = i$ is called the i^{th} *vertex operator of $[n]$* .
- For each $n \in \mathbb{N}$ we use the notation η^n to denote the unique operator from $[n]$ to $[0]$.

Unless doing so would introduce an ambiguity, we will tend to reduce notational clutter by dropping the superscripts of these elementary operators.

Observation 4 (the simplicial identities). The following classical relationships hold in Δ_+ and are sufficient to fully characterise equalities between composites of elementary face and degeneracy operators in Δ_+ :

- for any pair $j < i \in [n+1]$ we have $\delta_i^{n+1} \circ \delta_j^n = \delta_j^{n+1} \circ \delta_{i-1}^n$, and

- for any pair $j \leq i \in [n-1]$ we have $\sigma_i^{n-1} \circ \sigma_j^n = \sigma_j^{n-1} \circ \sigma_{i+1}^n$.
- for all $j \in [n]$ and $i \in [n-1]$ we have

$$\sigma_i^{n-1} \circ \delta_j^n = \begin{cases} \delta_j^{n-1} \circ \sigma_{i-1}^{n-2} & \text{if } j < i, \\ \text{id}_{[n-1]} & \text{if } j = i \text{ or } j = i+1, \\ \delta_{j-1}^{n-1} \circ \sigma_i^{n-2} & \text{if } j > i+1. \end{cases}$$

Notation 5 (partition operators). We say that a pair $p, q \in \mathbb{N}$ is a *partition* of $n \in \mathbb{N}$ if $p + q = n$. For each such partition we have:

- face operators $\mathbb{I}_1^{p,q}: [p] \longrightarrow [n]$ given by $\mathbb{I}_1^{p,q}(i) = i$ and $\mathbb{I}_2^{p,q}: [q] \longrightarrow [n]$ given by $\mathbb{I}_2^{p,q}(j) = j + p$, and
- degeneracy operators $\pi_1^{p,q}: [n] \longrightarrow [p]$ given by

$$\pi_1^{p,q}(i) = \begin{cases} i & \text{when } i \leq p \text{ and} \\ p & \text{when } i > p. \end{cases}$$

and $\pi_2^{p,q}: [n] \longrightarrow [q]$ given by:

$$\pi_2^{p,q}(i) = \begin{cases} 0 & \text{when } i < p \text{ and} \\ i - p & \text{when } i \geq p. \end{cases}$$

We call these *partition operators* and, as is easily verified, they satisfy the following *partition identities*:

$$\begin{aligned} \pi_1^{p,q} \circ \mathbb{I}_1^{p,q} &= \text{id}_{[p]} & \pi_2^{p,q} \circ \mathbb{I}_2^{p,q} &= \text{id}_{[q]} \\ \mathbb{I}_1^{p+q,r} \circ \mathbb{I}_1^{p,q} &= \mathbb{I}_1^{p,q+r} & \mathbb{I}_1^{p+q,r} \circ \mathbb{I}_2^{p,q} &= \mathbb{I}_2^{p,q+r} \circ \mathbb{I}_1^{q,r} & \mathbb{I}_2^{p,q+r} \circ \mathbb{I}_2^{q,r} &= \mathbb{I}_2^{p+q,r} \\ \pi_1^{p,q} \circ \pi_1^{p+q,r} &= \pi_1^{p,q+r} & \pi_2^{p,q} \circ \pi_1^{p+q,r} &= \pi_1^{q,r} \circ \pi_2^{p,q+r} & \pi_2^{q,r} \circ \pi_2^{p,q+r} &= \pi_2^{p+q,r} \end{aligned} \quad (1)$$

Observation 6 (duals of simplicial operators). There exists a canonical functor $(-)^{\circ}$ from Δ_+ to itself which “maps each ordinal to its dual as an ordered set”. Explicitly, $[n]^{\circ} = [n]$ and if $\alpha: [n] \longrightarrow [m]$ is a simplicial operator then, for each $i \in [n]$, $\alpha^{\circ}(i) = m - \alpha(n - i)$. Clearly this dual functor is *strictly involutive* in the sense that the diagram

$$\begin{array}{ccc} \Delta_+ & \xrightarrow{\text{id}_{\Delta_+}} & \Delta_+ \\ & \parallel & \\ & (-)^{\circ} & \\ & \Delta_+ & \end{array}$$

of functors commutes (on the nose). In other words, the functor obtained by composing $(-)^{\circ}$ with itself is the identity on Δ_+ . Notice also that the identities

$$\begin{aligned} (\delta_i^n)^{\circ} &= \delta_{n-i}^n & (\sigma_i^n)^{\circ} &= \sigma_{n-i}^n & (\varepsilon_i^n)^{\circ} &= \varepsilon_{n-i}^n \\ (\mathbb{I}_1^{p,q})^{\circ} &= \mathbb{I}_2^{q,p} & (\mathbb{I}_2^{p,q})^{\circ} &= \mathbb{I}_1^{q,p} & (\pi_1^{p,q})^{\circ} &= \pi_2^{q,p} & (\pi_2^{p,q})^{\circ} &= \pi_1^{q,p} \end{aligned} \quad (2)$$

hold between elementary operators and their duals.

Observation 7 (a useful characterisation of face and degeneracy operators). The following facts about simplicial operators $\alpha: [n] \longrightarrow [m]$ are sometimes of use:

- α is a degeneracy operator iff $\alpha(0) = 0$, $\alpha(n) = m$ and for all $j \leq j' \in [n]$ we have $\alpha(j') - \alpha(j) \leq j' - j$.
- α is a face operator iff for all $j \leq j' \in [n]$ we have $\alpha(j') - \alpha(j) \geq j' - j$.

- (iii) α is a face operator iff there exists a simplicial operator $\tilde{\alpha}: [n] \longrightarrow [m-n]$ such that $\alpha(j) = j + \tilde{\alpha}(j)$ for all $j \in [n]$.

Observation 8 (face-degeneracy factorisation). Every simplicial operator $\alpha: [n] \longrightarrow [m]$ factors **uniquely** into a composite $\alpha^f \circ \alpha^d$ where α^f is a face operator and α^d is a degeneracy. Furthermore, let $\text{im}(\alpha)$ denote the subset of $[m]$ given by:

$$\text{im}(\alpha) = \{i \in [m] \mid (\exists j \in [n]) \alpha(j) = i\}$$

It is a trivial, but nonetheless useful, fact that one simplicial operator $\alpha: [m] \longrightarrow [n]$ factors though another $\beta: [r] \longrightarrow [n]$, that is to say there is some $\gamma: [m] \longrightarrow [r]$ with $\beta \circ \gamma = \alpha$, if and only if $\text{im}(\alpha) \subseteq \text{im}(\beta)$.

2.2. The Algebraist's Δ and 2-Categories. We quickly review the theory of 2-categories, which will be useful for expressing some of the “meta-theory” developed in the remainder of this section. Later on, in section 4, we will take a second bite at the 2-category cherry and consider them “in the small” when we review the algebraic theory of ω -categories.

Observation 9 (the cartesian closed category $\underline{Cat}$). We will let $\underline{Cat}$ denote the (huge) category of all (large) categories and functors between them.

This category is cartesian closed, where the “function space” from a category $\mathcal{C}$ to a category $\mathcal{D}$ is the *functor category* $[\mathcal{C}, \mathcal{D}]$ which has:

- **Objects** functors from $\mathcal{C}$ to $\mathcal{D}$,
- **Maps** natural transformations between such functors,
- **Composition** the usual, point-wise, composition of natural transformations.

By definition, this function space is characterised by the adjunction

$$\underline{Cat} \begin{array}{c} \xleftarrow[- \times \mathcal{C}]{} \\ \xrightarrow[\perp]{} \\ \xrightarrow{[\mathcal{C}, *]} \end{array} \underline{Cat}$$

where $\times$ denotes the cartesian product of categories. In other words, there exists a natural bijection between functors $F: \mathcal{B} \times \mathcal{C} \longrightarrow \mathcal{D}$ and $\hat{F}: \mathcal{B} \longrightarrow [\mathcal{C}, \mathcal{D}]$. It follows (see [19]) that $\underline{Cat}$ is a rich “universe” over which we might enrich the homsets of other categories. This observation leads to the following definition:

Definition 10 (2-category). A *2-category* is a category enriched in the cartesian closed category $\underline{Cat}$. Correspondingly, a *2-functor* is simply an $\underline{Cat}$ -enriched functor between 2-categories.

The canonical reference for the theory of enriched categories is, of course, Kelly’s comprehensive book [19] - to which we recommend the reader.

Observation 11 (2-categories explicitly). Fundamentally, a 2-category is simply a category $\underline{\mathcal{C}}$ in which each homset $\underline{\mathcal{C}}(C, D)$ is itself a category (object of $\underline{Cat}$). This immediately implies that our 2-category contains 3 sorts of objects:

- **0-cells** which are the objects of $\underline{\mathcal{C}}$ (for which we use symbols $A, B, C, \dots$),
- **1-cells** which are the objects of the homsets $\underline{\mathcal{C}}(A, B)$ (for which we use symbols $F, G, H, \dots$),
- **2-cells** which are the arrows of the homsets $\underline{\mathcal{C}}(A, B)$ (for which we use Greek symbols $\lambda, \mu, \nu, \dots$).

Commensurate with the two layers of “category-ness” involved in $\underline{\mathcal{C}}$ we get two distinct category structures on these cells:


- **horizontal** with objects which are 0-cells, arrows which are 1-cells **and** 2-cells and compositional structure given by the quadruple $(\circ, \text{dom}_0, \text{cod}_0, \text{id}_0)$. Often we write this composition simply as juxtaposition.
- **vertical** with objects which are 1-cells, arrows which are 2-cells and compositional structure given by $(\cdot, \text{dom}_1, \text{cod}_1, \text{id}_1)$.

In diagrams and running text we will tend to use single arrows $F: A \longrightarrow B$ to denote 1-cells (with $A = \text{dom}_0(F)$ and $B = \text{cod}_0(F)$) and double arrows $\lambda: F \Longrightarrow G$ to denote 2-cells (with $F = \text{dom}_1(\lambda)$ and $G = \text{cod}_1(\lambda)$).

These category structures must satisfy a number of compatibility conditions which bind them together intimately. Most important amongst these are:

- **globularity:** for all 2-cells $\lambda \in \underline{\mathcal{C}}$ we have $\text{dom}_0(\text{dom}_1(\lambda)) = \text{dom}_0(\text{cod}_1(\lambda))$ and $\text{cod}_0(\text{dom}_1(\lambda)) = \text{cod}_0(\text{cod}_1(\lambda))$.
- **middle four interchange:** if we have 2-cells λ, λ', μ and μ' then the equality $(\mu' \circ \lambda') \cdot (\mu \circ \lambda) = (\mu' \cdot \mu) \circ (\lambda' \cdot \lambda)$ holds whenever the various composites involved are defined. Notice that the composite on the left of this equality is defined iff that on its right is defined.

The globularity condition on a 2-cell λ implies that we may picture it as follows:



where $F = \text{dom}_1(\lambda)$, $G = \text{cod}_1(\lambda)$, $A = \text{dom}_0(\lambda)$ and $B = \text{cod}_0(\lambda)$.

Finally, a 2-functor $\mathcal{F}: \underline{\mathcal{C}} \longrightarrow \underline{\mathcal{D}}$ may be thought of explicitly as a map which maps the cells of $\underline{\mathcal{C}}$ to those of $\underline{\mathcal{D}}$ in a way which acts functorially on both horizontal *and* vertical category structures

Observation 12 (whiskering). We commonly identify 0-cells with their corresponding identity 1-cells (under id_0), and 1-cells in turn with their identity 2-cells (under id_1). For instance, we will often use this convention, and the one that allows us to replace $\circ$ by juxtaposition, to write things like

- $G\lambda$ for $(\text{id}_1(G)) \circ \lambda$ and
- μF for $\mu \circ (\text{id}_1(F))$.

These two derived operations are so useful that they have been respectively dubbed left and right *whiskering* by Street [35]. Indeed it is possible (and quite informative) to re-cast the theory of 2-categories purely in terms of these whiskering operations and vertical composition alone. That this is the case follows directly from the observation that the identity and middle four interchange rules imply that for horizontally composable 2-cells $\lambda: F \Longrightarrow F'$ and $\mu: G \Longrightarrow G'$ we have

$$(G'\lambda) \cdot (\mu F) = \mu \circ \lambda = (\mu F') \cdot (G\lambda)$$

Observation 13 ($\underline{\mathcal{Cat}}$ as a 2-category). We know, from Kelly [19], that we may immediately enrich the cartesian closed category $\underline{\mathcal{Cat}}$ over itself, by making $[\mathcal{B}, \mathcal{C}]$ the enriched homset between categories $\mathcal{B}$ and $\mathcal{C}$. This 2-category structure is often best described explicitly in terms of whiskering operations, it has:

- **0-cells** (large) categories, **1-cells** functors between these and **2-cells** natural transformations between those.
- **vertical** composition of 2-cells given by the usual “point-wise” composite of natural transformations.
- **left whiskering** of a natural transformation $\lambda: F \Longrightarrow F': \mathcal{B} \longrightarrow \mathcal{C}$ by a functor $G: \mathcal{C} \longrightarrow \mathcal{D}$ is formed by applying G “point-wise” to the components of λ , that is $(G\lambda)_b \stackrel{\text{def}}{=} G(\lambda_b)$ for each $b \in \text{obj}(\mathcal{B})$.
- **right whiskering** of a natural transformation $\mu: G \Longrightarrow G': \mathcal{C} \longrightarrow \mathcal{D}$ by a functor $F: \mathcal{B} \longrightarrow \mathcal{C}$ is obtained by re-indexing the components of μ using the action of the functor F on objects, that is $(\mu F)_b \stackrel{\text{def}}{=} \mu_{F(b)}$ for each $b \in \text{obj}(\mathcal{B})$.

Observation 14 (Δ_+ as a 2-category). Since each object of Δ_+ is an ordered set it follows that each of its homsets $\Delta_+([n], [m])$ possesses a natural partial order $\leq$, given by

$$\alpha \leq \beta \text{ if and only if } (\forall j \in [n])(\alpha(j) \leq \beta(j))$$

Furthermore, since each map in Δ_+ is order preserving, we also know that composition preserves these partial orders, in the sense that if $\alpha \leq \beta$ in $\Delta_+([n], [m])$ and $\alpha' \leq \beta'$ in $\Delta_+([m], [r])$ then $(\alpha' \circ \alpha) \leq (\beta' \circ \beta)$ in $\Delta_+([n], [r])$.

It follows that, under these partial orders on its homsets, Δ_+ becomes a partial order enriched category or, in other words, a 2-category each homset of which is a partial order.

In fact, this is simply the full sub-2-category of $\underline{\text{Cat}}$ on those 0-cells obtained by considering each totally ordered set $[n]$ ($n \in \mathbb{N}$) as a category in the usual way. That is to say, think of $[n]$ as a category with objects $\{0, 1, \dots, n\}$ and a unique arrow $i \longrightarrow j$ for each pair of integers with $i \leq j$.

It is worth noting that the following useful inequalities (2-cells) hold between (composites of) face and degeneracy operators:

- for all $n \geq 1$ and $j \in [n-1]$ we have $\delta_j^n \circ \sigma_j^{n-1} > \text{id}_{[n]}$ and $\delta_{j+1}^n \circ \sigma_j^{n-1} < \text{id}_{[n]}$,
- if $n \geq 1$ and $j \leq k \in [n]$ then $\delta_k^n \leq \delta_j^n$ and $\sigma_j^{n-1} \leq \sigma_k^{n-1}$.

Observation 15 (adjoints in the 2-category Δ_+). In the sequel we will have occasion to consider adjoint pairs $\alpha \dashv \beta$ of simplicial operators. These can be defined and understood abstractly in the 2-category Δ_+ , in terms of the general theory of adjoint pairs in 2-categories as expounded (for instance) by Kelly and Street in their classic review article [20] or by Street in [30].

However, here we prefer to directly exploit the identification of the 2-category Δ_+ as a full sub-2-category of $\underline{\text{Cat}}$ and utilise the traditional definition of adjunction. In such terms, a simplicial operator $\alpha: [n] \longrightarrow [m]$ is left adjoint to $\beta: [m] \longrightarrow [n]$ if for all $i \in [n]$ and $j \in [m]$ we have $\alpha(i) \leq j$ iff $i \leq \beta(j)$. Equivalently, this condition holds if the inequalities $\text{id}_{[n]} \leq \beta \circ \alpha$ and $\alpha \circ \beta \leq \text{id}_{[m]}$ hold.

Of course, the usual properties of adjunctions hold for those in Δ_+ , in particular the left (resp. right) adjoint to an operator α (if such a thing exists) is unique. Furthermore, the following classical observations characterise adjunctions of simplicial operators:

- α has a (necessarily unique) *left* adjoint $\alpha^l: [m] \longrightarrow [n]$ iff $\alpha(n) = m$, in which case $\alpha^l(i) = \min\{j \in [n] \mid i \leq \alpha(j)\}$.
- Dually, α has a (necessarily unique) *right* adjoint $\alpha^r: [m] \longrightarrow [n]$ iff $\alpha(0) = 0$, in which case $\alpha^r(i) = \max\{j \in [n] \mid i \geq \alpha(j)\}$.
- It follows that if α is a degeneracy operator then it has both left and right adjoints.

- (iv) If $\alpha \dashv \beta$ is an adjoint pair and either α is a degeneracy operator or β is a face operator then $\alpha \circ \beta = \text{id}_{[m]}$.
- (v) Dually, if $\alpha \dashv \beta$ is an adjoint pair and either α is a face operator or β is a degeneracy operator then $\beta \circ \alpha = \text{id}_{[n]}$.
- (vi) The simplicial identities and inequalities of observation 4 demonstrate that σ_j^{n-1} has **right adjoint** δ_j^n and **left adjoint** δ_{j+1}^n .

Finally, notice that the dual functor $(-)^{\circ}$ may be naturally extended to a 2-functor which is **contravariant on 2-cells**. By this we mean to say that if α and β are simplicial operators with $\alpha \leq \beta$ then $\beta^{\circ} \leq \alpha^{\circ}$. It follows that $(-)^{\circ}$ carries left (resp. right) adjoints to right (resp. left) adjoints. In other words, $(\alpha^{\circ})^r = (\alpha^l)^{\circ}$ and $(\alpha^{\circ})^l = (\alpha^r)^{\circ}$ when these adjoints exist.

2.3. The Algebraist's Δ and Monoidal Categories. In this subsection, we recall a few standard results regarding the algebraic content of Δ_+ . For proofs of these results, and the details of many other interesting facets of the algebra of Δ_+ , we refer the reader to Mac Lane [22].

Observation 16 (Δ_+ by generators and relations). We can go further than we did in the last subsection and factor every face (resp. degeneracy) operator as a (canonical) composite of elementary face (resp. degeneracy) operators. It follows that the elementary operators generate Δ_+ , but more is true.

We said above that the simplicial identities served to fully characterise the compositional relationships between elementary operators. This can be made precise by observing that Δ_+ may be *presented* in terms of the elementary operators as its generators and the simplicial identities as its relations.

In other words, in order to define a functor $F: \Delta_+ \longrightarrow \mathcal{E}$ it suffices to specify its action on objects and elementary operators and then check that the images of these elementary operators satisfy the simplicial identities in $\mathcal{E}$. It also implies that we may test the naturality of a family of maps between two functors $F, F': \Delta_+ \longrightarrow \mathcal{E}$ by testing it with respect to the elementary operators alone.

Recall 17 (monoidal categories). Recall (from say Mac Lane [22]) that a *monoidal category* $(\mathcal{E}, \otimes, I)$ consists of a category $\mathcal{E}$ equipped with a bifunctor $\otimes: \mathcal{E} \times \mathcal{E} \longrightarrow \mathcal{E}$ called a *tensor product*, an object $I \in \mathcal{E}$, and natural isomorphisms:

$$\begin{array}{llll}
 X \otimes (Y \otimes Z) & \xrightarrow{\alpha_{X,Y,Z}} & (X \otimes Y) \otimes Z & \text{associativity} \\
 X \otimes I & \xrightarrow{\iota_X^r} & X & \text{right identity} \\
 I \otimes X & \xrightarrow{\iota_X^l} & X & \text{left identity}
 \end{array}$$

These isomorphisms must also satisfy a number of *coherence conditions*, two of which relate the identity isomorphisms to the associativity and one, called the *pentagon condition*, which relates the various associativities involved in a four fold tensor product (see Mac Lane [22] for greater detail).

A *monoidal functor* $F: (\mathcal{E}, \otimes, I) \longrightarrow (\mathcal{E}', \otimes', I')$ consists of a functor $F: \mathcal{E} \longrightarrow \mathcal{E}'$ of underlying categories and a natural family of morphisms:

$$\begin{array}{lll}
 F(X) \otimes' F(Y) & \xrightarrow{\mu_{X,Y}} & F(X \otimes Y) \quad \text{product comparison} \\
 I' & \xrightarrow{\nu} & F(I) \quad \text{identity comparison}
 \end{array}$$

These morphisms must also satisfy coherence conditions, the first relating the identity comparison to right and left identity isomorphisms and the second relating the product comparisons to associativity isomorphisms (again see Mac Lane [22] for greater detail). We say that a monoidal functor is *strong* (resp. *strict*) if each of these comparison maps is in fact an isomorphism (resp. an identity) in $\mathcal{E}'$.

A *monoidal natural transformation* $\eta: F \Longrightarrow F'$ between monoidal functors F and F' is simply a natural transformation the components of which commute with the product and identity comparisons on its domain and codomain. In other words, we must have $\eta_I \cdot \nu = \nu'$ and $\eta_{X \otimes Y} \cdot \mu_{X,Y} = \mu'_{X,Y} \cdot (\eta_X \otimes \eta_Y)$.

Notice that, we may extend the usual composites of functors and natural transformations to provide corresponding composites of monoidal functors and monoidal natural transformations. To do this all we need worry about is how to define the comparison maps for the composite of a pair of monoidal functors $F: (\mathcal{E}, \otimes, I) \longrightarrow (\mathcal{E}', \otimes', I')$ and $F': (\mathcal{E}', \otimes', I') \longrightarrow (\mathcal{E}'', \otimes'', I'')$. However, a moment's reflection reveals that the most natural candidates for these are the composites

$$F'(F(X)) \otimes'' F'(F(Y)) \xrightarrow{\mu'_{F(X), F(Y)}} F'(F(X) \otimes' F(Y)) \xrightarrow{F'(\mu_{X,Y})} F'(F(X \otimes Y))$$

for tensor products and

$$I'' \xrightarrow{\nu'} F'(I') \xrightarrow{F'(\nu)} F'(F(I))$$

for identities. Verifying the various coherence conditions for this structure, as well as checking that it is well behaved with respect to horizontal composition of monoidal natural transformations, is a matter of routine calculation. In this way we lift the structure of the 2-category Cat (of categories, functors and natural transformations) to get a 2-category Mon with 0-cells monoidal categories, 1-cells monoidal functors between them and 2-cells that are monoidal natural transformations between those.

Recall 18 (monoids). As the name implies, monoidal categories bear a strong relationship to *monoids*, that is algebraic structures $(X, *, e)$ where $*$ is an associative binary operation on X for which $e \in X$ is a two sided identity. This relationship finds its clearest expression in two interesting observations:

External: Monoidal categories are a convenient structure within which to interpret the monoid concept. If $(\mathcal{E}, \otimes, I)$ is a monoidal category then a monoid $(X, *, e)$ in $\mathcal{E}$ consists of an underlying object $X \in \mathcal{E}$, a *multiplication* map $*: X \otimes X \longrightarrow X$ and a *unit* map $e: I \longrightarrow X$. This data must satisfy the diagrammatic conditions

$$\begin{array}{ccccc} X \otimes (X \otimes X) & \xrightarrow[\cong]{\alpha_{X,X,X}} & (X \otimes X) \otimes X & \xrightarrow{* \otimes X} & X \otimes X \\ X \otimes * \downarrow & & & & \downarrow * \\ X \otimes X & \xrightarrow{\quad \quad \quad} & & & X \\ & & * & & \\ X \otimes I & \xrightarrow{X \otimes e} & X \otimes X & \xleftarrow{e \otimes X} & I \otimes X \\ & \searrow \cong & \downarrow * & \swarrow \cong & \\ & \iota_X^r & X & \iota_X^l & \end{array}$$

which are simply the usual associativity and identity conditions for a monoid in disguise.

Many common algebraic structures turn out to be monoids in a suitable monoidal category. For instance, rings are really no more than monoids within the category of abelian groups equipped with the usual tensor product of groups as its monoidal structure.

We form a category $\underline{\text{Mon}}(\mathcal{E}, \otimes, I)$ of monoids in $(\mathcal{E}, \otimes, I)$, in which a *monoid morphism* f from $(X, *, e)$ to $(X', *, e')$ is simply a map of the underlying objects $f: X \longrightarrow X'$ in $\mathcal{E}$ which commutes with the monoid structures on its domain and codomain, or in other words for which $f \circ e = e'$ and $f \circ * = *' \circ (f \otimes f)$.

Internal: Monoidal categories in which each of the structural isomorphisms $\alpha_{X,Y,Z}$, ι_X^r and ι_X^l are actually *identities* are said to be *strict*. In other words, a strict monoidal category $(\mathcal{C}, \otimes, I)$ consists of a category equipped with a tensor structure for which the usual associativity and identity conditions $\lambda \otimes (\mu \otimes \nu) = (\lambda \otimes \mu) \otimes \nu$, $\lambda \otimes \text{id}_I = \lambda$ and $\text{id}_I \otimes \lambda = \lambda$ hold “on the nose” for all arrows $\lambda, \mu, \nu \in \mathcal{C}$. Equivalently, such a structure is no more nor less than a monoid in the (huge) monoidal category $(\underline{\text{Cat}}, \times, 1)$.

Notice that, if $\underline{\mathcal{C}}$ is a 2-category then each endo-category $\underline{\mathcal{C}}(A, A)$ on a 0-cell $A \in \underline{\mathcal{C}}$ is the underlying category of a strict monoidal category, with tensor given by horizontal composition $\circ$ and identity $\text{id}_0(A)$. Conversely, each strict monoidal category $(\mathcal{E}, \otimes, I)$ gives rise to a 2-category $\Omega\mathcal{E}$, called its *(one-point) suspension*, with a single 0-cell $*$, hom-category $\Omega\mathcal{E}(*, *) = \mathcal{E}$, identity $\text{id}_0(*) = I$, and horizontal composition $\otimes$.

Observation 19 (relating monoids and monoidal functors). It is worth observing that monoids and monoid maps bear a close relationship to monoidal functors and monoidal natural transformation. To be precise, it is clear that the one object, one arrow “terminal” category $\mathcal{I}$ admits a unique strict monoidal structure, with respect to which we may consider monoidal functors $G: \mathcal{I} \longrightarrow (\mathcal{E}, \otimes, I)$. On examining the data for such monoidal functor, it becomes clear that it amounts to no more nor less than an object $G(1)$, an identity comparison $\nu: I \longrightarrow G(1)$ and a product comparison $\mu_{1,1}: G(1) \otimes G(1) \longrightarrow G(1)$ satisfying coherence conditions that are identical to the axioms required of a monoid $(G(1), \mu_{1,1}, \nu)$ in $\mathcal{E}$. Furthermore, the data and conditions for a monoidal natural transformation $\eta: G \Longrightarrow G'$ correspond exactly to those for a monoid map between the monoids corresponding to G and G' . In other words, the category $\underline{\text{Mon}}(\mathcal{E}, \otimes, I)$ and the hom-category $\underline{\text{Mon}}(\mathcal{I}, (\mathcal{E}, \otimes, I))$ are canonically equivalent.

This observation has many useful consequences, for instance it implies that a monoidal functor $F: (\mathcal{E}, \otimes, I) \longrightarrow (\mathcal{E}', \otimes', I')$ may be lifted to a functor between categories of monoids $\underline{\text{Mon}}(F): \underline{\text{Mon}}(\mathcal{E}, \otimes, I) \longrightarrow \underline{\text{Mon}}(\mathcal{E}', \otimes', I')$. To be precise, this is simply the post composition functor $\underline{\text{Mon}}(\mathcal{I}, F)$ between corresponding hom-categories of the 2-category $\underline{\text{Mon}}$.

Observation 20 (Δ_+ as a strict monoidal category). In order to see why Δ_+ might be of use to Algebraists, it is first necessary to observe that ordinal addition of its objects extends to a functor $\oplus: \Delta_+ \times \Delta_+ \longrightarrow \Delta_+$ where $[n] \oplus [m] \stackrel{\text{def}}{=} [n + m + 1]$ and the *direct sum* $\beta \oplus \alpha$ of operators $\alpha: [n] \longrightarrow [n']$ and $\beta: [m] \longrightarrow [m']$ is defined by

$$\beta \oplus \alpha(j) = \begin{cases} \alpha(j) & \text{if } j \leq n, \\ n' + 1 + \beta(j - n - 1) & \text{if } j > n. \end{cases}$$

It is easily seen that the triple $(\Delta_+, \oplus, [-1])$ satisfies the strict associativity and identity conditions discussed in recollection 18 and thus it is a strict monoidal category. Notice also

that direct sums interact gracefully with the involution $(-)^{\circ}: \Delta \longrightarrow \Delta$, in particular if α and β are simplicial operators then we have $(\alpha \oplus \beta)^{\circ} = \beta^{\circ} \oplus \alpha^{\circ}$.

Observation 21 (generators for Δ_+ as a strict monoidal category). The strict monoidal category $(\Delta_+, \oplus, [-1])$ contains a canonical monoid with underlying object $[0]$, identity $\delta_0^0: [-1] \longrightarrow [0]$ and multiplication $\sigma_0^0: [0] \oplus [0] = [1] \longrightarrow [0]$, for which the associativity and unit conditions may be trivially verified.

Notice that all elementary face and degeneracy operators may be obtained as iterated direct sums of the structural components of the monoid $([0], \sigma_0^0, \delta_0^0)$ as follows:

- each object $[n]$ may be obtained from the underlying object $[0]$ as an iterated direct sum $[0]^{\oplus(n+1)}$,
- each elementary face operator δ_i^n may be obtained from the unit map δ_0^0 as an iterated direct sum $[0]^{\oplus(n-i)} \oplus \delta_0^0 \oplus [0]^{\oplus i}$, and
- each elementary degeneracy operator σ_i^n may be obtained from the multiplication map σ_0^0 as an iterated direct sum $[0]^{\oplus(n-i)} \oplus \sigma_0^0 \oplus [0]^{\oplus i}$.

Of course we know, by observation 16, that every simplicial operator may be expressed as a composite of elementary face and degeneracy operators and it follows that every operator may be obtained as a composite of iterated direct sums of the structural components of $([0], \sigma_0^0, \delta_0^0)$.

In fact much more is true for this monoid, as demonstrated by the following classical result which states that it actually *freely* generates $(\Delta_+, \oplus, [-1])$ in some suitable sense made precise in the statement of the lemma.

Lemma 22 (the universal property of Δ_+). *The monoid $([0], \sigma_0^0, \delta_0^0)$ in $(\Delta_+, \oplus, [-1])$ is universal, in the sense that for any other monoid $(X, *, e)$ in a strict monoidal category $(\mathcal{E}, \otimes, I)$ there exists a unique strict monoidal functor $\tilde{X}: (\Delta_+, \oplus, [-1]) \longrightarrow (\mathcal{E}, \otimes, I)$ such that the monoid*

$$\underline{\text{Mon}}(\tilde{X})([0], \sigma_0^0, \delta_0^0) = (\tilde{X}([0]), \tilde{X}(\sigma_0^0), \tilde{X}(\delta_0^0))$$

*is equal to $(X, *, e)$.*

Proof. Standard, see Mac Lane [22]. It is worth observing, however, that observation 21 immediately implies that the functor $\tilde{X}$ is completely determined by $\tilde{X}([n]) = X^{\otimes(n+1)}$, $\tilde{X}(\delta_i^n) = X^{\otimes(n-i)} \otimes e \otimes X^{\otimes i}$ and $\tilde{X}(\sigma_i^n) = X^{\otimes(n-i)} \otimes * \otimes X^{\otimes i}$. $\square$

2.4. Simplicial Sets.

Definition 23 (simplicial sets and simplicial maps). The category Simp of *simplicial sets* and *simplicial maps* between them is simply the functor category $[\Delta^{\text{op}}, \underline{\text{Set}}]$, where Set denotes the (large) category of all (small) sets and functions between them.

Observation 24 (simplicial sets as partial right actions). In practise, it is easier to think of a simplicial set as a single set endowed with a partially defined right action of the simplicial operators.

To be more precise, this description presents a simplicial set as a triple $(X, \dim, \cdot)$ where:

- X is a (small) set and $\dim: X \longrightarrow \mathbb{N}$ is a function, which assigns to each element of X a *dimension*,
- $x \cdot \alpha \in X$ is defined for any element $x \in X$ and simplicial operator $\alpha: [m] \longrightarrow [n]$ for which $\dim(x) = n$, in which case we say that x and α are *compatible*, and we have $\dim(x \cdot \alpha) = m$, and

- this action satisfies the equations $x \cdot \text{id}_{[n]} = x$ and $(x \cdot \alpha) \cdot \beta = x \cdot (\alpha \circ \beta)$ whenever they are well defined (i.e. when $\alpha: [m] \longrightarrow [n]$, $\beta: [r] \longrightarrow [m]$ and $\dim(x) = n$).

In time honoured fashion, we will usually “overload” $\dim$ and $\cdot$, using them to denote the dimension and action functions of whichever simplicial set we happen to be discussing at any given time. This substantially simplifies our notation and obviates the need to explicitly name these functions when introducing a new simplicial set.

Furthermore, we will often refer to the elements of simplicial set X as its *simplices* and say that $y \in X$ is a *face* of a simplex $x \in X$ if there is some simplicial operator α such that $x \cdot \alpha = y$.

We will often use the notation X_n to denote the set of n -simplices $\{x \in X \mid \dim(x) = n\}$ of X and if $\alpha: [m] \longrightarrow [n]$ is a simplicial operator then we adopt the notation $X_\alpha: X_n \longrightarrow X_m$ for the function which maps $x \in X_n$ to $X_\alpha(x) \stackrel{\text{def}}{=} x \cdot \alpha \in X_m$.

Observation 25 (simplicial maps as action preserving functions). When simplicial sets are expressed in this way it is most natural to consider simplicial maps to be action preserving functions. In other words, a simplicial map $f: (X, \dim, \cdot) \longrightarrow (Y, \dim, \cdot)$ consists of a function $f: X \longrightarrow Y$ of sets satisfying:

- $\dim(f(x)) = \dim(x)$ for each $x \in X$, and
- $f(x \cdot \alpha) = f(x) \cdot \alpha$ for each $x \in X$ and $\alpha \in \Delta$ with $[\dim(x)] = \text{cod}(\alpha)$.

Observation 26 (simplicial subsets). A subset Y of a simplicial set X is called a *simplicial subset*, denoted by $Y \subseteq_s X$, if it is closed in X under the action of Δ . If $Y \subseteq_s X$ then Y becomes a simplicial set, by inheriting the action structure of X , and the inclusion becomes a simplicial map $Y \xrightarrow{\subseteq_s} X$.

Notice that intersections and unions of simplicial subsets of X , as mere subsets, are again closed in X under the action of Δ , in other words they are themselves simplicial subsets of X .

Suppose that A is a subset of a simplicial set X then A^* , its *simplicial closure* in X , is the smallest simplicial subset of X that contains A . Explicitly:

$$A^* = \{x \in X \mid (\exists \alpha \in \Delta, y \in A) \text{ s.t. } x = y \cdot \alpha\}$$

Observation 27 (why *simplicial* sets?). The books by Gabriel and Zisman [13] and May [24] are considered the canonical references for the theory of simplicial sets. In particular, they explain how these structures may be used to provide combinatorial representations of topological structures and expounds at some length upon their homotopy theory.

For our purposes here, it is worth simply observing the following:

- If x is an n -dimensional element of a simplicial set X , then $x \cdot \varepsilon_i^n$ is thought of as its i^{th} vertex and $x \cdot \delta_i^n$ is thought of as its unique $(n-1)$ -dimensional face **not** containing that vertex.
- The simplicial identity for elementary face maps ensures that the various faces of a simplex agree appropriately at their boundaries.

Definition 28 (degenerate simplices). We say that an r -simplex x of a simplicial set X is *degenerate at k* if $x = x' \cdot \sigma_k^{r-1}$ for some $(r-1)$ -simplex x' . Of course σ_k^{r-1} has right inverses δ_k^r and δ_{k+1}^r , therefore such an x' is unique, if it exists, and $x' = x \cdot \delta_k^r = x \cdot \delta_{k+1}^r$.

We also say that a simplex is *degenerate* if it is degenerate at k for some k . Equivalently, $x \in X$ is degenerate iff there is a simplicial operator α and a compatible simplex $x' \in X$ with

$\dim(x') < \dim(x)$ and $x = x' \cdot \alpha$. We often use the notation $\tilde{X}$ to denote the set of those simplices of X which are **not** degenerate.

Observation 29. Let x be an simplex in a simplicial set X , then

- (a) x is degenerate iff there exists some k such that x is degenerate at k .
- (b) x is degenerate at k iff $x = x' \cdot \beta$ for a simplex $x' \in X$ and a simplicial operator β with $\beta(k) = \beta(k+1)$.
- (c) if x is degenerate at k then $x \cdot \alpha$ is degenerate for any face operator α with $k, k+1 \in \text{im}(\alpha)$.

Lemma 30 (the Eilenberg-Zilber lemma). *If x is a simplex of a simplicial set X then there exists a **unique** pair $(\bar{x}, \alpha^x)$ such that $\bar{x} \in X$ is non-degenerate, α^x is a degeneracy operator and $x = \bar{x} \cdot \alpha^x$. We call this unique pair $(\bar{x}, \alpha^x)$ the EZ-decomposition of x .*

Proof. See the proof in [13]. □

Observation 31 (limits and colimits in Simp). As a category of functors into the (small) complete and cocomplete category Set, Simp also has all (small) limits and colimits. These are formed in the usual way for such categories, that is to say *point-wise* in Set.

It is useful to expand upon this definition in terms of the partial action description of simplicial sets given above. To this end, suppose that $\mathbb{C}$ is a small category and that $D: \mathbb{C} \longrightarrow \text{Simp}$ is a functor (which we call a *diagram* of $\mathbb{C}$ in Simp):

Limits The elements of $\lim(D)$ are families $\{x_i \in D(i)\}_{i \in \text{obj}(\mathbb{C})}$ satisfying the conditions that:

- for each pair $i, j \in \text{obj}(\mathbb{C})$ we have $\dim(x_i) = \dim(x_j)$,
- if $f: i \longrightarrow j$ is an arrow in $\mathbb{C}$ then $D(f)(x_i) = x_j$.

The dimension of $\mathbf{x}$ is the common dimension of its components x_i , and the action of an operator α is given by $\{x_i\}_{i \in \text{obj}(\mathbb{C})} \cdot \alpha = \{x_i \cdot \alpha\}_{i \in \text{obj}(\mathbb{C})}$.

Colimits We form the colimit $\text{colim}(D)$ in Set, whose elements are equivalence classes of pairs (i, x) with $i \in \text{obj}(\mathbb{C})$ and $x \in D(i)$ under the equivalence relation $\sim$ generated by the relation $(i, x) \sim_1 (j, y)$ which holds when there exists an arrow $f: i \longrightarrow j$ in $\mathbb{C}$ with $D(f)(x) = y$. Following tradition, we use the notation $[i, x]$ to denote the equivalence class of a pair (i, x) under this equivalence relation.

Notice that if $f: i \longrightarrow j$ witnesses that $(i, x) \sim_1 (j, y)$, then $\dim(y) = \dim(D(f)(x)) = \dim(x)$ and if α is an operator with codomain $[\dim(x)]$ then $y \cdot \alpha = f(x) \cdot \alpha = f(x \cdot \alpha)$. It follows that the operations $\dim([i, x]) = \dim(x)$ and $[i, x] \cdot \alpha = [i, x \cdot \alpha]$ on the equivalence classes of $\sim$ are well defined and make $\text{colim}(D)$ into a simplicial set, the colimit of D in Simp.

Observation 32. The fact that limits and colimits in Simp are constructed point-wise in Set implies that we may immediately “lift” many of their properties from there. For instance the following useful result, which holds in Set and thus immediately lifts to Simp, allows us to describe a union of simplicial subsets as a certain kind of colimit, sometimes called a *wide pushout*, whose vertices are (intersections of) those subsets. To be precise, if Y is a simplicial set and $Y^{(i)} \subseteq Y$ (for $i = 1, 2, \dots, r$) is a family of simplicial subsets with $Y = \bigcup_l Y^{(l)}$ then the cocone of inclusion maps $Y^{(i)} \hookrightarrow Y$ and $Y^{(i)} \cap Y^{(j)} \hookrightarrow Y$ ($i < j$) presents Y as the colimit of a diagram $D_{\{Y^{(i)}\}}$ in Simp consisting of the stratified sets $Y^{(i)}$ and $Y^{(i)} \cap Y^{(j)}$ and

inclusions $Y^{(i)} \cap Y^{(j)} \hookrightarrow Y^{(i)}$ and $Y^{(i)} \cap Y^{(j)} \hookrightarrow Y^{(j)}$ ($i < j$) between them:

$$\begin{array}{ccc}
 & Y^{(i)} \cap Y^{(j)} & \\
 \swarrow & \downarrow & \searrow \\
 Y^{(i)} & & Y^{(j)} \\
 \swarrow & \downarrow & \searrow \\
 & Y &
 \end{array}
 \quad (3)$$

Observation 33 (connected components). For each set $S \in \underline{\text{Set}}$ we have a corresponding *discrete* simplicial set $\text{dis}(S)$ defined by $\text{dis}(S)_n = S$ ($\forall n \in \mathbb{N}$) and $\text{dis}(S)_\alpha = \text{id}_S$ (for all simplicial operators $\alpha \in \Delta$). This construction clearly provides us with a functor $\text{dis}: \underline{\text{Set}} \longrightarrow \underline{\text{Simp}}$ which is easily shown to be fully faithful. It is also clear that a simplicial set is in the replete image of this functor iff each of its simplices of dimension > 0 is degenerate.

We also have an adjunction $\Pi_0 \dashv \text{dis}$ and the set $\Pi_0(X)$ derived from a simplicial set X is known as its *set of connected components*, which may be constructed explicitly using the coequaliser

$$X_1 \begin{array}{c} \xrightarrow{X_{\varepsilon_0^1}} \\ \xrightarrow{X_{\varepsilon_1^1}} \end{array} X_0 \xrightarrow{q_X} \Pi_0(X)$$

in $\underline{\text{Set}}$. In other words, we start with the set of 0-simplices of X and form $\Pi_0(X)$ from it by identifying any pair of 0-simplices which are the 0-faces of some 1-simplex.

Observation 34 (standard simplices in $\underline{\text{Simp}}$). The classical Yoneda functor $\Delta: \Delta \longrightarrow \underline{\text{Simp}}$ carries each object $[n]$ of Δ to the *representable* simplicial set $\Delta[n] \stackrel{\text{def}}{=} \Delta(-, [n])$. This is called the *standard n -simplex* and is given explicitly by:

- $\Delta[n]_m = \Delta([m], [n])$ the set of simplicial operators $\gamma: [m] \longrightarrow [n]$,
- if $\beta: [r] \longrightarrow [m]$ is a simplicial operator that is compatible with $\gamma \in \Delta[n]_m$ then β and γ are composable and we may define $\gamma \cdot \beta = \gamma \circ \beta \in \Delta[n]_r$.

Furthermore, it carries the operator $\alpha: [n] \longrightarrow [m]$ to a simplicial map $\Delta(\alpha): \Delta[n] \longrightarrow \Delta[m]$ given by post-composition $\Delta(\alpha)(\gamma) = \alpha \circ \gamma$.

Given an n -simplex $x \in X$ in a simplicial set X , we adopt the notation $\ulcorner x \urcorner: \Delta[n] \longrightarrow X$ for the simplicial map defined by $\ulcorner x \urcorner(\gamma) \stackrel{\text{def}}{=} x \cdot \gamma$. Of course, Yoneda's lemma applied in the simplicial set context states that the function $\ulcorner \cdot \urcorner: X_n \longrightarrow \underline{\text{Simp}}(\Delta[n], X)$ is inverse to the evaluation function which maps $f \in \underline{\text{Simp}}(\Delta[n], X)$ to the n -simplex $f(\text{id}_{[n]})$ in X .

Observation 35 (a couple of observations regarding $\Delta[n]$). An r -simplex α of $\Delta[n]$ is non-degenerate if and only if, as a simplicial operator $\alpha: [r] \longrightarrow [n]$, it is injective (a face operator). It follows that, if $r > n$ then no such simplicial operator can be injective and so all of the r -simplices of $\Delta[n]$ must be degenerate. Consequently, the only non-degenerate n -simplex of $\Delta[n]$ is the identity $\text{id}_{[n]}: [n] \longrightarrow [n]$.

Notation 36 (the boundaries of standard simplices). The *boundary* $\partial\Delta[n]$ of the standard n -simplex $\Delta[n]$ is the simplicial subset generated by the set of all the $(n-1)$ -faces $\{\delta_i^n \mid i = 0, \dots, n\}$ of $\Delta[n]$. Equivalently, $\partial\Delta[n]$ is the largest simplicial subset of $\Delta[n]$ which does not contain $\text{id}_{[n]}$. A simplex α of $\Delta[n]$ is in $\partial\Delta[n]$ iff it is **not** a degeneracy operator.

We can express $\partial\Delta[n]$ as the union $\bigcup_{i=0}^n \{\delta_i^n\}^*$ of the simplicial subsets $\{\delta_i^n\}^* \subseteq \partial\Delta[n]$ which are the images of the simplicial injections $\Delta(\delta_i^n): \Delta[n-1] \hookrightarrow \Delta[n]$ and are thus isomorphic

to $\Delta[n-1]$. Also, if $i < j$ then we have $\alpha \in \{\delta_i^n\}^* \cap \{\delta_j^n\}^*$ iff α factors through both δ_i^n and δ_j^n iff $i, j \notin \text{im}(\alpha)$ iff $\alpha \in \{\delta_j^n \circ \delta_i^{n-1}\}^*$ therefore $\{\delta_i^n\}^* \cap \{\delta_j^n\}^* = \{\delta_j^n \circ \delta_i^{n-1}\}^*$ which is the image of the simplicial injection $\Delta(\delta_j^n \circ \delta_i^{n-1}) = \Delta(\delta_i^n \circ \delta_{j-1}^{n-1}): \Delta[n-2] \hookrightarrow \Delta[n]$ and is thus isomorphic to $\Delta[n-2]$. Furthermore, it is also clear that under these isomorphisms the inclusions of $\{\delta_i^n\}^* \cap \{\delta_j^n\}^*$ into $\{\delta_i^n\}^*$ and $\{\delta_j^n\}^*$ correspond to the simplicial injections $\Delta(\delta_{j-1}^{n-1}): \Delta[n-2] \hookrightarrow \Delta[n-1]$ and $\Delta(\delta_i^{n-1}): \Delta[n-2] \hookrightarrow \Delta[n-2]$ respectively.

Applying observation 32 and the isomorphisms of the last paragraph we see that $\partial\Delta[n]$ may be expressed as the wide pushout

$$\begin{array}{ccc}
 & \Delta[n-2] & \\
 \Delta(\delta_{j-1}^{n-1}) \swarrow & \downarrow & \searrow \Delta(\delta_i^{n-1}) \\
 \Delta[n-1] & \partial\Delta[n] & \Delta[n-1] \\
 \Delta(\delta_i^n) \swarrow & & \searrow \Delta(\delta_j^n)
 \end{array}$$

of standard simplices. Applying Yoneda's lemma, and the colimiting property of this wide pushout, we see that a simplicial map $f: \partial\Delta[n] \longrightarrow X$ corresponds to a family of $(n-1)$ -simplices $x_0, x_1, \dots, x_n \in X_{n-1}$ satisfying the simplicial identities $x_i \cdot \delta_j^{n-1} = x_j \cdot \delta_{i-1}^{n-1}$ for $j < i$. This data is often referred to as a $(n-1)$ -dimensional cycle in X and we say that such a cycle is the *boundary* of an n -simplex $x \in X$ if and only if $x_i = x \cdot \delta_i^n$ for each $i = 0, 1, \dots, n$.

Notation 37 (horns). For each $n \in \mathbb{N}$ and $k = 0, \dots, n$, the simplicial set $\Lambda_k[n]$, called the *standard $(n-1)$ -dimensional k -horn*, is the smallest simplicial subset of $\Delta[n]$ containing the set of $(n-1)$ -simplices $\{\delta_i^n \mid i = 0, \dots, k-1, k+1, \dots, n\}$. We say that such a horn is an *inner horn* whenever $0 < k < n$ otherwise we say that it is an *outer horn*.

Arguing as in notation 36, we see that a simplicial map $f: \Lambda_k[n] \longrightarrow X$ corresponds to a family of $(n-1)$ -simplices $x_0, x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n \in X_{n-1}$ satisfying the simplicial identities $x_i \cdot \delta_j^{n-1} = x_j \cdot \delta_{i-1}^{n-1}$ for $j < i$. This data is often referred to as a $(n-1)$ -dimensional k -horn in X . We say that an n -simplex $x \in X$ *fills* such a horn if and only if $x_i = x \cdot \delta_i^n$ for each $i = 0, \dots, k-1, k+1, \dots, n$.

Observation 38 (the dual of a simplicial set). The *dual* X° of a simplicial set X is obtained by pre-composing with the functor $(-)^{\circ}: \Delta \longrightarrow \Delta$. In action terms, X° has the same set of simplices as X but an action $*$ given by $x * \alpha = x \cdot \alpha^\circ$.

Notice that the dualising functor $(-)^{\circ}: \mathbf{Simp} \longrightarrow \mathbf{Simp}$ is again strictly involutive, in the sense that its composite with itself is the identity on $\mathbf{Simp}$. This is an immediate consequence of the fact that $(-)^{\circ}$ is strictly involutive on Δ .

2.5. Semi-Simplicial Sets.

Definition 39 (semi-simplicial sets). Let Δ_f (resp. Δ_{f+}) denote the subcategory of face operators in Δ (resp. Δ_+). The category $\mathbf{SSimp}$ of *semi-simplicial sets* and *semi-simplicial maps* between them is simply the functor category $[\Delta_f^{\text{op}}, \mathbf{Set}]$.

Notation 40 (the category of pointed objects). We may also characterise the monoidal category of face operators $(\Delta_{f+}, \oplus, [-1])$ in the spirit of lemma 22. However, to do so, we must first introduce the simple notion of a *pointed object* (X, e) in a monoidal category $(\mathcal{E}, \otimes, I)$, which consists of an object $X \in \mathcal{E}$ equipped with a map $e: I \longrightarrow X$ called a

point. A map of pointed objects $f: (X, e) \longrightarrow (X', e')$ is simply a map of underlying objects $f: X \longrightarrow X'$ which “preserves the point” in the sense that $f \circ e = e'$; in other words the category of pointed objects $\underline{\mathbf{Pt}}(\mathcal{E}, \otimes, I)$ is simply the comma category $I \downarrow \mathcal{E}$.

If $F: (\mathcal{E}, \otimes, I) \longrightarrow (\mathcal{E}', \otimes', I')$ is a monoidal functor then we may define a functor $\underline{\mathbf{Pt}}(F)$ between corresponding categories of points by mapping a pointed object (X, e) in $(\mathcal{E}, \otimes, I)$ to $(F(X), F(e) \circ \nu)$ in $(\mathcal{E}', \otimes', I')$ and a point preserving map $f: (X, e) \longrightarrow (X', e')$ to $F(f)$.

Lemma 41 (the universal property of Δ_{f+}). *The pointed object $([0], \delta_0^0)$ in $(\Delta_{f+}, \oplus, [-1])$ is universal, in the sense that if (X, e) is a pointed object in a monoidal category $(\mathcal{E}, \otimes, I)$ then there exists an unique strict monoidal functor $\tilde{X}: (\Delta_{f+}, \oplus, [-1]) \longrightarrow (\mathcal{E}, \otimes, I)$ such that the pointed object $\underline{\mathbf{Pt}}(\tilde{X})([0], \delta_0^0) = (\tilde{X}([0]), \tilde{X}(\delta_0^0))$ is equal to (X, e) .*

Proof. Standard, see Mac Lane [22]. It is worth remarking that observation 21 immediately implies that the functor $\tilde{X}$ is completely determined by $\tilde{X}([n]) = X^{\otimes(n+1)}$ and $\tilde{X}(\delta_i^n) = X^{\otimes(n-i)} \otimes e \otimes X^{\otimes i}$. $\square$

Observation 42 (the free simplicial set generated by a semi-simplicial set). In terms of the partial action description, a semi-simplicial set $(X, \dim, \cdot)$ is merely a set X equipped with a dimension function “dim” and a partial right action “ $\cdot$ ” by simplicial face operators. Under this interpretation, a semi-simplicial map $f: (X, \dim, \cdot) \longrightarrow (Y, \dim, \cdot)$ is simply a function from X to Y which preserves dimensions and right actions by face operators.

Clearly then, every simplicial set (resp. map) is also a semi-simplicial set (resp. map), giving rise to a canonical forgetful functor $U: \underline{\mathbf{Simp}} \longrightarrow \underline{\mathbf{SSimp}}$. Since limits and colimits are constructed point-wise in these categories it follows that this functor preserves them and thus has both left and right adjoints.

2.6. Analysing Products of Simplicial Sets - the Theory of Shuffles. The following observations, regarding products of standard simplices, are classical and are applied extensively in proving the “extension” lemmas of sections 7 and 8.

Observation 43 (nerves of partially ordered sets). Let $\underline{\mathbf{Ord}}$ denote the category of all partially ordered sets and order preserving maps between them. We know that Δ is a full subcategory of $\underline{\mathbf{Ord}}$ and it follows that we may extend the Yoneda functor $\Delta: \Delta \longrightarrow \underline{\mathbf{Simp}}$ to a functor $\Delta: \underline{\mathbf{Ord}} \longrightarrow \underline{\mathbf{Simp}}$ which carries the partially ordered set P to the simplicial set $\Delta(P)$ given by “homming out” of the subcategory Δ and into P . In other words, $\Delta(P)$ has n -simplices which are order preserving maps $x: [n] \longrightarrow P$ and the simplicial operator $\alpha: [m] \longrightarrow [n]$ acts on such a simplex by pre-composition $x \cdot \alpha = x \circ \alpha: [m] \longrightarrow P$. If $f: P \longrightarrow Q$ is an order preserving map then $\Delta(f)$ is the simplicial map obtained by post-composing each simplex of $\Delta(P)$ by f to obtain a simplex of $\Delta(Q)$. We call $\Delta(P)$ the *nerve* of the partially ordered set P .

We may construct a left adjoint to the nerve functor $\Delta: \underline{\mathbf{Ord}} \longrightarrow \underline{\mathbf{Simp}}$ by left Kan extending the inclusion functor $\Delta \hookrightarrow \underline{\mathbf{Ord}}$ along the Yoneda embedding $\Delta: \Delta \longrightarrow \underline{\mathbf{Simp}}$ (cf. Kelly [19] for instance). It follows that the nerve construction preserves all limits and in particular that it preserves the product $[n] \times [m]$ giving a canonical isomorphism $\Delta([n] \times [m]) \cong \Delta[n] \times \Delta[m]$ under which we shall usually identify these simplicial sets.

More explicitly, recall from observations 34 and 31 that the r -simplices of $\Delta[n] \times \Delta[m]$ are pairs (α, β) of simplicial operators $\alpha: [r] \longrightarrow [n]$ and $\beta: [r] \longrightarrow [m]$ and that these are acted upon by right composition $(\alpha, \beta) \cdot \gamma = (\alpha \circ \gamma, \beta \circ \gamma)$. The corresponding r -simplex in $\Delta([n] \times [m])$ is the unique order preserving map $[r] \longrightarrow [n] \times [m]$ induced by the universal

property of $[n] \times [m]$ applied to the pair (α, β) . As is traditional, we shall blur our notation a little by using (α, β) to denote both the pair of simplicial operators which represent a simplex in $\Delta[n] \times \Delta[m]$ and the single order preserving map $(\alpha, \beta): [r] \longrightarrow [n] \times [m]$, given by $(\alpha, \beta)(i) = (\alpha(i), \beta(i))$, which represents the corresponding simplex in $\Delta([n] \times [m])$.

In general, our basic notational conventions for simplicial operators extend in a natural way to partially ordered sets and maps. For instance, if $f: P \longrightarrow Q$ is an arrow of $\mathbf{Ord}$ then we use the notation $\text{im}(f)$ to denote its image $\{f(p) \in Q \mid p \in P\}$. Consequently, under the convention of the last paragraph, it follows that $\text{im}(\alpha, \beta)$ denotes the subset $\{(\alpha(i), \beta(i)) \mid i \in [r]\} \subseteq [n] \times [m]$ which we think of as being the set of vertices of the r -simplex (α, β) in $\Delta[n] \times \Delta[m]$. We leave other such mild generalisations to the imagination of the reader.

Definition 44 (shuffles). The non-degenerate $(n + m)$ -simplices of $\Delta[n] \times \Delta[m]$ are called *shuffles*. Exploiting the identification of $\Delta[n] \times \Delta[m]$ with the nerve $\Delta([n] \times [m])$, as discussed in the last observation, we may think of a shuffle as a (strict) path of maximal length $(n + m)$ in the ordered set $[n] \times [m]$ as depicted in figure 1.

Lemma 45 (properties of products and shuffles). *The following properties hold for the simplicial set $\Delta[n] \times \Delta[m]$:*

- (1) *Integer addition gives a strictly order preserving surjective map $+: [n] \times [m] \longrightarrow [n + m]$, to which we may apply the nerve functor of observation 43 to give a simplicial map $+: \Delta[n] \times \Delta[m] \longrightarrow \Delta[n + m]$ which carries a simplex (α, β) to the simplex $\alpha + \beta$ given point-wise by $(\alpha + \beta)(j) = \alpha(j) + \beta(j)$.*
- (2) *A simplex (α, β) is degenerate in $\Delta[n] \times \Delta[m]$ if and only if $\alpha + \beta$ is degenerate in $\Delta[n + m]$. It follows that any simplex of $\Delta[n] \times \Delta[m]$ of dimension greater than $(n + m)$ is degenerate.*
- (3) *The projection map that takes an $(n + m)$ -simplex (α, β) of $\Delta[n] \times \Delta[m]$ to the operator $\alpha: [n + m] \longrightarrow [n]$ establishes a bijection between the set of shuffles of $\Delta[n] \times \Delta[m]$ and the set of degeneracy operators $\alpha: [n + m] \longrightarrow [n]$.*
- (4) *There exists a bijection between the set of degeneracy operators $\alpha: [n + m] \longrightarrow [n]$ and the set of simplicial operators $\gamma: [n] \longrightarrow [m]$ with $\gamma(n) = m$.*
- (5) *Suppose that $\gamma: [n] \longrightarrow [m]$ is a simplicial operator with $\gamma(n) = m$ and that $(\alpha_\gamma, \beta_\gamma)$ is the shuffle it corresponds to under the bijections of (3) and (4). Let (α, β) be an arbitrary r -simplex of $\Delta[n] \times \Delta[m]$, then the following propositions hold:*
 - (a) *if (α, β) is an r -dimensional face of $(\alpha_\gamma, \beta_\gamma)$ then it is the face obtained by applying the operator $\alpha + \beta: [r] \longrightarrow [n + m]$ to the given shuffle, or in other words:*

$$(\alpha, \beta) = (\alpha_\gamma, \beta_\gamma) \cdot (\alpha + \beta) = (\alpha_\gamma \circ (\alpha + \beta), \beta_\gamma \circ (\alpha + \beta))$$
 - (b) *for an $l \in [r]$ we have $\alpha(l) = \alpha_\gamma \circ (\alpha + \beta)(l)$ and $\beta(l) = \beta_\gamma \circ (\alpha + \beta)(l)$ if and only if the inequalities*


$$\gamma(\alpha(l) - 1) \leq \beta(l) \leq \gamma(\alpha(l)) \tag{4}$$

hold (in which we take $\gamma(-1) \stackrel{\text{def}}{=} 0$ where necessary),

- (c) *consequently, (α, β) is a face of $(\alpha_\gamma, \beta_\gamma)$ if and only if the inequalities of (4) hold for all $l \in [r]$.*

- (6) *Every simplex in $\Delta[n] \times \Delta[m]$ is a face of some shuffle.*

Proof. These results are fundamentally classical in nature, and we leave their detailed verification up to the reader. However we sketch the main plot points.

FIGURE 1. A shuffle (α, β) in $\Delta[n] \times \Delta[m]$

Part (1) is clear from the statement and it is easily verified that part (2) follows from the observation that $+: [n] \times [m] \longrightarrow [n + m]$ is strictly order preserving. Part (3) is a direct consequence of the fact that a simplex (α, β) is a shuffle if and only if $\alpha + \beta = \text{id}_{[n+m]}$, which itself follows directly from (1) and (2).

Part (4) is more interesting, the intuition here is that a shuffle (α, β) , as depicted in figure 1, may be fully specified by providing a list of “plateau” levels $\gamma(0), \gamma(1), \dots, \gamma(n-1)$, one for each column in the figure (as marked). A moment’s reflection reveals that $\gamma(i)$, the height of the “plateau” in the i^{th} column, may be given by the formula:

$$\gamma(i) = \max \{j \in [n + m] \mid \alpha(j) \leq i\} - i \quad (5)$$

Of course, from observation 15 we know that the degeneracy operator α has a right adjoint α^r and that $\alpha^r(i) = \max\{j \in [n + m] \mid \alpha(j) \leq i\}$, so we may re-express this as $\gamma(i) = \alpha^r(i) - i$. Furthermore, observation 15(iv) reveals that α^r is a face operator and so we can apply observation 7(iii) to construct $\gamma: [n] \longrightarrow [m]$ as the unique simplicial operator satisfying the equation:

$$\text{id}_{[n]} + \gamma = \alpha^r$$

It is now a routine matter to apply observations 7 and 15 and show that this construction provides the bijection postulated in the statement of (4).

Given the intuition we developed in the last paragraph, regarding the relationship between shuffles (α, β) and operators $\gamma: [n] \longrightarrow [m]$, the meaning of inequality (4) in the statement of part (5) should now be clear. It simply states that in the l^{th} vertex of (α, β) is a vertex of the shuffle $(\alpha_\gamma, \beta_\gamma)$ if and only if its vertical ordinate $\beta(l)$ lies between the plateaus of $(\alpha_\gamma, \beta_\gamma)$ immediately to either side of its horizontal ordinate $\alpha(l)$ (the levels of which we know to be $\gamma(\alpha(l) - 1)$ and $\gamma(\alpha(l))$ respectively). Given this observation, it is a matter of straightforward calculation, using the results we have already established, to show that the results of part (5) also hold.

Finally, to demonstrate that part (6) holds, we start with the r -simplex (α, β) and define a simplicial operator $\gamma: [n] \longrightarrow [m]$ by

$$\gamma(i) = \begin{cases} \min\{\beta(l) \mid l \in [r] \wedge \alpha(l) > i\} & \text{if } \alpha(r) > i, \\ m & \text{otherwise.} \end{cases} \quad (6)$$

for which it is a matter of routine verification to demonstrate that the condition of part (5)(c) holds. It follows that with γ defined in this way we can infer that our simplex (α, β) is a face of the shuffle $(\alpha_\gamma, \beta_\gamma)$. $\square$

Notation 46. We will adopt the following notational conventions with respect to the results embodied in the previous lemma:

- Rather than consider simplicial operators $\gamma: [n] \longrightarrow [m]$ with $\gamma(n) = m$, which were useful in proving the results of the lemma, we will instead work with the set of *all* simplicial operators $\gamma: [n-1] \longrightarrow [m]$ which, whenever necessary, we implicitly extend to $[n]$ by setting $\gamma(n) = m$.
- The notation $(\alpha_\gamma, \beta_\gamma)$ will be used to denote the shuffle associated with a simplicial operator $\gamma: [n-1] \longrightarrow [m]$ (as in lemma 45(5)).
- Given an r -simplex (α, β) in $\Delta[n] \times \Delta[m]$, the notation $\gamma_{(\alpha, \beta)}: [n-1] \longrightarrow [m]$ will be used to denote the simplicial operator defined in display (6) of the proof of lemma 45(6).

Observation 47 (A linear ordering of shuffles). The set of simplicial operators from $[n-1]$ to $[m]$ and thus, by the bijection of lemma 45, the set of shuffles in $\Delta[n] \times \Delta[m]$ may be linearly ordered by the “lexicographic ordering” relation $\triangleleft$, defined by

$$\gamma \triangleleft \tau \iff (\exists i \in [n-1])(\gamma(0) = \tau(0) \wedge \gamma(1) = \tau(1) \wedge \cdots \wedge \gamma(i-1) = \tau(i-1) \wedge \gamma(i) < \tau(i))$$

which extends the (strict) point-wise ordering of observation 14, in other words whenever $\gamma < \tau$ (ie whenever $\gamma \leq \tau$ and $\gamma \neq \tau$) then we also have $\gamma \triangleleft \tau$. In particular, the shuffle $(\pi_1^{n,m}, \pi_2^{n,m})$ of partition operators (notation 5) is minimal under the linear ordering $\triangleleft$ and $(\pi_2^{m,n}, \pi_1^{m,n})$ is maximal under the same relation. We also often use the notation $\trianglelefteq$ to denote the non-strict version of $\triangleleft$, in other words $\gamma \trianglelefteq \tau$ if and only if $\gamma \triangleleft \tau$ or $\gamma = \tau$.

Let $\#(n, m)$ denote the cardinality of the set of shuffles of $\Delta[n] \times \Delta[m]$. We know, from the fact that $\triangleleft$ is a linear ordering, that there exists a unique enumeration of these shuffles (respectively the simplicial operators from $[n-1]$ to $[m]$) $\{(\alpha_i, \beta_i)\}_{i=1}^{\#(n,m)}$ (respectively $\{\gamma_i\}_{i=1}^{\#(n,m)}$) which is compatible with $\triangleleft$ in the sense that $(\alpha_i, \beta_i) \triangleleft (\alpha_{i'}, \beta_{i'})$ (respectively $\gamma_i \triangleleft \gamma_{i'}$) if and only if $i < i'$. Of course, since $\triangleleft$ extends the point-wise order $<$, it follows that in order to demonstrate that $i < i'$ it is sufficient to show that $\gamma_i < \gamma_{i'}$.

Observation 48. Notice that if (α, β) is an r -simplex of $\Delta[n] \times \Delta[m]$ then the associated operator $\gamma: [n-1] \longrightarrow [m]$ of display (6) in the proof of lemma 45(6) is the upper bound, in the point-wise ordering $\leq$, of the set of operators $\tau: [n-1] \longrightarrow [m]$ for which (α, β) is a face of $(\alpha_\tau, \beta_\tau)$.

To prove this suppose that $\tau: [n-1] \longrightarrow [m]$ is such an operator and observe that the definition of γ tells us that for each $i \in [n-1]$ either $m = \gamma(i)$ or there exists an $l \in [r]$ with $\alpha(l) > i$ (equivalently $\alpha(l) - 1 \geq i$) and $\gamma(i) = \beta(l)$. In the first case we know that $\tau(i) \leq m = \gamma(i)$, since m is the largest integer the codomain of τ and in the second case we may apply lemma 45(5) to show that $\tau(\alpha(i) - 1) \leq \beta(i)$ which we combine with the (in)equalities of the last sentence and order preservation by τ to give $\tau(i) \leq \tau(\alpha(l) - 1) \leq \beta(l) = \gamma(i)$ as required. So, by the point-wise definition of the ordering $\leq$ on simplicial operators, it follows that $\gamma \geq \tau$.

In particular, applying the last observation we see that if w is the unique integer with $\gamma = \gamma_w$ then it may be characterised as being the largest integer for which (α, β) is a face of the shuffle (α_w, β_w) .

3. SOME CATEGORICAL BACKGROUND

3.1. Reflective Full Sub-Categories. We will often be interested in a context in which we wish to study a reflective full subcategory of some “carrier” category. In these cases it is often inconvenient to work in the subcategory alone. Instead, it is usually better to make explicit calculations in the carrier category and then to reflect the results of these exploits into the subcategory.

Observation 49 (reflective sub-categories). A *reflective full subcategory* $\mathcal{E}'$ of $\mathcal{E}$ is simply a full subcategory which is *replete*, in the sense that it is closed under isomorphisms in $\mathcal{E}$, and for which the inclusion $I: \mathcal{E}' \hookrightarrow \mathcal{E}$ has a left adjoint $F: \mathcal{E} \longrightarrow \mathcal{E}'$.

We follow Kelly [19] (page 53) by selecting this adjunction so that its counit is the identity and concentrating our attention on the pair (L, η) consisting of the endo-functor $L = IF: \mathcal{E} \longrightarrow \mathcal{E}$ and the unit $\eta: \text{id}_{\mathcal{E}} \Longrightarrow L$ of the adjunction $F \dashv I$. It is easily seen that this pair satisfies the identities $LL = L$ and $\eta L = L\eta = \text{id}_L$ (see observations 12 and 13 for more regarding the 2-categorical notation here) making it into an *idempotent monad* on $\mathcal{E}$.

Indeed, any such idempotent monad (L, η) on $\mathcal{E}$ gives rise to a reflective full subcategory $\mathcal{E}_L$ on those objects $A \in \mathcal{E}$ which are isomorphic to $L(X)$ for some $X \in \mathcal{E}$; we might say that $\mathcal{E}_L$ is the *replete image* of L . Notice that an object $A \in \mathcal{E}$ is in $\mathcal{E}_L$ if and only if the corresponding component of the unit η_A is an isomorphism. It may be easily demonstrated that the required left adjoint to the inclusion $\mathcal{E}_L \hookrightarrow \mathcal{E}$ is provided by L itself. In this way we get a 1-1 correspondence between idempotent monads on $\mathcal{E}$ and its reflective full sub-categories.

We will usually find it easier to identify these two concepts and work exclusively with idempotent monads (L, η) . Consequently, we refer to the L in such a structure as a *reflector* and the natural transformation η as its *unit*. Usually if we say that $\mathcal{E}_L$ is a reflective full subcategory of $\mathcal{E}$ we take it as assumed that we have an associated idempotent monad (L, η) which defines $\mathcal{E}_L$ in the way given above.

In general, we will use capitals from the end of the alphabet $X, Y, Z \dots$ to denote objects in $\mathcal{E}$ and those at the beginning $A, B, C \dots$ to denote objects in the subcategory $\mathcal{E}_L$.

For the next few observations we work within the context of a fixed category $\mathcal{E}$ and a chosen reflective full subcategory $\mathcal{E}_L$.

Definition 50 (L -invertible arrows). An arrow $f: X \longrightarrow Y$ in $\mathcal{E}$ is said to be *L -invertible* if and only if its image under the reflector L is an isomorphism.

Observation 51 (orthogonality). Equivalently, by Yoneda’s lemma, an arrow f is L -invertible if and only if it is *orthogonal* to each object $A \in \mathcal{E}_L$, in the sense that if $g: X \longrightarrow A$ is any map with $A \in \mathcal{E}_L$ then it admits a unique extension $\bar{g}: Y \longrightarrow A$ along f (i.e. $\bar{g} \circ f = g$):

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow g & \nearrow \exists! \bar{g} \\ & A & \end{array}$$

We often define reflective full sub-categories in terms of orthogonality conditions. In these cases, we start with a set S of arrows $\mathcal{E}$, sometimes called a *regulus*, and define $\mathcal{E}_{\perp S}$ to be the full subcategory of those objects in $\mathcal{E}$ which are orthogonal to all of the arrows in S . Usually,

proving that such an $\mathcal{E}_{\perp S}$ is actually reflective in $\mathcal{E}$ is a matter of applying some form of completion process expressed as a transfinite construction, as described in Kelly [18].

In general, if we are given a set of L-invertible arrows S then we say that it is *adequate to detect* objects of $\mathcal{E}_L$ if we can show that any object in $\mathcal{E}$ which is orthogonal to all arrows in S is an object of $\mathcal{E}_L$. Observe that this condition holds iff $\mathcal{E}_{\perp S} = \mathcal{E}_L$.

Observation 52 (fundamental properties of reflective full sub-categories and L-invertible arrows). The following observations follow directly by standard elementary categorical arguments, which we leave up to the reader:

- 1) If $D: \mathbb{C} \longrightarrow \mathcal{E}_L$ is a diagram with limit $\lim(D)$ in $\mathcal{E}$ then $\lim(D)$ is actually in the subcategory $\mathcal{E}_L$. We simply say that $\mathcal{E}_L$ is *closed in $\mathcal{E}$ under all limits which exist there*.
- 2) If $D: \mathbb{C} \longrightarrow \mathcal{E}_L$ is a diagram then we may form its colimit in $\mathcal{E}_L$ by constructing $\text{colim}(D)$ in $\mathcal{E}$ and reflecting it into $\mathcal{E}_L$ using L . The corresponding colimit cocone in $\mathcal{E}_L$ is obtained by composing the colimit cocone in $\mathcal{E}$ with the unit map $\eta_{\text{colim}(D)}: \text{colim}(D) \longrightarrow L(\text{colim}(D))$.
- 3) The condition $L\eta = \text{id}_L$ satisfied by our idempotent monad (L, η) may be usefully restated as saying that for each $X \in \mathcal{E}$ the component $\eta_X: X \longrightarrow L(X)$ of the unit η is L-invertible.
- 4) If $\mathcal{E}_{L'}$ is another reflective full subcategory of $\mathcal{E}$ then $\mathcal{E}_L$ is contained in $\mathcal{E}_{L'}$ if and only if every L' -invertible arrow is L-invertible.
- 5) The class of L-invertible arrows is closed under composition, right cancellation of epimorphisms / L-invertibles and left cancellation of L-invertibles. In other words:
 - if $f: X \longrightarrow Y$ and $g: Y \longrightarrow Z$ are both L-invertible then so is their composite $g \circ f: X \longrightarrow Z$,
 - if $g \circ f: X \longrightarrow Z$ is L-invertible and $f: X \longrightarrow Y$ is either epimorphic or L-invertible then the factor $g: Y \longrightarrow Z$ is also L-invertible, and
 - if $g \circ f: X \longrightarrow Z$ is L-invertible and either $g: Y \longrightarrow Z$ is L-invertible or $f: X \longrightarrow Y$ is epimorphic then the factor $f: X \longrightarrow Y$ is also L-invertible.
- 6) Consequently, if the arrows $m: X \longrightarrow Y$ and $e: Y \longrightarrow X$ form an inclusion / retraction pair (that is $e \circ m = \text{id}_X$) then m is L-invertible if and only if e is L-invertible.
- 7) The class of L-invertible arrows is closed under colimits. More explicitly, if the transformation of diagrams $\gamma: D \Rightarrow D': \mathbb{C} \longrightarrow \mathcal{E}$ has the property that each of its components γ_c ($c \in \text{obj}(\mathbb{C})$) is L-invertible, then the induced map $\bar{\gamma}: \text{colim}(D) \longrightarrow \text{colim}(D')$ is also L-invertible.
- 8) The class of L-invertible arrows is closed under pushout. Explicitly, if the square

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ h \downarrow & \lrcorner & \downarrow k \\ Z & \xrightarrow{g} & W \end{array}$$

is a pushout in $\mathcal{E}$ and f is L-invertible then so is g .

Observation 53 (L-almost concepts). We are often interested in proving categorical results about $\mathcal{E}_L$ by making calculations within $\mathcal{E}$. This usually allows us to exploit simpler explicit descriptions of objects in $\mathcal{E}$ than are available to us for the corresponding objects in $\mathcal{E}_L$. However, in order to do this we will need to generalise certain categorical concepts. Broadly speaking if we are given a categorical property “X” of a given structure then we say that such a structure in $\mathcal{E}$ *L-almost has property “X”* if applying L to it gives us a structure which has

property “X” in $\mathcal{E}_L$. For instance an arrow $f: X \longrightarrow Y$ is L-almost an epimorphism if the arrow $L(f): L(X) \longrightarrow L(Y)$ is an epimorphism in $\mathcal{E}_L$.

In particular, if $D: \mathbb{C} \longrightarrow \mathcal{E}$ is a diagram then we say that the cocone $i: D \twoheadrightarrow X$ in $\mathcal{E}$ is *L-almost colimiting* iff the cone $L(i): L(D(-)) \twoheadrightarrow L(X)$ is a colimiting cocone in $\mathcal{E}_L$. Notice that if the colimit $\text{colim}(D)$ exists in $\mathcal{E}$ then we know that our cocone induces a unique arrow $\hat{i}: \text{colim}(D) \longrightarrow X$ and that this is L-invertible if and only if our cocone is L-almost colimiting.

Most abstract properties of L-almost constructions may be proved simply by reflecting them into the subcategory $\mathcal{E}_L$ and then exploiting the corresponding traditional categorical theorem in that context.

Observation 54 (partial adjoints). If $F: \mathcal{E} \longrightarrow \mathcal{F}$ is a functor then, following Kelly [19] (page 50), we say that F has a *partial right adjoint* U defined on a full subcategory $\mathcal{F}'$ of $\mathcal{F}$ if there exists a functor $U: \mathcal{F}' \longrightarrow \mathcal{E}$ and a family of isomorphisms

$$\mathcal{F}(F(X), A) \cong \mathcal{E}(X, U(A))$$

which is natural in $X \in \mathcal{E}$ and $A \in \mathcal{F}'$.

Suppose now that we are given reflective full sub-categories $\mathcal{E}_L$ and $\mathcal{F}_{L'}$ of $\mathcal{E}$ and $\mathcal{F}$ respectively. In such cases, we are predominantly interested in those partial right adjoints which are defined on $\mathcal{F}_{L'}$ **and** map into $\mathcal{E}_L$. The following results hold by elementary arguments, in which we assume that $U: \mathcal{F}_{L'} \longrightarrow \mathcal{E}$ is a partial right adjoint to $F: \mathcal{E} \longrightarrow \mathcal{F}$:

- (1) the image of U is contained in $\mathcal{E}_L$ if and only if F carries all L-invertible arrows in $\mathcal{E}$ to L' -invertible arrows in $\mathcal{F}$.
- (2) if S is a set of L-invertible arrows which is adequate to detect objects of $\mathcal{E}_L$ and F maps all elements of S to L' -invertible arrows in $\mathcal{F}$ then the image of U is contained in $\mathcal{E}_L$.
- (3) if the image of U is contained in $\mathcal{E}_L$ then it is (genuinely) right adjoint to the composite $L'F: \mathcal{E}_L \longrightarrow \mathcal{F}_{L'}$.
- (4) under the same conditions the functor F preserves almost colimits, in the sense that if $D: \mathbb{C} \longrightarrow \mathcal{E}$ is a diagram and the cocone $i: D \twoheadrightarrow X$ is L-almost colimiting in $\mathcal{E}$ then the cocone $Fi: FD \twoheadrightarrow FX$ is L' -almost colimiting in $\mathcal{F}$.

The proofs of these propositions are a matter of routine abstract category theory and are left up to the reader. When speaking of partial right adjoints we will distinguish those that satisfy the above property simply by notationally restricting their codomains. In other words, we will simply say that $U: \mathcal{F}_{L'} \longrightarrow \mathcal{E}_L$ is a partial right adjoint to $F: \mathcal{E} \longrightarrow \mathcal{F}$.

We’ll also make heavy use of the following, slightly non-standard, version of Day’s reflection theorem [10] for monoidal biclosed categories:

Theorem 55 (Day’s reflection theorem). *Suppose that $(\mathcal{E}, \otimes, I)$ is a monoidal category and that $\mathcal{E}_L$ is a reflective full subcategory of $\mathcal{E}$ with associated idempotent monad (L, η) . Also assume that for objects $X, Y \in \mathcal{E}$ the right and left tensoring functors $- \otimes Y: \mathcal{E} \longrightarrow \mathcal{E}$ and $X \otimes -: \mathcal{E} \longrightarrow \mathcal{E}$ have partial right adjoints $* \Leftarrow Y: \mathcal{E}_L \longrightarrow \mathcal{E}_L$ and $X \Rightarrow *: \mathcal{E}_L \longrightarrow \mathcal{E}_L$ respectively.*

As usual, we may enrich the family of functors $ \Leftarrow Y$ (resp. $X \Rightarrow *$) with a canonical functorial structure in the variable Y (resp. X), thereby forming them into a bi-functor $\Leftarrow: \mathcal{E}_L \times \mathcal{E}^{\text{op}} \longrightarrow \mathcal{E}_L$ (resp. $\Rightarrow: \mathcal{E}^{\text{op}} \times \mathcal{E}_L \longrightarrow \mathcal{E}_L$).*

Under these assumptions, the following important results hold:

- (1) We may reflect the monoidal structure $(\mathcal{E}, \otimes, I)$ onto $\mathcal{E}_L$, making it into a monoidal biclosed category with:
- tensor product $A \otimes_L B \stackrel{\text{def}}{=} L(A \otimes B)$ and
 - left and right closures $A \Rightarrow *$ and $* \Leftarrow B$.
- (2) An arrow $f: X \longrightarrow Y$ in $\mathcal{E}$ is L -invertible if and only if $A \Leftarrow f: A \Leftarrow Y \longrightarrow A \Leftarrow X$ (or dually $f \Rightarrow A: Y \Rightarrow A \longrightarrow X \Rightarrow A$) is an isomorphism for each $A \in \mathcal{E}_L$.
- (3) The right and left tensoring functors $- \otimes Y, X \otimes -: \mathcal{E} \longrightarrow \mathcal{E}$ both preserve L -almost colimits.
- (4) If $D: \mathbb{C} \longrightarrow \mathcal{E}$ is a diagram then a cocone $i: D \dashrightarrow X$ is L -almost colimiting if and only if for each $A \in \mathcal{E}_L$ the contravariant functor $A \Leftarrow *: \mathcal{E}^{\text{op}} \longrightarrow \mathcal{E}_L$ (or dually $* \Rightarrow A: \mathcal{E}^{\text{op}} \longrightarrow \mathcal{E}_L$) maps it to a limit cone in $\mathcal{E}_L$.

Proof. Our proof of (1) follows Day's original argument of [10], which revolves around the observation that if X and Y are objects in $\mathcal{E}$ then the arrows $\eta_X \otimes Y: X \otimes Y \longrightarrow L(X) \otimes Y$, $X \otimes \eta_Y: X \otimes Y \longrightarrow X \otimes L(Y)$ and $\eta_X \otimes \eta_Y: X \otimes Y \longrightarrow L(X) \otimes L(Y)$ are all L -invertible. This follows from the observation that each component of the unit η is L -invertible (cf. observation 52(3)), that the left and right tensoring functors $X \otimes -$ and $- \otimes Y$ both preserve L -invertibility (by observation 54(1) and the assumption that they have appropriate partial right adjoints) and that L -invertible arrows are closed under composition.

This immediately allows us to enrich the bi-functor $\otimes_L$ with a monoidal structure which makes L into a strong monoidal functor $L: (\mathcal{E}, \otimes, I) \longrightarrow (\mathcal{E}_L, \otimes_L, L(I))$. For instance, we may then construct an associativity isomorphism for $\otimes_L$ as the composite

$$\begin{aligned} A \otimes_L (B \otimes_L C) &= L(A \otimes L(B \otimes C)) \xrightarrow{L(A \otimes \eta_{(B \otimes C)})^{-1}} L(A \otimes (B \otimes C)) \\ &\xrightarrow{L(\alpha_{A, B, C})} L((A \otimes B) \otimes C) \\ &\xrightarrow{L(\eta_{(A \otimes B)} \otimes C)} L(L(A \otimes B) \otimes C) = (A \otimes_L B) \otimes_L C \end{aligned}$$

in which the isomorphisms in the first and last line are a consequence of the L -invertibility result of the previous paragraph. To enrich $L: \mathcal{E} \longrightarrow \mathcal{E}_L$ to a strong monoidal functor we have the isomorphism $L(\eta_X \otimes \eta_Y): L(X \otimes Y) \longrightarrow L(L(X) \otimes L(Y)) \stackrel{\text{def}}{=} L(X) \otimes_L L(Y)$ obtained by applying L to the L -invertible arrow $\eta_X \otimes \eta_Y$. Furthermore, the fact that the bi-functors $\Leftarrow$ and $\Rightarrow$ provide us with right and left closures for this structure follows directly from observation 54(3).

To prove (2) observe that if $A \in \mathcal{E}_L$ then $A \Leftarrow f$ is defined to be the unique arrow making the diagram

$$\begin{array}{ccc} \mathcal{E}(Z \otimes Y, A) & \xrightarrow{\mathcal{E}(Z \otimes f, A)} & \mathcal{E}(Z \otimes X, A) \\ \cong \downarrow & & \downarrow \cong \\ \mathcal{E}(Z, A \Leftarrow Y) & \xrightarrow{\mathcal{E}(Z, A \Leftarrow f)} & \mathcal{E}(Z, A \Leftarrow X) \end{array}$$

commute for each $Z \in \mathcal{E}$. By Yoneda's lemma, $A \Leftarrow f$ is an isomorphism iff the map at the bottom of this diagram is an isomorphism for each Z and, since the verticals in the diagram are isomorphisms, it follows that this happens iff the map at the top of the diagram is an isomorphism. However, by definition, this latter condition simply states that $Z \otimes f$ is orthogonal to A for all $Z \in \mathcal{E}$.

So if $A \Leftarrow f$ is an isomorphism for each $A \in \mathcal{E}_L$ then it follows, since f is isomorphic to $I \otimes f$, that f is orthogonal to each $A \in \mathcal{E}_L$ and thus that f is L -invertible. Conversely, if f is L -invertible then we know that $Z \otimes -$ preserves L -invertibility, so $Z \otimes f$ is also L -invertible for each $Z \in \mathcal{E}$ and thus orthogonal to any $A \in \mathcal{E}_L$, thereby demonstrating that $A \Leftarrow f$ is an isomorphism for each $A \in \mathcal{E}_L$ by the argument of the last paragraph.

Part (3) is simply a restatement of observation 54(4) so all that remains is to prove part (4). To do so we simply apply the reflector L , and the natural family of isomorphisms $A \Leftarrow \eta_X: A \Leftarrow L(X) \xrightarrow{\cong} A \Leftarrow X$ arising from part (2) of this theorem (and the fact that each η_X is L -invertible), to reflect the desired result regarding L -almost colimits into a corresponding characterisation of (genuine) colimits in the biclosed monoidal category $\mathcal{E}_L$. This latter result follows by an elementary argument in the theory of biclosed monoidal categories, which we leave up to the reader. $\square$

3.2. LFP Categories and LE-Theories. In this subsection we simply summarise those parts of the theory of locally finitely presentable categories and left exact theories that we apply in the subsequent text. For greater detail we refer the reader to any number of standard presentations of this material, including Gabriel and Ulmer's book [12], Kennison's work on limit preserving functors [21], Barr and Wells' textbook [3] and Kelly's book on enriched category theory [19].

Definition 56 (weighted limits and colimits). Recall, from Kelly [19], that the colimit of a diagram $D: \mathbb{C} \longrightarrow \mathcal{E}$ *weighted* or *indexed* by $W: \mathbb{C}^{\text{op}} \longrightarrow \underline{\text{Set}}$ is an object $\text{colim}(W, D)$ of $\mathcal{E}$ which represents the functor $C_{W,D}(E) = [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](W, \mathcal{E}(D(*), E))$ from $\mathcal{E}$ to $\underline{\text{Set}}$. The unit of this representation is a natural transformation $\iota: W \longrightarrow \mathcal{E}(D(*), \text{colim}(W, D))$ called its *colimiting cylinder*. In this context, the traditional colimit of D is referred to as its *conical colimit* and it may be obtained by weighting our colimit by the functor which maps each object of $\mathbb{C}$ to a singleton set.

This weighted colimit construction (wherever it exists) is covariantly functorial in each argument. To be precise, any natural transformation of weights $\gamma: W \longrightarrow V \in [\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$ induces an unique arrow $\text{colim}(\gamma, D): \text{colim}(W, D) \longrightarrow \text{colim}(V, D)$ in $\mathcal{E}$ which represents the family of pre-composition functions $- \circ \gamma: [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](V, \mathcal{E}(D(*), E)) \longrightarrow [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](W, \mathcal{E}(D(*), E))$. Similarly, if $\phi: D \longrightarrow D'$ is a natural transformation of diagrams then the induced arrow $\text{colim}(W, \phi): \text{colim}(W, D) \longrightarrow \text{colim}(W, D')$ represents the family of post-compositions $\mathcal{E}(\phi_*, E) \circ -: [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](W, \mathcal{E}(D(*), E)) \longrightarrow [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](W, \mathcal{E}(D'(*), E))$.

Dually the limit $\lim(U, D) \in \mathcal{E}$ of our diagram weighted by $U: \mathbb{C} \longrightarrow \underline{\text{Set}}$ represents the functor $L_{U,D}(E) = [\mathbb{C}, \underline{\text{Set}}](U, \mathcal{E}(E, D(*)))$. It is worth remarking that under the duality involved here $\lim(U, D)$ becomes contravariantly functorial in its first variable and covariantly so in its second.

Let $\mathcal{Y}: \mathbb{C} \longrightarrow [\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$ denote the usual Yoneda functor, which maps an object $c \in \mathbb{C}$ to the representable $\mathcal{Y}_c = \mathbb{C}(-, c)$, and observe that Yoneda's lemma immediately implies that $\text{colim}(W, \mathcal{Y}) \cong W$ and that $\text{colim}(\mathcal{Y}_c, D) \cong D(c)$. It is also worth recalling that from the weight W we may construct a small comma category $\mathbb{G}(W) = \mathcal{Y} \downarrow W$, called the *Grothendieck category* of W . By Yoneda's lemma we can describe $\mathbb{G}(W)$ as a category whose objects are pairs (c, x) , where c is an object of $\mathbb{C}$ and $x \in W(c)$, and in which an arrow $f: (c, x) \longrightarrow (c', x')$ is simply an arrow $f: c \longrightarrow c'$ of $\mathbb{C}$ such that $W(f')(x') = x$. Composing the obvious projection functor $p: \mathbb{G}(W) \longrightarrow \mathbb{C}$ with the diagram $D: \mathbb{C} \longrightarrow \mathcal{E}$ and taking the conical colimit of this, if it exists, we obtain an object which represents the functor $C_{W,D}$ of the last paragraph. In other

words the weighted limit $\text{colim}(W, D)$ exists if and only if the conical colimit $\text{colim}_{\mathbb{G}(W)}(D \circ p)$ does and in that case these colimits are isomorphic.

Definition 57 (filtered colimits). We say that a small, non-empty category $\mathbb{D}$ is *filtered* iff

- given any two objects $d, d' \in \mathbb{D}$ there is a third object $\bar{d} \in \mathbb{D}$ to which they both map, and
- given any parallel pair of arrows $f, g: d \longrightarrow d'$ in $\mathbb{D}$ there exists a $h: d' \longrightarrow \bar{d}$ with $h \circ f = h \circ g$.

A *filtered colimit* is a (conical) colimit on a diagram whose indexing category is filtered.

Most importantly, in the category Set filtered colimits bear a simple description which allows us to easily demonstrate that they commute with all finite limits.

We say that a functor $F: \mathcal{E} \longrightarrow \mathcal{F}$ is *finitely accessible* iff it preserves all the (small) filtered colimits which exist in $\mathcal{E}$.

Definition 58 (locally finitely presentable categories). We say that a category $\mathcal{E}$ is *locally finitely presentable* (or simply an *LFP-category*) iff it is (equivalent to a) reflective full subcategory of some functor category $[\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$ for which $\mathbb{C}$ is a small category and the reflector L associated with $\mathcal{E}$ is finitely accessible.

In general, we say that an object E in some category $\mathcal{E}$ is *finitely presentable* (or simply *FP*) iff the associated representable functor $\mathcal{E}(A, -): \mathcal{E} \longrightarrow \underline{\text{Set}}$ preserves all (small) filtered colimits. Using this concept, we may provide a simple “intrinsic” characterisation of LFP-categories as being those locally small and small cocomplete categories $\mathcal{E}$ which possess a small dense set of finitely presentable objects.

Observation 59 (reflective full sub-categories of LFP-categories). If (L, η) is an idempotent monad on an LFP-category $\mathcal{E}$ for which L is finitely accessible then it is a routine matter to demonstrate that the corresponding reflective full subcategory $\mathcal{E}_L$ is also an LFP-category.

In this context, it is quite common to specify such reflective full sub-categories by supplying a defining *FP-regulus* T , which is simply a (essentially small) set of arrows in $\mathcal{E}$ all of whose domains and codomains are FP-objects. Applying theorem 10.2 of [18], we see that the full subcategory $\mathcal{E}_{\perp T}$ of objects orthogonal to T is reflective in $\mathcal{E}$. Furthermore, an immediate consequence of our assumption of finitely presentable domains and codomains is that $\mathcal{E}_{\perp T}$ is closed in $\mathcal{E}$ under filtered colimits. It follows that the reflector associated with $\mathcal{E}_{\perp T}$ must preserve filtered colimits and that we may apply the result of the previous paragraph to demonstrate that this subcategory is also LFP.

Definition 60 (left exact theories). LFP-categories may be presented explicitly in terms of a kind of categorical theory $\mathbb{T} = (\mathbb{C}, T)$ called a *left exact theory* (or simply an *LE-theory*) which consists of a small category $\mathbb{C}$ and an FP-regulus T in the presheaf category $[\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$. We use the notation $\mathbb{T}\text{-}\underline{\text{Alg}}$ to denote the reflective full-subcategory $[\mathbb{C}^{\text{op}}, \underline{\text{Set}}]_{\perp T}$ of objects orthogonal to the arrows in the FP-regulus T , call its objects $\mathbb{T}$ -*algebras* and use the notation $L_{\mathbb{T}}$ for the corresponding finitely accessible reflector.

We may generalise the $\mathbb{T}$ -algebra notion representably to any category $\mathcal{E}$, by defining such algebras to be those contravariant functors $A: \mathbb{C}^{\text{op}} \longrightarrow \mathcal{E}$ satisfying the condition that for each object $E \in \mathcal{E}$ the functor $\mathcal{E}(E, A(*)) \in [\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$ is actually a $\mathbb{T}$ -algebra. We use the notation $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E})$ to denote the full subcategory of the functor category $[\mathbb{C}^{\text{op}}, \mathcal{E}]$ of those functors which are $\mathbb{T}$ -algebras in $\mathbb{C}$, and refer to the arrows of $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E})$ as $\mathbb{T}$ -*algebra morphisms*.

However, we are usually only interested in considering $\mathbb{T}$ -algebras in categories $\mathcal{E}$ which possess all finite limits. These are called *LE-categories* and functors between them which preserve all finite limits are called *LE-functors*. Notice that when W is a finitely presented object of $[\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$ we may express it as a finite colimit of representables and consequently show that $\lim(W, A)$ may be constructed as a finite limit in $\mathcal{E}$. So if $\mathcal{E}$ is an LE-category then for each $\gamma: W \longrightarrow V$ in the regulus T we may form the arrow $\lim(\gamma, A): \lim(V, A) \longrightarrow \lim(W, A)$ in $\mathcal{E}$ representing the pre-composition $- \circ \gamma: [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](V, \mathcal{E}(E, A(*))) \longrightarrow [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](W, \mathcal{E}(E, A(*)))$ for each object $E \in \mathcal{E}$ and therefore we see, by Yoneda's lemma, that $\mathcal{E}(E, A(*))$ is orthogonal to γ for all objects $E \in \mathcal{E}$ if and only if $\lim(\gamma, A)$ is an isomorphism. It follows that in the context of LE-categories the $\mathbb{T}$ -algebra property may be re-expressed as stating that the arrow $\lim(\gamma, A): \lim(V, A) \longrightarrow \lim(W, A)$ is an isomorphism for each arrow $\gamma: W \longrightarrow V$ in the FP-regulus T .

Having expressed our $\mathbb{T}$ -algebra property in terms of finite limits we immediately see that if $F: \mathcal{E} \longrightarrow \mathcal{F}$ is an LE-functor then the post-composition functor $F \circ -: [\mathbb{C}^{\text{op}}, \mathcal{E}] \longrightarrow [\mathbb{C}^{\text{op}}, \mathcal{F}]$ carries $\mathbb{T}$ -algebras in $\mathcal{E}$ to $\mathbb{T}$ -algebras in $\mathcal{F}$. Therefore it restricts to a functor from $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E})$ to $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{F})$ which we will sometimes call $\mathbb{T}\text{-}\underline{\text{Alg}}(F)$, although in many cases it is convenient to overload our notation and simply think of F itself as having been extended or lifted to a functor $F: \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}) \longrightarrow \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{F})$. We often say that $\mathbb{T}\text{-}\underline{\text{Alg}}(F)$ is obtained by *applying* F *point-wise to the $\mathbb{T}$ -algebras in its domain*.

Observation 61 (more about LE-theories). Sometimes it is more convenient to present LE-theories in terms of *finite sketches* rather than FP-reguli. A finite sketch on a small category $\mathbb{C}$ consists of a family of diagrams $D_i: \mathbb{D}_i \longrightarrow \mathbb{C}$ ($i \in I$), for which each $\mathbb{D}_i$ is a finite category, and a family of cocones $\iota_i: D_i \longrightarrow c_i$ ($i \in I$). We say that a functor $A: \mathbb{C}^{\text{op}} \longrightarrow \underline{\text{Set}}$ is an algebra for such a sketch if it carries each of the specified cocones to a limiting cone in $\underline{\text{Set}}$.

From a sketch we may construct an FP-regulus whose members are obtained by applying the Yoneda functor $\mathcal{Y}: \mathbb{C} \longrightarrow [\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$ to each of its diagrams and cocones, taking the colimit of the former to obtain finitely presented objects $\text{colim}_{\mathbb{D}_i} (\mathcal{Y} \circ D_i) \in [\mathbb{C}^{\text{op}}, \underline{\text{Set}}]$ and forming the arrows $f_i: \text{colim}(\mathcal{Y} \circ D_i) \longrightarrow \mathcal{Y}_{c_i}$ induced by the latter. A trivial application of Yoneda's lemma now demonstrates that the algebras for our sketch and its associated FP-regulus coincide.

These definitions are motivated by the observation that a $\mathbb{T}$ -algebra in $\underline{\text{Set}}$ for an LE-theory is really no-more-nor-less than a family of basic underlying sets equipped with a family of operations whose domains and codomains are finite limits of those basic sets and for which a specified class of finitary equational conditions must hold. In particular these operations and equational conditions are coded into the structure of $\mathbb{C}$ while the FP-regulus T (or sketch) specifies precisely which limits constitute the domains and codomains of each operation. We often think of each of these operations as being partially defined upon equationally specified subsets of products of basic sets.

We refer the reader to [3] for a thorough exposition of theory of LE-theories, including a proof of Kennison's theorem which demonstrates that $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E})$ is a reflective full subcategory of $[\mathbb{C}^{\text{op}}, \mathcal{E}]$ (for all sufficiently cocomplete LE-categories $\mathcal{E}$). It is also straightforward to show that $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E})$ is closed in $[\mathbb{C}^{\text{op}}, \mathcal{E}]$ under all the limits and filtered colimits which exist in there. Combining these two results, it follows that $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E})$ is an LFP-category for any LFP-category $\mathcal{E}$.

Observation 62 (tensor products of LE-theories). Thus far we know that if $\mathbb{T}$ and $\mathbb{S} = (\mathbb{D}, S)$ are LE-theories and $\mathcal{E}$ is an LFP-category then the category $\mathbb{S}\text{-}\underline{\text{Alg}}(\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}))$ is also LFP and

may, therefore, be presented as a category of algebras for some LE-theory in $\mathcal{E}$. However, for many purposes it is useful to have an explicit presentation of this theory in terms of $\mathbb{T}$ and $\mathbb{S}$ themselves. To this end we define their *tensor product* $\mathbb{T} \otimes \mathbb{S}$ to be the LE-theory with underlying category $\mathbb{C} \times \mathbb{D}$ and FP-regulus $T \otimes S$ in $[(\mathbb{C} \times \mathbb{D})^{\text{op}}, \underline{\text{Set}}]$ given by

$$T \otimes S = \{\gamma \circ \pi_{\mathbb{C}}^{\text{op}} : W \circ \pi_{\mathbb{C}}^{\text{op}} \longrightarrow V \circ \pi_{\mathbb{C}}^{\text{op}} \mid \gamma : W \longrightarrow V \in T\} \cup \\ \{\gamma' \circ \pi_{\mathbb{D}}^{\text{op}} : W' \circ \pi_{\mathbb{D}}^{\text{op}} \longrightarrow V' \circ \pi_{\mathbb{D}}^{\text{op}} \mid \gamma' : W' \longrightarrow V' \in S\}$$

where $\pi_{\mathbb{C}} : \mathbb{C} \times \mathbb{D} \longrightarrow \mathbb{C}$ and $\pi_{\mathbb{D}} : \mathbb{C} \times \mathbb{D} \longrightarrow \mathbb{D}$ are the canonical projection functors.

This definition is designed to ensure that a $\mathbb{T} \otimes \mathbb{S}$ -algebra in an LE-category $\mathcal{E}$ is precisely a functor $A : \mathbb{C} \times \mathbb{D} \longrightarrow \mathcal{E}$ satisfying the condition that for each object $c \in \mathbb{C}$ the functor $A(c, -) : \mathbb{D} \longrightarrow \mathcal{E}$ is a $\mathbb{S}$ -algebra and for each object $c' \in \mathbb{D}$ the functor $A(-, c') : \mathbb{C} \longrightarrow \mathcal{E}$ is a $\mathbb{T}$ -algebra. Furthermore, it is clear that the canonical isomorphisms

$$[\mathbb{C}^{\text{op}}, [\mathbb{D}^{\text{op}}, \mathcal{E}]] \cong [(\mathbb{C} \times \mathbb{D})^{\text{op}}, \mathcal{E}] \cong [\mathbb{D}^{\text{op}}, [\mathbb{C}^{\text{op}}, \mathcal{E}]]$$

of functor categories restrict to isomorphisms

$$\mathbb{T}\text{-}\underline{\text{Alg}}(\mathbb{S}\text{-}\underline{\text{Alg}}(\mathcal{E})) \cong \mathbb{T} \otimes \mathbb{S}\text{-}\underline{\text{Alg}}(\mathcal{E}) \cong \mathbb{S}\text{-}\underline{\text{Alg}}(\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}))$$

of categories of algebras which are natural in $\mathcal{E}$.

Observation 63 (Kan's construction - right adjoint functors from L-almost coalgebras). Of course, we say that a functor $C : \mathbb{C} \longrightarrow \mathcal{E}$ is a $\mathbb{T}$ -coalgebra if its dual is a $\mathbb{T}$ -algebra in $\mathcal{E}^{\text{op}}$. Taking the dual of our characterisation of $\mathbb{T}$ -algebras in LE-categories, we see that we can characterise $\mathbb{T}$ -coalgebras as being those functors $C : \mathbb{C} \longrightarrow \mathcal{E}$ such that for each arrow $\gamma : W \longrightarrow V$ in the FP-regulus T the induced arrow $\text{colim}(\gamma, C) : \text{colim}(W, C) \longrightarrow \text{colim}(V, C)$ is an isomorphism.

Our primary interest in $\mathbb{T}$ -coalgebras here is that each one gives rise to a right adjoint functor into the category of $\mathbb{T}$ -algebras, by an application of (a variant of) *Kan's construction* [17]. However, in the sequel we will actually be interested in taking the approach described in observation 53, that is we will work in a category $\mathcal{E}$ equipped with an idempotent monad (L, η) and use L-almost coalgebras to construct right adjoint functors with domain the reflective full subcategory $\mathcal{E}_L$.

Following the convention established in observation 53, we now define an L-almost $\mathbb{T}$ -coalgebra in $\mathcal{E}$ to be a functor $C : \mathbb{C} \longrightarrow \mathcal{E}$ for which the composite $L \circ C : \mathbb{C} \longrightarrow \mathcal{E}_L$ is a genuine $\mathbb{T}$ -coalgebra in $\mathcal{E}_L$. Using our characterisation of coalgebras in terms of colimits in $\mathcal{E}_L$ it follows that this is equivalent to saying that for each $\gamma : W \longrightarrow V$ in the FP-regulus T the induced arrow $\text{colim}(\gamma, C) : \text{colim}(W, C) \longrightarrow \text{colim}(V, C)$ in $\mathcal{E}$ is L-invertible. We use the notation $\mathbb{T}\text{-}\underline{\text{CoAlg}}_L(\mathcal{E})$ for the full subcategory of L-almost $\mathbb{T}$ -coalgebras in $[\mathbb{C}, \mathcal{E}]$. Again it follows that if the functor $F : \mathcal{E} \longrightarrow \mathcal{F}$ preserves L-almost finite colimits then it may be applied point-wise to provide a functor $\mathbb{T}\text{-}\underline{\text{CoAlg}}_L(F) : \mathbb{T}\text{-}\underline{\text{CoAlg}}_L(\mathcal{E}) \longrightarrow \mathbb{T}\text{-}\underline{\text{CoAlg}}_L(\mathcal{F})$.

Any L-almost $\mathbb{T}$ -coalgebra C gives rise to an adjoint pair

$$\begin{array}{ccc} & F_C & \\ & \curvearrowright & \\ \mathcal{E}_L & \xrightleftharpoons[\text{K}_C]{\perp} & \mathbb{T}\text{-}\underline{\text{Alg}} \end{array}$$

in which K_C is the *Kan functor* constructed by “homming out” of C . In other words, if E is an object of $\mathcal{E}_L$ then we define $K_C(E) = \mathcal{E}(C(-), E) : \mathbb{C}^{\text{op}} \longrightarrow \underline{\text{Set}}$ and if $f : E \longrightarrow E'$ is an arrow of $\mathcal{E}$ then $K_C(f)$ is the natural transformation with components $K_C(f)_c = \mathcal{E}(C(c), f) = f \circ -$ for $c \in \mathbb{C}$. That $K_C(E)$ is actually a $\mathbb{T}$ -algebra follows from the fact that we have the reflection

isomorphism $\mathcal{E}(C(-), E) \cong \mathcal{E}_L(L \circ C(-), E)$ for each $E \in \mathcal{E}_L$ and by definition we know that $L \circ C$ is a $\mathbb{T}$ -coalgebra in $\mathcal{E}_L$ iff the functor on the right of this isomorphism is a $\mathbb{T}$ -algebra (in $\underline{\text{Set}}$) for each object $E \in \mathcal{E}_L$.

This construction is clearly contravariantly functorial in the L -almost coalgebra C so we get a functor $K_*: \mathbb{T}\text{-CoAlg}_L(\mathcal{E})^{\text{op}} \longrightarrow [\mathcal{E}_L, \mathbb{T}\text{-Alg}]$. If a functor $G: \mathcal{E}_L \longrightarrow \mathbb{T}\text{-Alg}$ is isomorphic to the Kan functor on some L -almost coalgebra C then we say that G is *representable* or that it is *represented by* C .

Each $\mathbb{T}$ -algebra A can, of course, be considered to be a weight on $\mathbb{C}$ and so we may define $F_C(A)$ to be the colimit $\text{colim}(A, L \circ C)$ in $\mathcal{E}_L$. That this does indeed provide a left adjoint to K_C follows from the calculation:

$$\begin{aligned} \mathcal{E}_L(\text{colim}(A, L \circ C), E) &\cong [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](A, \mathcal{E}_L(L \circ C(-), E)) && \text{definition of weighted colimit,} \\ &\cong [\mathbb{C}^{\text{op}}, \underline{\text{Set}}](A, \mathcal{E}(C(-), E)) && \text{reflection isomorphism,} \\ &\cong \mathbb{T}\text{-Alg}(A, K_C(E)) && \text{definition of } K_C \end{aligned}$$

Finally, we say that an L -almost coalgebra C in $\mathcal{E}$ is *finitely presented* iff it maps each object of $\mathbb{C}$ to a finitely presented object in $\mathcal{E}$. We will sometimes have use for the routine observation that if the reflector L is a finitely accessible functor and the L -almost coalgebra C is finitely presented then K_C also preserves all filtered colimits. The proof of this result is a routine matter of elementary category theory, which we leave up to reader.

Observation 64 (An internal version of Kan’s construction). For the remainder of this subsection, we’ll consider a context in which $\mathcal{E}$ is an LFP-category that comes equipped with a finitely accessible idempotent monad (L, η) and the monoidal and partial closures described in our version of Day’s reflection theorem (theorem 55). In such situations we will be interested in studying $\mathbb{T}$ -algebras in the LFP biclosed reflective full subcategory $\mathcal{E}_L$ and constructing functors $\mathcal{E}_L \longrightarrow \mathbb{T}\text{-Alg}(\mathcal{E}_L)$ using L -almost $\mathbb{T}$ -coalgebras in $\mathcal{E}$.

Using the tensor product and partial closures on $\mathcal{E}$ we may internalise the Kan’s construction to derive right adjoint functors from $\mathcal{E}_L$ to $\mathbb{T}\text{-Alg}(\mathcal{E}_L)$. To see that this is the case recall from theorem 55 that for each object $E \in \mathcal{E}_L$ the contravariant left closure functor $* \Rightarrow E: \mathcal{E}^{\text{op}} \longrightarrow \mathcal{E}_L$ inverts L -invertible arrows and carries L -almost colimits in $\mathcal{E}$ to limits in $\mathcal{E}_L$. It follows therefore, from our colimit and limit characterisations, that post-composition by $* \Rightarrow E$ carries L -almost $\mathbb{T}$ -coalgebras in $\mathcal{E}$ to $\mathbb{T}$ -algebras in $\mathcal{E}_L$ and thus provides a functor $\mathbb{T}\text{-CoAlg}_L(\mathcal{E})^{\text{op}} \longrightarrow \mathbb{T}\text{-Alg}(\mathcal{E}_L)$. The functoriality of this construction in $E \in \mathcal{E}_L$ is clear and we have therefore succeeded in constructing a functor $K_*^l: \mathbb{T}\text{-CoAlg}_L(\mathcal{E})^{\text{op}} \longrightarrow [\mathcal{E}_L, \mathbb{T}\text{-Alg}(\mathcal{E}_L)]$ which maps an L -almost $\mathbb{T}$ -coalgebra $C: \mathbb{C} \longrightarrow \mathcal{E}$ to the Kan functor $K_C^l: \mathcal{E}_L \longrightarrow \mathbb{T}\text{-Alg}(\mathcal{E}_L)$ obtained by “internally homming out of C ” or explicitly:

$$\begin{aligned} K_C^l(E) &\stackrel{\text{def}}{=} C(-) \Rightarrow E \quad \text{for each object } E \in \mathcal{E}_L, \text{ and} \\ K_C^l(f)_c &\stackrel{\text{def}}{=} C(c) \Rightarrow f \quad \text{for each arrow } f \in \mathcal{E}_L \text{ and object } c \in \mathbb{C}. \end{aligned}$$

Furthermore, if $\alpha: C \longrightarrow C'$ is an L -almost $\mathbb{T}$ -coalgebra map then it is mapped to the natural transformation $K_\alpha^l: K_{C'}^l \longrightarrow K_C^l$ whose component at the object $E \in \mathcal{E}_L$ is the $\mathbb{T}$ -algebra morphism $(K_\alpha^l)_E \stackrel{\text{def}}{=} \mathbb{T}\text{-Alg}(* \Rightarrow E)(\alpha)$ which consists of the family of arrows $\alpha_c \Rightarrow E: C'(c) \Rightarrow E \longrightarrow C(c) \Rightarrow E$ indexed by objects $c \in \mathbb{C}$.

If A is a $\mathbb{T}$ -algebra in $\mathcal{E}_L$ then we may tensor it with the L -almost $\mathbb{T}$ -coalgebra and compose the result with L to give a functor $L(A(-) \otimes C(*)): \mathbb{C}^{\text{op}} \times \mathbb{C} \longrightarrow \mathcal{E}_L$. Now we may define

$F_C^l(A)$, the value of the left adjoint to K_C^l on the algebra A to be a certain kind of colimit of this diagram in $\mathcal{E}_L$ called its *coend* by Kelly in [19] and denoted by $\int^{c \in \mathbb{C}} (L(A(c) \otimes C(c))) \in \mathcal{E}_L$.

Notice also that if we know that the tensor product of any pair of finitely presentable objects in $\mathcal{E}$ is always finitely presentable then we may show that $C(c) \Rightarrow -$ preserves all filtered colimits in $\mathcal{E}_L$ whenever $C(c)$ is finitely presentable. Furthermore, we also know that $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L)$ is closed in $[\mathbb{C}, \mathcal{E}_L]$ under filtered limits which are calculated point-wise in there. Combining these two facts, it follows therefore that the functor $K_C^l: \mathcal{E}_L \longrightarrow \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L)$ preserves all filtered colimits whenever C is a finitely presented L-almost coalgebra.

Observe that if I is the identity object of the monoidal structure on $\mathcal{E}$ then, since we generalised the $\mathbb{T}$ -algebra notion to arbitrary categories representably, we know that post-composition by the representable functor $\mathcal{E}_L(L(I), -) \cong \mathcal{E}(I, -): \mathcal{E}_L \longrightarrow \underline{\text{Set}}$ gives a functor $\mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}(I, -)): \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L) \longrightarrow \mathbb{T}\text{-}\underline{\text{Alg}}$. By partial adjointness we also have natural isomorphisms $\mathcal{E}(I, C(-) \Rightarrow E) \cong \mathcal{E}(C(-) \otimes I, E) \cong \mathcal{E}(C(-), E)$ for each object $E \in \mathcal{E}_L$ and we may use these to construct an isomorphism $\mathcal{E}(I, -) \circ K_C^l \cong K_C$ in $[\mathcal{E}_L, \mathbb{T}\text{-}\underline{\text{Alg}}]$ which is natural in the L-almost $\mathbb{T}$ -coalgebra C .

Finally, we could equally well have used the partial right closure functor $E \Leftarrow *$ in the calculations above and thereby obtain another functor $K_*^r: \mathbb{T}\text{-}\underline{\text{CoAlg}}_L(\mathcal{E})^{\text{op}} \longrightarrow [\mathcal{E}_L, \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L)]$. In general, we will distinguish these constructions in our text by referring to the former as the *left handed* construction and the latter as the *right handed* version. If our monoidal structure on $\mathcal{E}$ is augmented by a suitable duality operation $(-)^{\circ}: \mathcal{E} \longrightarrow \mathcal{E}$ then we may relate these constructions as (appropriately defined) duals of each other.

Observation 65 (tensor products of L-almost coalgebras). Assume again that $\mathcal{E}$ is as in the last observation, and suppose that $C \in \mathbb{T}\text{-}\underline{\text{CoAlg}}_L(\mathcal{E})$ and $D \in \mathbb{S}\text{-}\underline{\text{CoAlg}}_L(\mathcal{E})$ then the functor $C \otimes D: \mathbb{C} \times \mathbb{D} \longrightarrow \mathcal{E}$ defined by $(C \otimes D)(c, d) \stackrel{\text{def}}{=} C(c) \otimes D(d)$ is an L-almost $\mathbb{T} \otimes \mathbb{S}$ -coalgebra called the *tensor product* of C and D .

To show that this is indeed an L-almost coalgebra consider an object $c \in \mathbb{C}$ and observe that $C(c) \otimes -$ preserves L-almost colimits, since it has a partial right adjoint $C(c) \Rightarrow *$, and thus may be applied point-wise to the L-almost $\mathbb{S}$ -coalgebra D to show that $(C \otimes D)(c, -) = C(c) \otimes D(-)$ is also an L-almost $\mathbb{S}$ -coalgebra. By a dual argument $(C \otimes D)(-, d) = C(-) \otimes D(d)$ is an L-almost $\mathbb{T}$ -coalgebra for each object $d \in \mathbb{D}$. Consulting observation 62 it is clear that these two observations are enough to demonstrate that $C \otimes D$ is an L-almost $\mathbb{T} \otimes \mathbb{S}$ -coalgebra as required.

This definition is motivated by the observation that the monoidal and partial closure structures on $\mathcal{E}$ provide canonical isomorphisms

$$\begin{aligned} D(d) \Rightarrow (C(c) \Rightarrow E) &\cong (C(c) \otimes D(d)) \Rightarrow E \\ (E \Leftarrow C(c)) \Leftarrow D(d) &\cong E \Leftarrow (D(d) \otimes C(c)) \end{aligned}$$

which are natural in $E \in \mathcal{E}_L$, $c \in \mathbb{C}$ and $d \in \mathbb{D}$ and thus may be collected together to provide natural isomorphisms

$$\begin{array}{ccc} \mathcal{E}_L & \xrightarrow{K_C^l} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L) \\ K_{C \otimes D}^l \downarrow & \cong & \downarrow \mathbb{T}\text{-}\underline{\text{Alg}}(K_D^l) \\ \mathbb{T} \otimes \mathbb{S}\text{-}\underline{\text{Alg}}(\mathcal{E}_L) & \xrightarrow{\cong} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathbb{S}\text{-}\underline{\text{Alg}}(\mathcal{E}_L)) \end{array} \quad \begin{array}{ccc} \mathcal{E}_L & \xrightarrow{K_C^r} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L) \\ K_{D \otimes C}^r \downarrow & \cong & \downarrow \mathbb{T}\text{-}\underline{\text{Alg}}(K_D^r) \\ \mathbb{S} \otimes \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L) & \xrightarrow{\cong} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathbb{S}\text{-}\underline{\text{Alg}}(\mathcal{E}_L)) \end{array} \quad (7)$$

respectively. Sometimes it is useful to partially externalise this result and use the partial adjunction isomorphisms

$$\mathcal{E}_L(D(d), C(c) \Rightarrow E) \cong \mathcal{E}(C(c) \otimes D(d), E) \quad \mathcal{E}_L(D(d), E \Leftarrow C(c)) \cong \mathcal{E}(D(d) \otimes C(c), E)$$

which are natural in $E \in \mathcal{E}_L$, $c \in \mathbb{C}$ and $d \in \mathbb{D}$ to construct natural isomorphisms

$$\begin{array}{ccc} \mathcal{E}_L & \xrightarrow{K_C^l} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L) \\ K_{C \otimes D} \downarrow & \cong & \downarrow \mathbb{T}\text{-}\underline{\text{Alg}}(K_D) \\ \mathbb{T} \otimes \mathbb{S}\text{-}\underline{\text{Alg}} & \xrightarrow{\cong} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathbb{S}\text{-}\underline{\text{Alg}}) \end{array} \quad \begin{array}{ccc} \mathcal{E}_L & \xrightarrow{K_C^r} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathcal{E}_L) \\ K_{D \otimes C} \downarrow & \cong & \downarrow \mathbb{T}\text{-}\underline{\text{Alg}}(K_D) \\ \mathbb{S} \otimes \mathbb{T}\text{-}\underline{\text{Alg}} & \xrightarrow{\cong} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathbb{S}\text{-}\underline{\text{Alg}}) \end{array} \quad (8)$$

respectively.

Observation 66. Consider the concrete case where $\mathbb{S}$ is an LE-theory, $\mathcal{E}$ is the presheaf category $[\mathbb{D}^{\text{op}}, \underline{\text{Set}}]$ and L is the reflector for which $\mathcal{E}_L$ is the category of algebras $\mathbb{S}\text{-}\underline{\text{Alg}}$.

Applying Yoneda's lemma as in definition 56, we know that if $\gamma: W \longrightarrow V$ is any natural transformation of weights then $\text{colim}(\gamma, \mathcal{Y}): \text{colim}(W, \mathcal{Y}) \longrightarrow \text{colim}(V, \mathcal{Y})$ is actually isomorphic to γ itself, so it follows immediately, by our colimit characterisation of L -almost coalgebras, that the Yoneda functor $\mathcal{Y}: \mathbb{D} \longrightarrow [\mathbb{D}^{\text{op}}, \underline{\text{Set}}]$ is an L -almost $\mathbb{S}$ -coalgebra. Furthermore, applying Yoneda's lemma once again we see that the associated functor $K_{\mathcal{Y}}$ is actually the identity functor on $\mathbb{S}\text{-}\underline{\text{Alg}}$.

Consequently if C is any L -almost $\mathbb{T}$ -coalgebra in $[\mathbb{D}^{\text{op}}, \underline{\text{Set}}]$ then we may apply this result in the context of the essentially commutative squares of display (8), reducing their right hand verticals to identities and yielding essentially commutative triangles:

$$\begin{array}{ccc} & \mathbb{S}\text{-}\underline{\text{Alg}} & \\ K_{C \otimes \mathcal{Y}} \swarrow & \cong & \searrow K_C^l \\ \mathbb{T} \otimes \mathbb{S}\text{-}\underline{\text{Alg}} & \xrightarrow{\cong} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathbb{S}\text{-}\underline{\text{Alg}}) \end{array} \quad \begin{array}{ccc} & \mathbb{S}\text{-}\underline{\text{Alg}} & \\ K_{\mathcal{Y} \otimes C} \swarrow & \cong & \searrow K_C^r \\ \mathbb{S} \otimes \mathbb{T}\text{-}\underline{\text{Alg}} & \xrightarrow{\cong} & \mathbb{T}\text{-}\underline{\text{Alg}}(\mathbb{S}\text{-}\underline{\text{Alg}}) \end{array}$$

In practise the left hand diagonals of these triangles provide conveniently explicit external presentations of the internal constructions on their right hand diagonals.

4. DOUBLE CATEGORIES, 2-CATEGORIES AND N-CATEGORIES

4.1. Categories in the Small.

Observation 67. In subsection 2.2 we introduced 2-categories “in the gros sense”, that is to say our definitions were selected to make them a little more amenable to expressing some of the meta-theory in this work. In this section we re-introduce categories, 2-categories and their higher dimensional relatives the ω -categories “in the petit sense”. That is to say, our purpose here is to present them as objects of algebraic study in their own right rather than as contexts within which to study the properties of other mathematical entities.

Notice here that the distinction we make between the gros and the petit isn’t really a matter of set-theoretic size. On the one hand, it is true that all of our petit categories and n -categories will be (defined to be) set-theoretically small and that in our meta-theory the gros categories and 2-categories we manipulate will be set-theoretically large or huge respectively. However, the important distinction here really is one of intention, petit categories are inhabitants of our theory while our gros categories are denizens of our meta-theory.

Definition 68 (categories in the small). A (small) category $\mathbb{C}$ is a sextuple $(\mathbb{C}_a, \mathbb{C}_o, *, i, s, t)$ consisting of

- a set $\mathbb{C}_o$ whose elements we think of as being the *objects* of $\mathbb{C}$,
- a set $\mathbb{C}_a$ whose elements which we think of as being the *arrows* of $\mathbb{C}$,
- a function $i: \mathbb{C}_o \longrightarrow \mathbb{C}_a$ which we think of as mapping an object to the *identity* arrow on that object,
- two functions $s, t: \mathbb{C}_a \longrightarrow \mathbb{C}_o$ mapping each arrow to its *source* and *target* objects respectively. Collectively, these functions must satisfy the condition that i is a right compositional inverse for both s and t (that is $t \circ i = s \circ i = \text{id}_{\mathbb{C}_o}$). In other words for any object $c \in \mathbb{C}_o$ the source and target of the corresponding identity arrow $i(c)$ must be c itself. We adopt the notation $s' \stackrel{\text{def}}{=} i \circ s$ and $t' \stackrel{\text{def}}{=} i \circ t$ to denote the idempotent functions which map each arrow to the identity arrows associated with its source and target objects respectively.
- a partially defined binary composition operation $*$ on $\mathbb{C}_a$, such that $q * p \in \mathbb{C}_a$ is defined for each pair of arrows $p, q \in \mathbb{C}_a$ with $s(q) = t(p)$ (we say that such a pair is *compositionally compatible*), and
- this data should satisfy the equations

$$s(q * p) = s(p) \quad t(q * p) = t(q) \quad t'(p) * p = p \quad p * s'(p) = p \quad r * (q * p) = (r * q) * p$$

whenever the various composites involved are well defined.

When speaking of more than one category, we will tend to follow traditional practise by using the symbols $*, i, s$ and t “polymorphically” and rely on context to disambiguate the particular instance of these operators being applied in any given case.

A functor $f: \mathbb{C} \longrightarrow \mathbb{C}'$ consists of a function $f: \mathbb{C}_a \longrightarrow \mathbb{C}'_a$ between sets of arrows, which must satisfy the conditions

$$s'(f(p)) = f(s'(p))t'(f(p)) = f(t'(p)) \quad f(q * p) = f(q) * f(p)$$

wherever the composites involved are well defined. Of course, the composite of (the underlying functions of) two functors is itself a functor, and so the collection of all categories and functors itself possesses the structure of a (large) category, which we shall call Cat.

Notice that a functor $f: \mathbb{C} \longrightarrow \mathbb{C}'$ gives rise to a map $f_o \stackrel{\text{def}}{=} s \circ f \circ i: \mathbb{C}_o \longrightarrow \mathbb{C}'_o$ of sets of objects which is the unique function for which any one (and thus all) of the equalities $i \circ f_o = f \circ i$, $f_o \circ s = s \circ f$ and $f_o \circ t = t \circ f$ hold.

Observation 69 (single-sorted categories). The category notion given in the last definition is often referred to as the *two-sorted* theory of categories, since each such category possesses both a sort of arrows and a separately specified sort of objects. In order to avoid definitional burden, some authors have chosen not to carry around this separate sort of objects and have instead relied upon a *single-sorted* presentation. In this definition we don't distinguish objects and arrows, instead we identify each object with its associated identity arrow.

Under such a presentation, a category becomes a quadruple $(\mathbb{C}_a, *, s', t')$ where s' and t' are endo-functions on $*$ satisfying the “absorption properties” $s' \circ s' = t' \circ s' = s'$ and $t' \circ t' = s' \circ t' = t'$ and a pair of arrows p and q in $\mathbb{C}_a$ is composable iff $t'(p) = s'(q)$. No information is lost in the process and we may regain the sort of objects $\mathbb{C}_o$ simply by splitting the idempotent $s': \mathbb{C}_a \longrightarrow \mathbb{C}_a$. That is we let $\mathbb{C}_o = \{p \in \mathbb{C}_a \mid s'(p) = p\}$, the set of *identity* arrows for the composition operation $*$.

The absorption properties of s' and t' ensure that we may factor them simultaneously through the inclusion i of $\mathbb{C}_o$ into $\mathbb{C}_a$, giving us functions $s, t: \mathbb{C}_a \longrightarrow \mathbb{C}_o$ for which $s' = i \circ s$, $s \circ i = \text{id}_{\mathbb{C}_o}$, $t' = i \circ t$ and $t \circ i = \text{id}_{\mathbb{C}_o}$. This data provides us with the underlying structure of a two-sorted category, with i as the identities function and s and t as source and target functions respectively.

Given this discussion, we will sometimes take the liberty of passing without comment between single-sorted and two-sorted descriptions of categories. However in most cases we will prefer the two-sorted definition, simply because it will be convenient in many proofs to carry around an explicitly chosen presentation of each set of objects.

Observation 70 (categories as algebras for an LE-theory). The theory of categories may be expressed as an LE-theory $\mathbb{T}_{\text{Cat}}$ which has underlying category Δ and the set of horn inclusions $\Lambda_k[n] \xrightarrow{\subseteq_s} \Delta[n] \in \underline{\text{Simp}} = [\Delta^{\text{op}}, \underline{\text{Set}}]$ ($n = 2, 3, \dots, k = 1, \dots, n - 1$) as its FP-regulus. We let $L_{\text{Cat}}: \underline{\text{Simp}} \longrightarrow \underline{\text{Simp}}$ denote the reflector associated with this theory.

Alternatively, it is also common to present this LE-theory in terms of a finite sketch with pushout-like cocones of the form

$$\begin{array}{ccc}
 [0] & \xrightarrow{\varepsilon_p^p} & [p] \\
 \varepsilon_0^q \downarrow & \searrow & \downarrow \varepsilon_1^{p,q} \\
 [q] & \xrightarrow{\varepsilon_2^{p,q}} & [p+q]
 \end{array} \tag{9}$$

in Δ for each pair of integers $p, q \geq 1$. The intuition behind this presentation is that if $A \in \underline{\text{Simp}} = [\Delta^{\text{op}}, \underline{\text{Set}}]$ is a $\mathbb{T}_{\text{Cat}}$ -algebra then we would like to think of its set of n -simplices A_n as being the set of composable paths of arrows of length n in some category. Were this the case, then we should be able to form a path in A_{p+q} by adjoining a pair of compatible paths in A_p and A_q , which is precisely what the algebra condition with respect to the cocone in display (9) states.

If $A \in \underline{\text{Simp}}$ is a $\mathbb{T}_{\text{Cat}}$ -algebra we can see that it gives rise to a category, in the sense of definition 68, with

- Sets of objects A_0 and arrows A_1 , the sets of 0- and 1-dimensional simplices in A respectively.
- Identity, source and target functions $\mathbf{i} \stackrel{\text{def}}{=} A_{\eta^1}$, $\mathbf{s} \stackrel{\text{def}}{=} A_{\varepsilon_0^1}$ and $\mathbf{t} \stackrel{\text{def}}{=} A_{\varepsilon_1^1}$, for which we may apply the functoriality of A , along with elementary equations in Δ , to demonstrate the required identity rules $\mathbf{s} \circ \mathbf{i} = \mathbf{t} \circ \mathbf{i} = \text{id}_{A_0}$.
- From notation 37 we know that a simplicial map $f: \Lambda_1[2] \longrightarrow A$ corresponds to a composable pair of arrows in A_1 and since A is a $\mathbb{T}_{\text{Cat}}$ -algebra it is perpendicular to $\Lambda_1[2] \xrightarrow{\subseteq_s} \Delta[2]$ and it follows that the function $(-\cdot\delta_0^2, -\cdot\delta_2^2): A_2 \longrightarrow A_1 \mathbf{s} \times_{\mathbf{t}} A_1$ is an isomorphism with inverse which we'll denote by w . In other words, if p and q are a composable pair of arrows then there exists a (unique) 2-simplex $w(p, q) \in A_2$ with $p = w(p, q) \cdot \delta_2^2$ and $q = w(p, q) \cdot \delta_0^2$ and we define their composite $q * p$ to be the 3rd 1-dimensional face $w(p, q) \cdot \delta_1^2$ of $w(p, q)$. We say that $w(p, q)$ *witnesses* the composite $q * p$.
- The associativity of this composition operation can be established by taking a composable sequence p, q, r of 1-simplices and exploiting orthogonality to $\Lambda_1[2] \xrightarrow{\subseteq_s} \Delta[2]$ three times to build a 2-dimensional 1-horn which we fill to a 3-simplex using orthogonality to $\Lambda_1[3] \xrightarrow{\subseteq_s} \Delta[3]$. The four 2-dimensional faces of this 3-simplex witness the various composites involved in the two sides of the associativity condition, viz $r * (q * p)$ and $(r * q) * p$, however its 1-dimensional faces corresponding to these two composites coincide, and so the composites themselves must be equal as required.

If $f: A \longrightarrow A'$ is a simplicial map between $\mathbb{T}_{\text{Cat}}$ -algebras then it carries the 2-simplex witnessing a composite $q * p$ in A to the 2-simplex witnessing the composite $f(q) * f(p)$ in A' and so the action of f is functorial with respect to these category structures. It follows that this construction provides us with a functor from $\mathbb{T}_{\text{Cat}}\text{-Alg}(\text{Set})$ to Cat . and it is a matter of routine verification to demonstrate that this is an equivalence. It follows that the LE-theory $\mathbb{T}_{\text{Cat}}$ does indeed provide us with a suitable presentation of the theory of (small) categories. Hence forth, we shall move backward and forward between these presentations of the theory of categories without comment.

Observation 71 (relativising our theory). Generally, all of the theories introduced in this work, including the all important theory of complicial sets introduced later on, may be expressed as LE-theories. Furthermore, all of the theorems that we prove about these structures may be lifted *representably* (that is to say via an application of Yoneda's lemma) to an internal context where we take models of our LE-theories in an arbitrary LE-category.

Indeed, there will many places in the proofs that follow in which we will prove a theorem in Set and then apply it to algebras internal to some explicitly described LFP-category $\mathcal{E}$. In all such cases, however, it will be clear that the results we are applying immediately lift “point-wise” to corresponding results on structures defined in terms of the concrete algebras in $\mathcal{E}$.

Consequently, in future we will silently assume that the reader is aware of this process of *relativisation* and make no further comment when lifting results to an internal context. Furthermore, we will generally adopt the notational convention that if $\underline{\mathcal{C}}$ denotes the category of algebras of such a theory then $\underline{\mathcal{C}}(\mathcal{E})$ is used to denote the corresponding category of models in $\mathcal{E}$. As before, if $F: \mathcal{E} \longrightarrow \mathcal{E}'$ is an LE-functor then the corresponding functor of categories of algebras may sometimes be denoted by $\underline{\mathcal{C}}(F)$, but is most likely to simply be referred to by “overloading” the symbol F and saying that it lifts to a LE-functor $F: \underline{\mathcal{C}}(\mathcal{E}) \longrightarrow \underline{\mathcal{C}}(\mathcal{E}')$

4.2. Double Categories.

Definition 72 (double categories and double functors). A (small) double category is a model for the finite limit theory of categories in the category $\underline{\text{Cat}}$ of (small) categories and functors. We define the category $\underline{\text{Double}}$ to be $\underline{\text{Cat}}(\underline{\text{Cat}})$ and call its arrows *double functors*.

Alternatively we know, by observations 70 and 62, that a double category $\mathbb{D}$ may be presented more symmetrically as an algebra of the LE-theory $\mathbb{T}_{\underline{\text{Cat}}} \otimes \mathbb{T}_{\underline{\text{Cat}}}$, that is a functor $\mathbb{D}: \Delta^{\text{op}} \times \Delta^{\text{op}} \longrightarrow \underline{\text{Set}}$ such that each of the functors $\mathbb{D}([n], -), \mathbb{D}(-, [m]): \Delta^{\text{op}} \longrightarrow \underline{\text{Set}}$ are categories (presented as a $\mathbb{T}_{\underline{\text{Cat}}}$ -algebras) for all $[n], [m] \in \Delta$.

In more economical and explicit terms, we may present our double category $\mathbb{D}$ as having:

- a set $\mathbb{D}_s$ whose elements are called *squares*, sets $\mathbb{D}_v$ and $\mathbb{D}_h$ whose elements are called *vertical* and *horizontal* arrows respectively and a set $\mathbb{D}_o$ of *objects*.
- four category structures
 - $(\mathbb{D}_h, \mathbb{D}_o, *_h, \mathbf{i}_h, \mathbf{s}_h, \mathbf{t}_h)$ the *horizontal category of arrows*, denoted $\text{arr}_h(\mathbb{D})$,
 - $(\mathbb{D}_v, \mathbb{D}_o, *_v, \mathbf{i}_v, \mathbf{s}_v, \mathbf{t}_v)$ the *vertical category of arrows*, denoted $\text{arr}_v(\mathbb{D})$,
 - $(\mathbb{D}_s, \mathbb{D}_v, *_h, \mathbf{i}_h, \mathbf{s}_h, \mathbf{t}_h)$ the *horizontal category of squares*, denoted $\text{sq}_h(\mathbb{D})$, and
 - $(\mathbb{D}_s, \mathbb{D}_h, *_v, \mathbf{i}_v, \mathbf{s}_v, \mathbf{t}_v)$ the *vertical category of squares*, denoted $\text{sq}_v(\mathbb{D})$.

This data must satisfy the following rules:

- (i) the vertical structural functions $\mathbf{i}_v, \mathbf{s}_v$ and $\mathbf{t}_v$ of $\text{sq}_v(\mathbb{D})$ are functors with respect to the horizontal category structures $\text{arr}_h(\mathbb{D})$ and $\text{sq}_h(\mathbb{D})$,
- (ii) the horizontal structural functions $\mathbf{i}_h, \mathbf{s}_h$ and $\mathbf{t}_h$ of $\text{sq}_h(\mathbb{D})$ are functors with respect to the vertical category structures $\text{arr}_v(\mathbb{D})$ and $\text{sq}_v(\mathbb{D})$,
- (iii) the middle four interchange rule holds, that is for any quadruple of squares $\alpha, \beta, \gamma, \delta$ we have $(\delta *_h \gamma) *_v (\beta *_h \alpha) = (\delta *_v \beta) *_h (\gamma *_v \alpha)$ where the composite on the left is well defined iff that on the right is.

Notice that the middle four interchange rule is equivalent to saying that the horizontal composite map $*_h$ of $\text{sq}_h(\mathbb{D})$ is functorial with respect to the vertical category structure on $\text{sq}_v(\mathbb{D})$ and that of its pullback $\text{sq}_v(\mathbb{D})_{\mathbf{s}_h} \times_{\mathbf{t}_h} \text{sq}_v(\mathbb{D})$. Dually, this rule may be interpreted as postulating the functoriality of vertical composition $*_v$ with respect to horizontal category structures.

Under this presentation, a double functor $f: \mathbb{D} \longrightarrow \mathbb{D}'$ is simply a function $f: \mathbb{D}_s \longrightarrow \mathbb{D}'_s$ which is functorial with respect to both the vertical and the horizontal categories of squares of $\mathbb{D}$ and $\mathbb{D}'$. In other words, f is the underlying function of two functors $f: \text{sq}_v(\mathbb{D}) \longrightarrow \text{sq}_v(\mathbb{D}')$ and $f: \text{sq}_h(\mathbb{D}) \longrightarrow \text{sq}_h(\mathbb{D}')$.

Given a double category $\mathbb{D}$ presented as a $\mathbb{T}_{\underline{\text{Cat}}} \otimes \mathbb{T}_{\underline{\text{Cat}}}$ -algebra its corresponding explicit presentation may be obtained by:

- defining the various sets of squares and arrows in $\mathbb{D}$ to be:

$$\mathbb{D}_s \stackrel{\text{def}}{=} \mathbb{D}([1], [1]) \quad \mathbb{D}_o \stackrel{\text{def}}{=} \mathbb{D}([0], [0]) \quad \mathbb{D}_v \stackrel{\text{def}}{=} \mathbb{D}([1], [0]) \quad \mathbb{D}_h \stackrel{\text{def}}{=} \mathbb{D}([0], [1])$$

- letting the various categories $\text{arr}_h(\mathbb{D})$, $\text{arr}_v(\mathbb{D})$, $\text{sq}_h(\mathbb{D})$ and $\text{sq}_v(\mathbb{D})$ be those which correspond to the $\mathbb{T}_{\underline{\text{Cat}}}$ -algebras $D([0], -)$, $D(-, [0])$, $D([1], -)$ and $D(-, [1])$ respectively.

The isomorphisms of categories of algebras given at the end of observation 62, provide us with two distinct presentations of each “symmetrised” double category $\mathbb{D}$ as an internal category in $\underline{\text{Cat}}$, viz:

- its *horizontal presentation* $(\text{sq}_v(\mathbb{D}), \text{arr}_v(\mathbb{D}), *_h, \mathbf{i}_h, \mathbf{s}_h, \mathbf{t}_h)$, an internal category of vertical categories, and

- its *vertical presentation* $(\text{sq}_h(\mathbb{D}), \text{arr}_h(\mathbb{D}), *_v, \mathbf{i}_v, \mathbf{s}_v, \mathbf{t}_v)$, an internal category of horizontal categories.

Observation 73 (double categories in pictures). A square α of a double category $\mathbb{D}$ may literally be pictured as a square thus:

$$\begin{array}{ccc}
 & \mathbf{s}_v(\alpha) & \\
 \bullet & \xrightarrow{\quad} & \bullet \\
 \mathbf{s}_h(\alpha) \downarrow & \alpha & \downarrow \mathbf{t}_h(\alpha) \\
 \bullet & \xrightarrow{\quad} & \bullet \\
 & \mathbf{t}_v(\alpha) &
 \end{array}$$

Here, the vertical (resp. horizontal) lines represent the vertical (resp. horizontal) arrows which constitute the horizontal (resp. vertical) domain and codomain of α . It is a consequence of the functoriality of horizontal and vertical domain and codomain functions that adjacent pairs of sides in this diagram “meet up” at a corner, each of which is an object. Furthermore, two squares are horizontally (resp. vertically) composable if they abut at a common vertical (resp. horizontal) arrow. Finally, we can picture the middle four interchange rule in terms of the two possible ways of composing four squares, as shown in figure 2.

$$\begin{array}{ccccc}
 \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \alpha & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} & \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \beta & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} & \xrightarrow{\quad} & \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \beta *_h \alpha & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} \\
 \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \gamma & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} & \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \delta & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} & \xrightarrow{\quad} & \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \delta *_h \gamma & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} \\
 \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
 \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \gamma *_v \alpha & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} & \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & \delta *_v \beta & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} & \xrightarrow{\quad} & \begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \downarrow & (\delta *_h \gamma) *_v (\beta *_h \alpha) = \\ & (\delta *_v \beta) *_h (\gamma *_v \alpha) & \downarrow \\ \bullet & \xrightarrow{\quad} & \bullet \end{array}
 \end{array}$$

FIGURE 2. The middle four interchange rule for double categories.

4.3. 2-Categories and Double Categories with Connections. Now we are ready to give another definition of 2-categories, which is easily seen to be equivalent to that given in subsection 2.2:

Definition 74 (2-categories in the small). A category $\mathbb{C}$ is said to be *discrete* iff each of its arrows is an identity on some object. Equivalently $\mathbb{C}$ is discrete iff its identity function $\mathbf{i}: \mathbb{C}_o \longrightarrow \mathbb{C}_a$ is an isomorphism.

A 2-category is a double category $\mathbb{D}$ whose category of vertical arrows $\text{arr}_v(\mathbb{D})$ is discrete and a 2-functor is simply a double functor between 2-categories. We use the notation 2-Cat for the full subcategory of 2-categories in Double.

Observation 75 (The relationship to definition 10). Under our new definition, a 2-category is simply a double category in which each vertical arrow is an identity. In other words, if we depict identity arrows as equalities then we might depict the squares of a 2-category thus:

$$\begin{array}{ccc} \bullet & \xrightarrow{p} & \bullet \\ \parallel & \alpha & \parallel \\ \bullet & \xrightarrow{p'} & \bullet \end{array}$$

Contracting these equalities to a single point and depicting our square as a double arrow from its vertical domain to its vertical codomain we simply regain the “globular” 2-cell picture of observation 11.

Consequently, we will adopt the usual nomenclature for the structural components of 2-categories presented in this way, referring to their squares as 2-cells, horizontal arrows as 1-cells and objects as 0-cells.

We’ve seen that 2-categories may be considered simply to be special kinds of double categories. There is, however, a slightly less trivial construction by which we can build a double category out of a 2-category and which we shall need to apply in the sequel.

Observation 76 (the double category of pasting squares in a 2-category). If $\mathbb{C}$ is a 2-category then we may define a double category $\text{Sq}(\mathbb{C})$ with squares consisting of quintuples $(\psi, d_h, c_h, d_v, c_v)$ where c_h, d_h, c_v and d_v are all 1-cells (in $\mathbb{C}$), ψ is a 2-cell and this information fits together into a pasting square:

$$\begin{array}{ccc} \bullet & \xrightarrow{d_v} & \bullet \\ d_h \downarrow & \psi \swarrow & \downarrow c_h \\ \bullet & \xrightarrow{c_v} & \bullet \end{array}$$

The operations of horizontally and vertically pasting together such squares (see Kelly and Street [20] or Power [25]) suffices to provide us with natural horizontal and vertical composition operations. Furthermore, it is an immediate consequence of the theory of pasting that a 2-functor $f: \mathbb{C} \longrightarrow \mathbb{C}'$ gives rise to a double functor $\text{Sq}(f): \text{Sq}(\mathbb{C}) \longrightarrow \text{Sq}(\mathbb{C}')$, which acts “point-wise” on pasting squares as in the next diagram:

$$\begin{array}{ccc} \begin{array}{ccc} \bullet & \xrightarrow{d_v} & \bullet \\ d_h \downarrow & \psi \swarrow & \downarrow c_h \\ \bullet & \xrightarrow{c_v} & \bullet \end{array} & \longmapsto & \begin{array}{ccc} \bullet & \xrightarrow{f(d_v)} & \bullet \\ f(d_h) \downarrow & f(\psi) \swarrow & \downarrow f(c_h) \\ \bullet & \xrightarrow{f(c_v)} & \bullet \end{array} \end{array}$$

Observation 77 (duals of double categories and 2-categories). A double category $\mathbb{D}$ admits two basic kinds of duality:

- The *horizontal* and *vertical* duals $\mathbb{D}^{\text{hop}}$ and $\mathbb{D}^{\text{vop}}$ which are obtained by applying the usual categorical duality to its horizontal and vertical category structures respectively. We will often use $\mathbb{D}^{\text{hvp}}$ to denote the result of applying both of these dualities.
- The *reflection* dual $\mathbb{D}^{\text{r}}$ which is obtained by swapping the roles of these horizontal and vertical category structures.

Geometrically the first two dualities correspond to reversing the orientation of squares in one dimension or the other. The second corresponds to flipping or reflecting squares through their leading diagonals.

Of course, these dualities are all canonically functorial and involutive, in the sense that apply any one of them twice returns us to the double category we started with, and trivially interrelated via the (inter-derivable) identities $(\mathbb{D}^r)^{\text{hop}} = (\mathbb{D}^{\text{vop}})^r$ and $(\mathbb{D}^r)^{\text{vop}} = (\mathbb{D}^{\text{hop}})^r$.

2-categorists often use the notation $\mathbb{C}^{\text{op}}$ for the horizontal dual and $\mathbb{C}^{\text{co}}$ for the vertical dual of a 2-category $\mathbb{C}$. They also tend to use the notation $\mathbb{C}^{\text{coop}}$ to denote the 2-category obtained by applying both of these dualities.

Observation 78 (The 2-category of globs in a double category). For a double category $\mathbb{D}$, let $\text{Glob}(\mathbb{D})$ be the sub-double category of squares $\psi \in \mathbb{D}$ such that $t_h(\psi)$ and $s_h(\psi)$ are both identities in $\text{arr}_v(\mathbb{D})$. From this definition it is clear that $\text{Glob}(\mathbb{D})$ is actually a 2-category, which we call the *2-category of globs* in $\mathbb{D}$.

It is easily shown that a double functor $f: \mathbb{D} \longrightarrow \mathbb{D}'$ maps squares in $\text{Glob}(\mathbb{D})$ into $\text{Glob}(\mathbb{D}')$ and so restricts to a 2-functor $\text{Glob}(f): \text{Glob}(\mathbb{D}) \longrightarrow \text{Glob}(\mathbb{D}')$. It follows that Glob may be extended to a functor $\text{Glob}: \underline{\text{Double}} \longrightarrow \underline{2\text{-Cat}}$.

Notice that the 2-category of globs construction allows us to re-construct a 2-category from its double category of squares, in the sense that there exists a canonical family of isomorphisms $\mathbb{C} \cong \text{Glob}(\text{Sq}(\mathbb{C}))$ which is natural in the 2-category $\mathbb{C}$.

It follows that we may immediately infer that $\text{Sq}: \underline{2\text{-Cat}} \longrightarrow \underline{\text{Double}}$, the pasting squares functor, is faithful. However, it is **not** the case that this functor is also full, a fact which immediately begs two related questions:

- What extra structure might double categories of squares be asked to carry to restrict the maps between them and thus make the squares functor full?
- How can we characterise those double categories that arise by applying the squares functor to some 2-category?

The remainder of this subsection recalls the answers to these questions, in a form originally due to Chris Spencer and presented in Brown and Mosa [7].

Definition 79 (thin squares). A *double category with thinness* is a pair $(\mathbb{D}, t\mathbb{D})$ consisting of a double category $\mathbb{D}$ and a specified sub-double category $t\mathbb{D} \subseteq \mathbb{D}$ which contains all of the horizontal and vertical arrows of $\mathbb{D}$ (but which does not, in general, contain all of the squares of $\mathbb{D}$). It should come as no surprise that the squares of the *thinness structure* $t\mathbb{D}$ are said to be *thin*.

We form a category $\underline{\text{Double}}_T$ whose objects are double categories with thinness and whose arrows $f: (\mathbb{D}, t\mathbb{D}) \longrightarrow (\mathbb{D}', t\mathbb{D}')$ are double functors $f: \mathbb{D} \longrightarrow \mathbb{D}'$ which *preserve thinness* in the sense that they satisfy the condition $t\mathbb{D} \subseteq f^{-1}(t\mathbb{D}')$.

Clearly, in order to provide a double category $\mathbb{D}$ with a thinness structure it is enough to specify a subset $t\mathbb{D}_s$ of the set of squares of $\mathbb{D}$ which:

- contains all of the horizontal and vertical identity squares of $\mathbb{D}$, and
- is closed in $\mathbb{D}$ under horizontal and vertical composition of squares.

Observation 80. The double category of pasting squares in a 2-category $\mathbb{C}$ admits a canonical thinness structure under which we declare a square to be thin if its 2-cellular component is an identity in $\mathbb{C}$. If $f: \mathbb{C} \longrightarrow \mathbb{C}'$ is a 2-functor then it is clear that the associated double

functor $\text{Sq}(f): \text{Sq}(\mathbb{C}) \longrightarrow \text{Sq}(\mathbb{C}')$ preserves thinness with respect to these canonical structures. In other words, the pasting squares construction admits a canonical lifting to a functor $\text{Sq}: \underline{2}\text{-Cat} \longrightarrow \underline{\text{Double}}_T$.

Lemma 81 (Brown and Mosa [7]). *The functor $\text{Sq}: \underline{2}\text{-Cat} \longrightarrow \underline{\text{Double}}_T$ is fully faithful, furthermore a double category with thinness $(\mathbb{D}, t\mathbb{D})$ is in its replete image if and only if it satisfies the following conditions:*

- (i) *If α is a thin square such that $s_h(\alpha)$ and $t_h(\alpha)$ are identities in the category of vertical arrows $\text{arr}_v(\mathbb{D})$ then α is a vertical identity, diagrammatically:*

$$\text{if } \begin{array}{ccc} \bullet & \xrightarrow{p} & \bullet \\ \parallel & \alpha & \parallel \\ \bullet & \xrightarrow{p'} & \bullet \end{array} \text{ is thin then } \alpha = i_v(p) = i_v(p') \text{ (so } p = p') \text{ }$$

- (ii) *If α is a thin square such that $s_v(\alpha)$ and $t_v(\alpha)$ are identities in the category of horizontal arrows $\text{arr}_h(\mathbb{D})$ then α is a horizontal identity, diagrammatically:*

$$\text{if } \begin{array}{ccc} \bullet & \xlongequal{\quad} & \bullet \\ q \downarrow & \alpha & \downarrow q' \\ \bullet & \xlongequal{\quad} & \bullet \end{array} \text{ is thin then } \alpha = i_h(q) = i_h(q') \text{ (so } q = q') \text{ }$$

- (iii) *For each horizontal arrow $p \in \text{arr}_h(\mathbb{D})$ there exists a thin square μ_p such that $t_v(\mu_p) = p$, $s_v(\mu_p)$ is an identity in $\text{arr}_h(\mathbb{D})$ and $s_h(\mu_p)$ is an identity in $\text{arr}_v(\mathbb{D})$, diagrammatically:*

$$\text{for each } \bullet \xrightarrow{p} \bullet \text{ we have a thin } \begin{array}{ccc} \bullet & \xlongequal{\quad} & \bullet \\ \parallel & \mu_p & \downarrow \hat{p} \\ \bullet & \xrightarrow{p} & \bullet \end{array}$$

- (iv) *For each vertical arrow $q \in \text{arr}_v(\mathbb{D})$ there exists a thin square ν_q such that $s_h(\nu_q) = q$, $t_h(\nu_q)$ is an identity in $\text{arr}_v(\mathbb{D})$ and $t_v(\nu_q)$ is an identity in $\text{arr}_h(\mathbb{D})$, diagrammatically:*

$$\text{for each } \begin{array}{c} \bullet \\ q \downarrow \\ \bullet \end{array} \text{ we have a thin } \begin{array}{ccc} \bullet & \xrightarrow{\tilde{q}} & \bullet \\ q \downarrow & \nu_q & \parallel \\ \bullet & \xlongequal{\quad} & \bullet \end{array}$$

Note that the various squares μ_p and ν_q are called connection squares or simply connections and so a double category with thinness which satisfies the conditions above is often referred to as a double category with connections and we use $\underline{\text{Conn}}$ to denote the full subcategory of such things in $\underline{\text{Double}}_T$.

Proof. (sketch, see Brown and Mosa [7] for more detail) Notice that, as pointed out in observation 76, the 2-cells of $\text{Glob}(\text{Sq}(\mathbb{C}))$ are pretty much no more nor less than 2-cells in $\mathbb{C}$. A routine calculation shows that this correspondence extends to a (natural) isomorphism of 2-categories $\mathbb{C} \cong \text{Glob}(\text{Sq}(\mathbb{C}))$.

It remains to prove that for each double category with connections $(\mathbb{D}, t\mathbb{D})$ we have a natural isomorphism $(\mathbb{D}, t\mathbb{D}) \cong \text{Sq}(\text{Glob}(\mathbb{D}, t\mathbb{D}))$ in $\underline{\text{Double}}_T$. We do this in a number of steps:

1. Our conditions set up a bijection between the horizontal and vertical arrows of $\mathbb{D}$:

- (i) if p is a horizontal arrow then condition (iv) provides us with a thin square μ_p and, as a result, a vertical arrow $\hat{p} = t_h(\mu_p)$,
- (ii) if q is a vertical arrow then condition (iii) provides us with a thin square ν_q and a horizontal arrow $\tilde{q} = s_v(\nu_q)$,
- (iii) the horizontal arrow $\tilde{\hat{p}}$ is equal to p , which fact may be demonstrated by forming the horizontal composite $\nu_{\hat{p}} *_{\tilde{h}} \mu_p$ and appealing to condition (i).

$$\begin{array}{ccccc}
 & & & \xrightarrow{\tilde{\hat{p}}} & \\
 \bullet & \xrightarrow{\mu_p} & \bullet & & \bullet \\
 \parallel & & \downarrow \hat{p} & & \parallel \\
 \bullet & \xrightarrow{p} & \bullet & & \bullet
 \end{array}$$

- (iv) the vertical arrow $\widehat{\tilde{q}}$ is equal to q , which we demonstrate by forming the vertical composite $\nu_q *_{\tilde{v}} \mu_{\tilde{q}}$ and appealing to condition (ii).

$$\begin{array}{ccc}
 \bullet & \xrightarrow{\mu_{\tilde{q}}} & \bullet \\
 \parallel & & \downarrow \widehat{\tilde{q}} \\
 \bullet & \xrightarrow{\tilde{q}} & \bullet \\
 \downarrow q & & \parallel \\
 \bullet & & \bullet
 \end{array}$$

- (v) If p and q are a composable pair of vertical arrows then we have $\widetilde{q *_{\tilde{v}} p} = \tilde{q} *_{\tilde{h}} \tilde{p}$, which we can demonstrate by forming the composite $\nu_{q *_{\tilde{v}} p} *_{\tilde{h}} (\mu_{\tilde{q}} *_{\tilde{v}} p) *_{\tilde{h}} \mu_{\tilde{p}}$ appealing to condition (i).

$$\begin{array}{ccccc}
 \bullet & \xrightarrow{\mu_{\tilde{p}}} & \bullet & \xrightarrow{\mu_{\tilde{q}}} & \bullet \\
 \parallel & & \downarrow p & & \downarrow p \\
 \bullet & \xrightarrow{\tilde{p}} & \bullet & \xrightarrow{\tilde{q}} & \bullet \\
 \parallel & & \downarrow q & & \downarrow q \\
 \bullet & \xrightarrow{\tilde{p}} & \bullet & \xrightarrow{\tilde{q}} & \bullet
 \end{array}$$

These results provide us with a natural isomorphism between the horizontal category $\text{arr}_h(\mathbb{D})$ and the vertical category $\text{arr}_v(\mathbb{D})$. Consequently, from here on we identify these categories and simplify our notation by dropping explicit use of the accents $\sim$ and $\wedge$ to pass to and fro between horizontal and vertical arrows.

2. Each square $(\psi, d_h, c_h, d_v, c_v)$ in $\text{Sq}(\text{Glob}(\mathbb{D}, t\mathbb{D}))$ gives rise to a square ψ° in $\mathbb{D}$ of the form

$$\begin{array}{ccc}
 \bullet & \xrightarrow{d_v} & \bullet \\
 d_h \downarrow & \psi^\circ & \downarrow c_h \\
 \bullet & \xrightarrow{c_v} & \bullet
 \end{array}$$

obtained by taking the composite $(c_v * _h \nu_{d_h}) * _v \psi * _v (\mu_{c_h} * _h d_v)$, or diagrammatically:

$$\begin{array}{ccccc}
 \bullet & \xrightarrow{d_v} & \bullet & \xrightarrow{\quad} & \bullet \\
 \parallel & d_v & \parallel & \mu_{c_h} & \downarrow c_h \\
 \bullet & \xrightarrow{-d_v} & \bullet & \xrightarrow{-c_h} & \bullet \\
 \parallel & & \psi & & \parallel \\
 \bullet & \xrightarrow{-d_h} & \bullet & \xrightarrow{-c_v} & \bullet \\
 \downarrow d_h & \nu_{d_h} & \parallel & c_v & \parallel \\
 \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{c_v} & \bullet
 \end{array}$$

3. A square $\lambda \in \mathbb{D}$ gives rise to a pasting square $(\lambda^*, q, q', p, p')$ in $\text{Glob}(\mathbb{D}, t\mathbb{D})$ where $p = s_v(\lambda)$, $p' = t_v(\lambda)$, $q = s_h(\lambda)$, $q' = t_h(\lambda)$ and λ^* is the square given by the composite $\nu_{q'} * _h \lambda * _h \mu_q$, or diagrammatically:

$$\begin{array}{ccccc}
 \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{p} & \bullet & \xrightarrow{q'} & \bullet \\
 \parallel & \mu_q & \downarrow q & \lambda & \downarrow q' & \nu_{q'} & \parallel \\
 \bullet & \xrightarrow{q} & \bullet & \xrightarrow{p'} & \bullet & \xrightarrow{\quad} & \bullet
 \end{array}$$

4. It is a matter of routine calculation, involving heavy use of the middle four rule and conditions (i) and (ii), to demonstrate that these two operations are mutually inverse, respect thinness and preserve the composition structures of $(\mathbb{D}, t\mathbb{D})$ and $\text{Sq}(\text{Glob}(\mathbb{D}, t\mathbb{D}))$. It follows that they give rise to a (natural) isomorphism between these two double categories with connections, as required. $\square$

Observation 82 (duals and double categories of pasting squares). Later on we will need to understand the relationship between the various dualities of observation 77 and the constructions of the last lemma. For the horizontal and vertical dualities things are simple since it is easily seen that if $\mathbb{D}$ is any double category then $\text{Glob}(\mathbb{D}^{\text{hop}}) = \text{Glob}(\mathbb{D})^{\text{op}}$ and $\text{Glob}(\mathbb{D}^{\text{vop}}) = \text{Glob}(\mathbb{D})^{\text{co}}$. In general there is, however, no reason to believe that any simple relationship should hold between the 2-categories $\text{Glob}(\mathbb{D})$ and $\text{Glob}(\mathbb{D}^r)$.

One useful exception, which we shall apply later on, is the case when our double category is of the form $\text{Sq}(\mathbb{C})$ for some 2-category $\mathbb{C}$. In this case, it is easy to construct a canonical family of isomorphisms $\mathbb{C}^{\text{co}} \cong \text{Glob}(\text{Sq}(\mathbb{C})^r)$ which is natural in the 2-category $\mathbb{C}$. Equivalently, we can exploit the result of the last lemma to express this as a family of isomorphisms $\text{Glob}(\mathbb{D}, t\mathbb{D})^{\text{co}} \cong \text{Glob}(\mathbb{D}^r, t\mathbb{D}^r)$ which is natural in the double category with connections $(\mathbb{D}, t\mathbb{D}) \in \underline{\text{Conn}}$.

4.4. n -Categories and ω -Categories. One exception to preferring many sorted definitions over single sorted ones is that of ω -categories, which appears in the following form in Street [31]:

Definition 83 (ω -categories). An ω -category $\mathbb{C}$ consists of the following data

- a set of elements called *cells* and
- a family of (one-sorted) category structures $(*_i, s_i, t_i)$ one for each $i \in \mathbb{N}$,

which must satisfy the following conditions

- (i) if $i < j \in \mathbb{N}$ then category structures $(*_i, s_i, t_i)$ and $(*_j, s_j, t_j)$ constitute the horizontal and vertical category structures (respectively) of a 2-category structure on $\mathbb{C}$ and
- (ii) for each cell α there exists an $i \in \mathbb{N}$ such that α is an identity for the category structure $(*_i, s_i, t_i)$.

If $\mathbb{C}$ and $\mathbb{C}'$ are ω -categories then an ω -functor is simply a function $f: \mathbb{C} \longrightarrow \mathbb{C}'$ which is a functor with respect to the i^{th} category structure on $\mathbb{C}$ and $\mathbb{C}'$ for each $i \in \mathbb{N}$. As usual we use the notation $\omega\text{-}\underline{\text{Cat}}$ to denote the category of all ω -categories and ω -functors between them.

Observation 84 (*n*-cells and *n*-categories). A cell in an ω -category $\mathbb{C}$ is said to be an *n-cell* (for $n \in \mathbb{N}$) if it is an identity for the category structure $(*_n, s_n, t_n)$. Furthermore, we say that our cell is a *non-trivial n-cell* if it is an *n-cell* but it isn't an $(n-1)$ -cell. We will generally use the notation $\mathbb{C}_n \subseteq \mathbb{C}$ to denote the set of *n*-cells in $\mathbb{C}$. Condition (i) of definition 83 implies that if $m \geq n$ then any *n-cell* is also an *m-cell* (that is $\mathbb{C}_n \subseteq \mathbb{C}_m$).

An *n-category* is an ω -category in which every cell is an *n-cell* and $n\text{-}\underline{\text{Cat}}$ denotes the full subcategory of *n-categories* in $\omega\text{-}\underline{\text{Cat}}$. From the previous paragraph, we know that if $m \geq n$ then the category structure $(*_m, s_m, t_m)$ is *discrete*, in the sense that under $*_m$ cells may only compose with themselves. It follows that our latter definition of 2-category is entirely consistent with the one which we originally gave in definition 74, since all of its category structures aside from $*_0$ and $*_1$ are completely trivial.

Observe that the sources, targets and composites of *n*-cells are again *n*-cells, thus it follows that the ω -category structure of $\mathbb{C}$ restricts to provide us with an *n-category* structure on $\mathbb{C}_n$. Following Street [31], we adopt the notation $|\mathbb{C}|_n$ for this sub-*n-category* which we'll refer to as the *n-superstructure* of $\mathbb{C}$.

Of course, since an ω -functor $f: \mathbb{C} \longrightarrow \mathbb{D}$ preserves sources and targets it is clear that it maps *n*-cells to *n*-cells and thus restricts to an ω -functor $f: |\mathbb{C}|_n \longrightarrow |\mathbb{D}|_n$. This makes $|\cdot|_n$ into the underlying functor of an idempotent comonad on $\omega\text{-}\underline{\text{Cat}}$, the full image of which is the subcategory of *n-categories*.

The contrapositive of this “dimension preservation” property of the ω -functor f , the observation that if $f(c)$ is not an *n-cell* in $\mathbb{D}$ then c itself is not an *n-cell* in $\mathbb{C}$, is also, somewhat surprisingly, of use in its own right. In particular, it is often used to demonstrate the non-triviality of an *n-cell* in $\mathbb{C}$ by constructing a functor with that domain which maps our cell to a demonstrably non-trivial *n-cell* in its codomain.

The *n-superstructure* functor has a right adjoint $\text{Cos}_n: \omega\text{-}\underline{\text{Cat}} \longrightarrow \omega\text{-}\underline{\text{Cat}}$, we call the ω -category $\text{Cos}_n(\mathbb{C})$ the *n-cosuperstructure* of the ω -category $\mathbb{C}$. This may be constructed as the $(n+1)$ -category whose cells are pairs (c_s, c_t) of *n*-cells in $\mathbb{C}$ satisfying the “globularity” condition that $s_{n-1}(c_s) = s_{n-1}(c_t)$ and $t_{n-1}(c_s) = t_{n-1}(c_t)$. The category structure $(*_m, s_m, t_m)$ on $\text{Cos}_n(\mathbb{C})$ is given point-wise on components when $m < n$, is discrete when $m > n$ and is defined by

$$s_n(c_s, c_t) = (c_s, c_s) \quad t_n(c_s, c_t) = (c_t, c_t) \quad (c'_s, c'_t) *_m (c_s, c_t) = (c'_s, c_t)$$

when $m = n$. By construction, it is clear that we have a canonical isomorphism of ω -categories $|\mathbb{C}|_n \longrightarrow |\text{Cos}_n(\mathbb{C})|_n$, which carries an *n-cell* $c \in \mathbb{C}$ to the *n-cell* $(c, c) \in \text{Cos}_n(\mathbb{C})$. Furthermore, the remaining structure of $\text{Cos}_n(\mathbb{C})$ is completely determined by the fact that each pair of its *n*-cells which satisfy the globularity condition required by the *n*-source and *n*-target of an $(n+1)$ -cell do indeed bound a unique such cell. The adjoint transpose $\hat{f}: \mathbb{C} \longrightarrow \text{Cos}_n(\mathbb{D})$ of an ω -functor $f: |\mathbb{C}|_n \longrightarrow \mathbb{D}$ is given by $\hat{f}(c) = (f(s_n(c)), f(t_n(c)))$, which is well defined since each $c \in \mathbb{C}$ has $s_n(c), t_n(c) \in |\mathbb{C}|_n$ and is easily seen to be appropriately functorial.

As a left adjoint $|\cdot|_n$ preserves all (small) colimits in $\omega\text{-}\underline{\text{Cat}}$. On the other hand, (small) limits in $\omega\text{-}\underline{\text{Cat}}$ are constructed by taking the corresponding limit of underlying sets of cells in $\underline{\text{Set}}$ and defining category structures on this set point-wise, therefore it is also clear that $|\cdot|_n$ preserves all (small) limits. Of course, these preservation properties imply that $|\cdot|_n$ preserves any structure or property which can be described in terms of limits and colimits, such as (regular) monomorphisms and (regular) epimorphisms.

Observation 85 (ω -categories as enriched categories). Recall, from Street [31], that the category $\omega\text{-}\underline{\text{Cat}}$ is cartesian closed and equivalent to the category $(\omega\text{-}\underline{\text{Cat}})\text{-}\underline{\text{Cat}}$ of ω -category enriched categories (in the sense of Kelly [19]). The functor $P: \omega\text{-}\underline{\text{Cat}} \xrightarrow{\sim} (\omega\text{-}\underline{\text{Cat}})\text{-}\underline{\text{Cat}}$ carries an ω -category $\mathbb{C}$ to the ω -category enriched category $P(\mathbb{C})$ with:

- objects u which are the 0-cells of $\mathbb{C}$,
- hom- ω -categories $P(\mathbb{C})(u, v)$ with set of cells $\{c \in \mathbb{C} \mid s_0(c) = u \text{ and } t_0(c) = v\}$ and compositions which are restrictions of the compositions $*_1, *_2, \dots$ of $\mathbb{C}$ to this set,
- composition ω -functor $\circ: P(\mathbb{C})(v, w) \times P(\mathbb{C})(u, v) \longrightarrow P(\mathbb{C})(u, w)$ given by restricting the composition $*_0$ of $\mathbb{C}$, and corresponding identities $\text{id}_u = i_0(u) \in P(\mathbb{C})(u, u)$.

Notice also that this functor restricts to an equivalence $P: (n+1)\text{-}\underline{\text{Cat}} \xrightarrow{\sim} (n\text{-}\underline{\text{Cat}})\text{-}\underline{\text{Cat}}$ for each $n \in \mathbb{N}$.

It is sometimes instructive to think of ω -categories as being *oriented combinatorial CW-complexes* or *globular spaces* and under this analogy it is clearly natural to think of this functor as being a form of *path category construction*.

5. AN INTRODUCTION TO THE DECALAGE CONSTRUCTION

5.1. Nerves and Decalage. First we introduce the nerve construction and the decalage comonad. We also make precise the sense in which the latter is generic for the former.

Observation 86 (comonads and left actions of Δ_+^{op}). First recall, from say [3], that a *comonad* (G, ν, ε) on a category $\mathcal{C}$ is no-more-nor-less than a monoid in the strict monoidal category $([\mathcal{C}, \mathcal{C}]^{\text{op}}, \circ, \text{id}_{\mathcal{C}})$, where $\circ$ denotes the *horizontal* composition of the 2-category $\underline{\text{Cat}}$ and $\text{id}_{\mathcal{C}}$ denotes the identity functor on $\mathcal{C}$. So we know, by lemma 22, that such a comonad corresponds to a unique, strictly monoidal, functor:

$$(\Delta_+^{\text{op}}, \oplus, [-1]) \xrightarrow{\tilde{G}} ([\mathcal{C}, \mathcal{C}], \circ, \text{id}_{\mathcal{C}})$$

However, just as in classical monoid theory, any such representation corresponds, under the adjunction $- \times \mathcal{C} \dashv [\mathcal{C}, *]$, to a unique left action of the monoid Δ_+^{op} on the category $\mathcal{C}$. That is to say, it uniquely determines a functor

$$\Delta_+^{\text{op}} \times \mathcal{C} \xrightarrow{\odot_G} \mathcal{C}$$

which we shall usually write in infix form. This may be given explicitly by $\alpha \odot_G f \stackrel{\text{def}}{=} (\tilde{G}\alpha)f$, for all simplicial operators $\alpha \in \Delta_+^{\text{op}}$ and arrows $f \in \mathcal{C}$, and it satisfies the usual axioms for a monoid left action of the arrows of Δ_+^{op} on those of $\mathcal{C}$:

$$\begin{aligned} \text{id}_{[-1]} \odot_G f &= f && \text{for all arrows } f \in \mathcal{C}, \\ \beta \odot_G (\alpha \odot_G f) &= (\beta \oplus \alpha) \odot_G f && \text{for all arrows } f \in \mathcal{C} \text{ and } \alpha, \beta \in \Delta_+^{\text{op}}. \end{aligned}$$

Classically, it is also common to go a step further and transpose such actions once more, this time under the adjunction $\Delta_+^{\text{op}} \times - \dashv [\Delta_+^{\text{op}}, *]$, to give a functor

$$\mathcal{C} \xrightarrow{\hat{G}} [\Delta_+^{\text{op}}, \mathcal{C}] \tag{10}$$

satisfying rules which are transposes of those satisfied by $\odot_G$. Explicitly, this latter functor maps an object $C \in \mathcal{C}$ to a functor $\hat{G}(C) \in [\Delta_+^{\text{op}}, \mathcal{C}]$, which we might call an *augmented simplicial object in $\mathcal{C}$* , determined by

$$\begin{aligned} \hat{G}(C)([n]) &= G^{n+1}(C) \\ \hat{G}(C)(\delta_i^n) &= G^{n-i}(\varepsilon_{G^i(C)}) && \hat{G}(C)(\sigma_i^n) = G^{n-i}(\nu_{G^i(C)}) \end{aligned}$$

and maps an arrow $f: C \longrightarrow D \in \mathcal{C}$ to an *augmented simplicial map* $\hat{G}(f)$ with components:

$$\hat{G}(f)_n = G^{n+1}(f)$$

Definition 87 (nerves of comonads). Suppose that we are given a category $\mathcal{C}$ equipped with a functor $\Pi: \mathcal{C} \longrightarrow \underline{\text{Set}}$ and a comonad (G, ν, ε) on $\mathcal{C}$. As in observation 86 we may transpose the Δ_+^{op} -action corresponding our comonad to give the functor $\hat{G}$ of display (10). This we compose with the functor

$$[\Delta_+^{\text{op}}, \mathcal{C}] \xrightarrow{[i, \Pi]} [\Delta^{\text{op}}, \underline{\text{Set}}] \stackrel{\text{def}}{=} \underline{\text{Simp}}$$

which post-composes Π onto each $F: \Delta_+^{\text{op}} \longrightarrow \mathcal{C}$ and restricts the domain of the result to Δ^{op} by pre-composing it with the inclusion $i: \Delta^{\text{op}} \hookrightarrow \Delta_+^{\text{op}}$. We call this composite $N_{G, \Pi} \stackrel{\text{def}}{=} [i, \Pi] \circ \hat{G}$ the *nerve functor associated with (G, ν, ε) under $\Pi: \mathcal{C} \longrightarrow \underline{\text{Set}}$* .

Definition 88 (the decalage construction). Using the properties of Δ_+ as a strict monoidal category and the fact that $\underline{\text{Simp}} \stackrel{\text{def}}{=} [\Delta^{\text{op}}, \underline{\text{Set}}]$, we may define a canonical left action of Δ_+^{op} on $\underline{\text{Simp}}$ as follows:

$$\begin{array}{ccc} \Delta_+^{\text{op}} \times \underline{\text{Simp}} & \xrightarrow{\quad \odot \quad} & \underline{\text{Simp}} \\ \begin{array}{ccc} ([n], X) & \xrightarrow{\quad} & X \circ (- \oplus [n]) \\ \alpha \uparrow & \downarrow f & \downarrow f \circ (- \oplus \alpha) \\ ([m], Y) & \xrightarrow{\quad} & Y \circ (- \oplus [m]) \end{array} \end{array} \quad (11)$$

Showing that this is functorial and that it satisfies the axioms required of a left action is a matter of the routine application of the properties of $\oplus$ as the tensor of a strict monoidal category and of $\circ$ as the horizontal composition of the 2-category $\underline{\text{Cat}}$. Appealing to observation 86, we know that this left action corresponds to a unique comonad $(\text{Dec}, \epsilon, \partial)$ on $\underline{\text{Simp}}$, called the *decalage comonad*, which has:

- **underlying functor** given by the “section”:

$$\text{Dec} \stackrel{\text{def}}{=} ([0] \odot -): \underline{\text{Simp}} \longrightarrow \underline{\text{Simp}}$$

- **comultiplication** the natural transformation:

$$\epsilon \stackrel{\text{def}}{=} (\sigma_0^0 \odot -): \text{Dec} = ([0] \odot -) \longrightarrow ([1] \odot -) = \text{Dec} \circ \text{Dec}$$

- **counit** the natural transformation:

$$\partial \stackrel{\text{def}}{=} (\delta_0^0 \odot -): \text{Dec} = ([0] \odot -) \longrightarrow ([-1] \odot -) = \text{id}_{\underline{\text{Simp}}}$$

In a precise sense, the pair consisting of the decalage comonad and the connected components functor constitutes an “identity” for the nerve construction:

Lemma 89 (the nerve functor associated with the decalage comonad). *The nerve functor N_{Dec, Π_0} associated with the decalage comonad $(\text{Dec}, \epsilon, \partial)$ under the connected components functor $\Pi_0: \underline{\text{Simp}} \longrightarrow \underline{\text{Set}}$ of observation 33 is (isomorphic to) the identity functor on $\underline{\text{Simp}}$.*

Proof. We know, from the constructions in definitions 87 and 88, that if X is a simplicial set and $n \in \mathbb{N}$ then $N_{\text{Dec}, \Pi_0}(X)_n \stackrel{\text{def}}{=} \Pi_0([n] \odot X)$, where $\odot$ is the canonical left action defined in display (11). Of course we also know, from observation 33, that $\Pi_0([n] \odot X)$ may be constructed as the coequaliser of the pair

$$([n] \odot X)_1 \begin{array}{c} \xrightarrow{- \cdot \delta_0^1} \\ \xrightarrow{- \cdot \delta_1^1} \end{array} ([n] \odot X)_0$$

which is easily seen, from the definition of $[n] \odot X$ in display (11), to be equal to the pair:

$$X_{n+2} \begin{array}{c} \xrightarrow{- \cdot \delta_{n+1}^{n+2}} \\ \xrightarrow{- \cdot \delta_{n+2}^{n+2}} \end{array} X_{n+1}$$

Notice now that a direct application of the simplicial identities of observation 4 demonstrates that both squares in the following diagram of simplicial operators commute

$$\begin{array}{ccccc}
 [n] & \xrightarrow{\delta_{n+1}^{n+1}} & [n+1] & \xrightarrow{\sigma_n^n} & [n] \\
 \delta_{n+1}^{n+1} \downarrow & & \downarrow \delta_{n+2}^{n+2} & & \downarrow \delta_{n+1}^{n+1} \\
 [n+1] & \xrightarrow{\delta_{n+1}^{n+2}} & [n+2] & \xrightarrow{\sigma_{n+1}^{n+1}} & [n+1]
 \end{array}$$

and that the composites of the upper and lower horizontal pairs of operators evaluate to identities. Applying the simplicial set X to this diagram we get a corresponding diagram in $\mathbf{Set}$ satisfying these commutativity conditions. This demonstrates that the following diagram satisfies the conditions required of it to be a *split coequaliser* (cf. [3])

$$\begin{array}{ccccc}
 & \xleftarrow{-\cdot\sigma_n^{n+1}} & & \xleftarrow{-\cdot\sigma_n^n} & \\
 X_{n+2} & \xrightarrow{-\cdot\delta_{n+1}^{n+2}} & X_{n+1} & \xrightarrow{-\cdot\delta_{n+1}^{n+1}} & X_n \\
 & \xrightarrow{-\cdot\delta_{n+2}^{n+2}} & & &
 \end{array}$$

from which it follows that $N_{\text{Dec}, \Pi_0}(X)_n \stackrel{\text{def}}{=} \Pi_0([n] \odot X) \cong X_n$. It is now a routine matter to demonstrate that these isomorphisms are natural in $[n] \in \Delta^{\text{op}}$, making them the components of a simplicial isomorphism $N_{\text{Dec}, \Pi_0}(X) \cong X$, and in $X \in \mathbf{Simp}$, thus making the whole collection into a natural isomorphism $N_{\text{Dec}, \Pi_0} \cong \text{id}_{\mathbf{Simp}}$ as required. $\square$

5.2. Comonad Transformations and Simplicial Reconstruction.

Observation 90 (the 2-category $\mathbf{Cyl}$ of cylinders). As a step towards studying maps of comonads, we may define a certain 2-category $\mathbf{Cyl}$ whose 0-cells are arbitrary functors and whose hom-category $\mathbf{Cyl}(F, G)$, from one functor $F: \mathcal{C} \longrightarrow \mathcal{C}'$ to another $G: \mathcal{D} \longrightarrow \mathcal{D}'$, is defined to be the **comma** category $[\mathcal{C}, G] \downarrow [F, \mathcal{D}']$ (cf. [3] or [22]) from the post-composite $[\mathcal{C}, G] = G \circ -: [\mathcal{C}, \mathcal{D}] \longrightarrow [\mathcal{C}, \mathcal{D}']$ to the pre-composite $[F, \mathcal{D}'] = - \circ F: [\mathcal{C}', \mathcal{D}'] \longrightarrow [\mathcal{C}, \mathcal{D}']$.

More explicitly, this hom-category has:

- **objects (1-cells)**, called *squares*, (K, K', κ) which consist of a pair of functors $K: \mathcal{C} \longrightarrow \mathcal{D}$ and $K': \mathcal{C}' \longrightarrow \mathcal{D}'$ and a natural transformation $\kappa: G \circ K \Longrightarrow K' \circ F$; as in observation 76 this data is depicted as a *pasting square*

$$\begin{array}{ccc}
 \mathcal{C} & \xrightarrow{K} & \mathcal{D} \\
 F \downarrow & \searrow \kappa & \downarrow G \\
 \mathcal{C}' & \xrightarrow{K'} & \mathcal{D}'
 \end{array}$$

- **arrows (2-cells)**, called *cylinders*, $(\gamma, \gamma'): (K, K', \kappa) \longrightarrow (\bar{K}, \bar{K}', \bar{\kappa})$ consisting of a pair of natural transformations $\gamma: K \Longrightarrow \bar{K}$ and $\gamma': K' \Longrightarrow \bar{K}'$ which satisfy the equation $\bar{\kappa} \cdot (G\gamma) = (\gamma'F) \cdot \kappa$, and
- **vertical composition** of cylinders calculated by applying vertical composition point-wise to the components of each cylinder.

Just as in observation 76 the horizontal composite of a pair of 1-cells in $\underline{Cyl}$ is constructed by horizontal “pasting” of squares or more explicitly:

$$(L, L', \lambda) \circ (K, K', \kappa) \stackrel{\text{def}}{=} (K \circ L, K' \circ L', (L' \kappa) \cdot (\lambda K))$$

Finally, horizontal composition of cylinders is a trivial matter of point-wise composition of their underlying natural transformations.

Of course, pasting squares are familiar to us from our considerations of double categories in subsection 4.3. However, it is worth stressing that in the earlier context our pasting squares comprised dimension 2 of our double categories whereas here we’ve shifted dimensions and they now inhabit dimension 1 of $\underline{Cyl}$.

Observation 91 (cotensors and tensors in $\underline{Cyl}$). In [19] Kelly discusses a primitive kind of enriched weighted colimit construction, called a *cotensor*, which generalises the notion of an iterated product in un-enriched category theory. In the 2-categorical case, if $\underline{A}$ is a 2-category, $\mathcal{E}$ is a category in $\underline{Cat}$ and A' is a 0-cell in $\underline{A}$ then the cotensor $\mathcal{E} \pitchfork A' \in \underline{A}$ of A' by $\mathcal{E}$, if it exists, is the (essentially) unique 0-cell for which there is an isomorphism of categories

$$[\mathcal{E}, \underline{A}(A, A')] \cong \underline{A}(A, \mathcal{E} \pitchfork A')$$

which is 2-natural in $A \in \underline{A}$. For instance, it is a trivial consequence of the way the we defined the enrichment of $\underline{Cat}$ over itself that the cotensor of a category $\mathcal{D} \in \underline{Cat}$ by $\mathcal{E}$ is simply (isomorphic to) the functor category $[\mathcal{E}, \mathcal{D}]$.

We might guess that the cotensor of a 0-cell $G: \mathcal{D} \longrightarrow \mathcal{D}' \in \underline{Cyl}$ in our 2-category of cylinders by the category $\mathcal{E}$ is simply the post-composition functor $[\mathcal{E}, G]: [\mathcal{E}, \mathcal{D}] \longrightarrow [\mathcal{E}, \mathcal{D}']$. To prove this first recall, by the definition of $\underline{Cyl}(F, G)$ as a comma category, that we have a square

$$\begin{array}{ccc} \underline{Cyl}(F, G) & \xrightarrow{\Pi} & [\mathcal{C}, \mathcal{D}] \\ \Pi' \downarrow & \searrow \pi & \downarrow [\mathcal{C}, G] \\ [\mathcal{C}', \mathcal{D}'] & \xrightarrow{[F, \mathcal{D}']} & [\mathcal{C}, \mathcal{D}'] \end{array} \quad (12)$$

in which Π (resp. Π') is the functor which projects cylinders onto their first (resp. second) component and π is the natural transformation with $\pi_{(K, K', \kappa)} \stackrel{\text{def}}{=} \kappa$ for each square $(K, K', \kappa): F \longrightarrow G$. This square is, in fact, 2-universal (in the strict enriched sense of Kelly [19]) amongst all 2-cones of this form over the diagram consisting of the functors $[\mathcal{C}, G]$ and $[F, \mathcal{D}']$. Squares with this property are usually called *comma squares* by 2-categorists.

Taking two copies of this comma square, one for $\underline{Cyl}(F, G)$ (to which we apply the 2-functor $[\mathcal{E}, *]$) and the other for $\underline{Cyl}(F, [\mathcal{E}, G])$, we get a diagram

$$\begin{array}{ccccc} [\mathcal{E}, \underline{Cyl}(F, G)] & \xrightarrow{[\mathcal{E}, \Pi]} & [\mathcal{E}, [\mathcal{C}, \mathcal{D}]] & & \\ \downarrow [\mathcal{E}, \Pi'] & \searrow [\mathcal{E}, \pi] & \downarrow [\mathcal{E}, [\mathcal{C}, G]] & \searrow \cong & \\ [\mathcal{E}, [\mathcal{C}', \mathcal{D}']] & \xrightarrow{[\mathcal{E}, [F, \mathcal{D}']]} & [\mathcal{E}, [\mathcal{C}, \mathcal{D}']] & \xrightarrow{\underline{Cyl}(F, [\mathcal{E}, G])} & [\mathcal{C}, [\mathcal{E}, \mathcal{D}]] \\ & \searrow \cong & \downarrow \Pi' & \searrow \pi & \downarrow [\mathcal{C}, [\mathcal{E}, G]] \\ & & [\mathcal{C}', [\mathcal{E}, \mathcal{D}']] & \xrightarrow{[F, [\mathcal{E}, \mathcal{D}']]} & [\mathcal{C}, [\mathcal{E}, \mathcal{D}']] \end{array} \quad (13)$$

in which the dashed diagonal arrows are canonical isomorphisms derived from the cartesian closed structure on $\underline{Cat}$ and the skewed rectangles that they bound commute by the naturality of those constructions. The existence of the last of these diagonal arrows, depicted with a dotted line, follows from the 2-universal property of the comma square at the bottom right-hand of the diagram and it is the unique arrow making the remaining skewed rectangles commute. However, the square in the upper left hand of the diagram is also a comma square since it is constructed by applying $[\mathcal{E}, *]$, which as a right 2-adjoint preserves all 2-limits, to the comma square associated with $\underline{Cyl}(F, G)$ and so it follows that our dotted arrow is also an isomorphism as required. Proving the appropriate 2-naturality of this isomorphism in F , G and $\mathcal{E}$ is now a matter of routine verification.

For the sake of symmetry, we might also mention the dual concept which generalises iterated sums. The *tensor* of a 0-cell A in a 2-category $\underline{A}$ by a category $\mathcal{E}$, if it exists, is the (essentially) unique 0-cell $\mathcal{E} \otimes A$ for which there is an isomorphism of categories

$$[\mathcal{E}, \underline{A}(A, A')] \cong \underline{A}(\mathcal{E} \otimes A, A')$$

which is 2-natural in $A' \in \underline{A}$. Again, it is practically a tautology that the category $\mathcal{E} \times \mathcal{C}$ is the tensor in $\underline{Cat}$ of $\mathcal{C}$ by $\mathcal{E}$. It is also the case that the tensor of a 0-cell $F: \mathcal{C} \longrightarrow \mathcal{C}' \in \underline{Cyl}$ in our 2-category of cylinders by the category $\mathcal{E}$ is simply the functor $\mathcal{E} \times F: \mathcal{E} \times \mathcal{C} \longrightarrow \mathcal{E} \times \mathcal{C}'$, which applies F “in the second variable”. A proof of this fact may be constructed along the lines of that given for cotensors in $\underline{Cyl}$ above and is left as an exercise for the reader. $\square$

Observation 92 (comonad transformations). Since the comonad notion is entirely equational, we may follow Street [29] and interpret it in any 2-category; in particular we have an interest in studying comonads in $\underline{Cyl}$. If $F: \mathcal{C} \longrightarrow \mathcal{C}'$ is a functor then a comonad on the 0-cell F in $\underline{Cyl}$ is simply a monoid in the (dual) endo-category $\underline{Cyl}(F, F)^{op}$, which is a strict monoidal category under the horizontal composition of $\underline{Cyl}$ in the usual way. Now, since horizontal and vertical composition of cylinders is calculated point-wise, we know that the function which projects cylinders onto their first (respectively second) component provides us with a 2-functor $\Pi: \underline{Cyl} \longrightarrow \underline{Cat}$ (resp. $\Pi': \underline{Cyl} \longrightarrow \underline{Cat}$). It follows that this 2-functor restricts to a strict monoidal functor from $\underline{Cyl}(F, F)^{op}$ to $[\mathcal{C}, \mathcal{C}]^{op}$ (resp. $[\mathcal{C}', \mathcal{C}']^{op}$) and, consequently, that this carries our monoid to a comonad (G, ν, ε) on $\mathcal{C}$ (resp. (G', ν', ε') on $\mathcal{C}'$). The remaining data encapsulated in our monoid in $\underline{Cyl}(F, F)$ simply consists of a natural transformation $\gamma: F \circ G \Longrightarrow G' \circ F$ making (G, G', γ) into an endo-square (endo-1-cell in $\underline{Cyl}$) on F and satisfying the cylinder conditions:

- the pair (ν, ν') is a cylinder from (G, G', γ) to $(G, G', \gamma) \circ (G, G', \gamma)$, in other words we must have $(G'\gamma) \cdot (\gamma G) \cdot (F\nu) = (\nu'F) \cdot \gamma$, and
- the pair $(\varepsilon, \varepsilon')$ is a cylinder from (G, G', γ) to the identity square $(id_{\mathcal{C}}, id_{\mathcal{C}}, id_F)$, equivalently we require that $F\varepsilon = (\varepsilon'F) \cdot \gamma$.

The pair (F, γ) of a functor F and a natural transformation γ satisfying these conditions is known as a *comonad transformation* from (G, ν, ε) to (G', ν', ε') . Such transformations are said to be *strong* if γ is invertible.

Of course we know, from lemma 22, that our comonad transformation (G, G', γ) , as a monoid in $\underline{Cyl}(F, F)^{op}$, corresponds to a unique strict monoidal functor

$$(\Delta_+^{op}, \oplus, [-1]) \xrightarrow{(G, G', \gamma)^\sim} (\underline{Cyl}(F, F), \circ, (id_{\mathcal{C}}, id_{\mathcal{C}'}, id_F))$$

and by applying the isomorphism $[\Delta_+^{\text{op}}, \underline{\text{Cyl}}(F, F)] \cong \underline{\text{Cyl}}(F, [\Delta_+^{\text{op}}, F])$ of observation 91 it follows that this, in turn, corresponds to a square

$$F \xrightarrow{(G, G', \gamma)^\wedge} [\Delta_+^{\text{op}}, F]$$

which, just as in observation 86, satisfies rules that are transposes of those for a left action of Δ_+^{op} on F . Tracing through the constructions of lemma 22 and observations 86 and 91 it is easily seen that this square is of the form

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\hat{G}} & [\Delta_+^{\text{op}}, \mathcal{C}] \\ F \downarrow & \hat{\gamma} \Downarrow & \downarrow [\Delta_+^{\text{op}}, F] \\ \mathcal{C}' & \xrightarrow{\hat{G}'} & [\Delta_+^{\text{op}}, \mathcal{C}'] \end{array} \quad (14)$$

in which the horizontal functors $\hat{G}$ and $\hat{G}'$ are those which correspond to the comonads (G, ν, ε) and (G', ν', ε') (respectively) as in observation 86. At the same time it is easily demonstrated that, for objects $C \in \mathcal{C}$ and $[n] \in \Delta_+^{\text{op}}$, the component $(\hat{\gamma}_C)_n$ of the natural transformation in square (14) is equal to $\gamma_C^{(n+1)}$, where $\gamma^{(n+1)}$ is the natural transformation in the square:

$$(G^{n+1}, (G')^{n+1}, \gamma^{(n+1)}) \stackrel{\text{def}}{=} (G, G', \gamma)^\sim([n])$$

Since this latter square is simply equal to the $(n+1)$ -fold iterated horizontal power of the square (G, G', γ) in $\underline{\text{Cyl}}$, it follows that our original transformation of comonads (F, γ) is **strong** if and only if the corresponding natural transformation $\hat{\gamma}$ of display (14) is an isomorphism. $\square$

Lemma 93. *Suppose that we are given a categories $\mathcal{C}$ and $\mathcal{C}'$, along with*

- *comonads (G, ν, ε) on $\mathcal{C}$ and (G', ν', ε') on $\mathcal{C}'$,*
- *a strong comonad transformation (F, γ) from the (G, ν, ε) to (G', ν', ε') , and*
- *functors $\Pi: \mathcal{C} \longrightarrow \underline{\text{Set}}$ and $\Pi': \mathcal{C}' \longrightarrow \underline{\text{Set}}$ along with an isomorphism*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{C}' \\ \Pi \searrow & \xrightarrow[\alpha]{\cong} & \swarrow \Pi' \\ & \underline{\text{Set}} & \end{array}$$

then we may construct a natural isomorphism

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{C}' \\ N_{G, \Pi} \searrow & \xrightarrow[N_{\gamma, \alpha}]{\cong} & \swarrow N_{G', \Pi'} \\ & \underline{\text{Simp}} & \end{array}$$

in which the diagonals are the nerve functors associated with our comonads as in definition 87.

Proof. Consider the following diagram

$$\begin{array}{ccccc}
 \mathcal{C} & \xrightarrow{\hat{G}} & [\Delta_+^{\text{op}}, \mathcal{C}] & \xrightarrow{[\Delta_+^{\text{op}}, \Pi]} & [\Delta_+^{\text{op}}, \underline{\text{Set}}] \xrightarrow{[i, \underline{\text{Set}}]} \underline{\text{Simp}} \\
 \downarrow F & \hat{\gamma} \Downarrow \cong & \downarrow [\Delta_+^{\text{op}}, F] & \downarrow [\Delta_+^{\text{op}}, \alpha] \Downarrow \cong & \\
 \mathcal{C}' & \xrightarrow{\hat{G}'} & [\Delta_+^{\text{op}}, \mathcal{C}'] & \xrightarrow{[\Delta_+^{\text{op}}, \Pi']} &
 \end{array}$$

in which

- the left hand square is that derived from the strong comonad transformation (F, γ) as in observation 92,
- the upper composite of functors (from $\mathcal{C}$ to $\underline{\text{Simp}}$) is, by definition, the nerve functor associated with (G, ν, ε) under Π .
- the lower composite of functors (from $\mathcal{C}'$ to $\underline{\text{Simp}}$) is, by definition, the nerve functor associated with (G', ν', ε') under Π' .

It follows that the natural transformation obtained by pasting this diagram provides us with the isomorphism required in the statement of the lemma. $\square$

The following formalises a common application of the theory of decalage:

Corollary 94 (simplicial reconstruction). *Suppose that we are given a category $\mathcal{C}$, along with*

- *a comonad (G, ν, ε) on $\mathcal{C}$,*
- *a strong comonad transformation (F, γ) from the decalage comonad $(\text{Dec}, \epsilon, \partial)$ on $\underline{\text{Simp}}$ to (G, ν, ε) , and*
- *a functor $\Pi: \mathcal{C} \longrightarrow \underline{\text{Set}}$ along with an isomorphism*

$$\begin{array}{ccc}
 \underline{\text{Simp}} & \xrightarrow{F} & \mathcal{C} \\
 \Pi_0 \searrow & \xrightarrow[\alpha]{\cong} & \swarrow \Pi \\
 & \underline{\text{Set}} &
 \end{array}$$

then the nerve functor $N_{G, \Pi}: \mathcal{C} \longrightarrow \underline{\text{Simp}}$ associated with (G, ν, ε) under Π is a left pseudo-inverse of $F: \underline{\text{Simp}} \longrightarrow \mathcal{C}$. In other words, there exists a natural isomorphism $N_{G, \Pi} \circ F \cong \text{id}_{\underline{\text{Simp}}}$.

Proof. Simply apply the previous lemma to the data given in the statement to obtain a natural isomorphism

$$\begin{array}{ccc}
 \underline{\text{Simp}} & \xrightarrow{F} & \mathcal{C} \\
 N_{\text{Dec}, \Pi_0} \searrow & \xrightarrow[\cong]{} & \swarrow N_{G, \Pi} \\
 & \underline{\text{Simp}} &
 \end{array}$$

and observe, by lemma 89, that the nerve functor N_{Dec, Π_0} is isomorphic to the identity functor on $\underline{\text{Simp}}$. $\square$

In other words, this lemma simply shows that under the conditions of its statement we may use the comonad (G, ν, ε) to reconstruct each simplicial set X from the object $F(X) \in \mathcal{C}$.

Observation 95 (semi-simplicial versions). In this paper we will apply a version of the simplicial reconstruction lemma above, but it will suffice for us to reconstruct semi-simplicial structures only. Of course, all of the theory in this appendix may be restricted to consider only semi-simplicial sets, copointed endo-functors (rather than comonads), their transformations (appropriately defined) and the corresponding subcategory Δ_{f+} (in the place of Δ_+). Proofs of these semi-simplicial results remain substantially unchanged from their simplicial counterparts discussed above, so we leave the (trivial) details up to the reader.

6. STRATIFICATIONS AND FILTERINGS OF SIMPLICIAL SETS

6.1. Stratified Simplicial Sets.

Definition 96 (stratified simplicial sets). A *stratified simplicial set* (or sometimes just a *stratified set*) is a pair (X, tX) where X is a simplicial set and tX is a subset¹ of its simplices, which are said to be *thin*, satisfying the conditions that

- no 0-simplex of X is in tX ,
- all of the degenerate simplices of X are in tX .

A *stratified simplicial map* (or sometimes just a *stratified map*) $f: (X, tX) \longrightarrow (Y, tY)$ is a simplicial map $f: X \longrightarrow Y$ such that $tX \subseteq f^{-1}(tY)$. In other words, a stratified simplicial map is a simplicial map that *preserves thinness*.

The composite of two stratified simplicial maps again preserves thinness, so we may collect stratified simplicial sets and maps together into a category Strat.

We will often drop explicit mention of the set of thin simplices tX and elect to notationally confuse a stratified set with its underlying simplicial set by simply declaring that X (rather than (X, tX)) is a stratified set. In such cases, the set of thin simplices associated with the stratified sets $X, Y, Z, \dots$ will always be denoted by $tX, tY, tZ, \dots$ respectively.

Observation 97 (unity and identity of opposites). The forgetful functor $U: \text{Strat} \longrightarrow \text{Simp}$, which takes (X, tX) to X , admits both right **and** left adjoints. The right adjoint takes a simplicial set X to $R(X) = (X, X \setminus X_0)$, in which all simplices of dimension > 0 are thin, whereas the left adjoint takes X to $L(X) = (X, X_d)$ where $X_d = \{x \in X \mid x \text{ is degenerate}\}$, making as few simplices thin as possible. Both of the functors R, L are fully faithful, so this sequence of adjoints is a *unity and identity of opposites* (a situation studied in general by F. W. Lawvere).

In general, we will identify Simp with its image under L , that is the full subcategory of Strat of those stratified sets such that every thin simplex is degenerate. In particular, we will assume, without comment, that the standard simplicial sets such as $\Delta[n]$ and $\Lambda_k[n]$ also live in Strat.

Observation 98 (duals of stratified simplicial sets). The dual $(X, tX)^\circ$ of a stratified simplicial set (X, tX) is the pair (X°, tX°) where X° is the dual simplicial set introduced in observation 38 and $tX^\circ \stackrel{\text{def}}{=} tX$.² Quite clearly this pair constitutes a well defined stratified simplicial set.

If $f: (X, tX) \longrightarrow (Y, tY)$ is a stratified simplicial map then the dual simplicial map $f^\circ: X^\circ \longrightarrow Y^\circ$ carries tX° into tY° . It follows that f° is a stratified map from $(X, tX)^\circ$ to $(Y, tY)^\circ$ and that we may extend our dual operation to a functor $(-)^\circ: \text{Strat} \longrightarrow \text{Strat}$ which makes the diagram of functors

$$\begin{array}{ccc} \text{Strat} & \xrightarrow{(-)^\circ} & \text{Strat} \\ U \downarrow & & \downarrow U \\ \text{Simp} & \xrightarrow{(-)^\circ} & \text{Simp} \end{array}$$

¹Note that tX is merely a subset of X , **not** a simplicial subset, in general it will not be closed in X under the action of Δ .

²It is often quite useful to distinguish tX and tX° notationally, even though they are identical as sets.

commute. Yet again this dual functor is strictly involutive, in that its composite with itself is the identity functor on Strat.

Notation 99 (stratified simplicial subsets). A pair (Y, tY) is a *stratified simplicial subset* (or just a *stratified subset*) of a stratified set (X, tX) , denoted $(Y, tY) \subseteq_s (X, tX)$, if:

- Y is a simplicial subset of X ,
- tY contains all of the degenerate simplices of Y , and
- tY is a subset of tX .

It follows from this definition that (Y, tY) inherits a stratified simplicial structure from (X, tX) and that the inclusion $Y \subseteq_s X$ becomes a stratified map $(Y, tY) \xrightarrow{\subseteq_s} (X, tX)$.

Intersections and unions of stratified subsets $(Y_i, tY_i) \subseteq_s (X, tX)$ are given by the formulae:

$$\bigcup_i (Y_i, tY_i) \stackrel{\text{def}}{=} \left(\bigcup_i Y_i, \bigcup_i tY_i \right) \quad \bigcap_i (Y_i, tY_i) \stackrel{\text{def}}{=} \left(\bigcap_i Y_i, \bigcap_i tY_i \right)$$

Notation 100 (regular, entire and inclusive stratified maps). We say that a stratified map $f: X \longrightarrow Y$ is

- **regular** if it *reflects thin simplices*, which means that whenever $f(x)$ is thin in Y it follows that x is thin in X . In other words, f is regular iff $f^{-1}(tY) = tX$.
- **entire** if it is surjective on simplices, that is to say if its underlying simplicial map is surjective.
- **an inclusion** if it is injective on simplices, that is to say if its underlying simplicial map is injective.

Sometimes we will adopt notation $\mathcal{R}$, $\mathcal{E}$ and $\mathcal{I}$ to denote the classes of regular, entire and inclusive stratified maps respectively.

Notice that a stratified map is entire (resp. an inclusion) if and only if it is an epimorphism (resp. a monomorphism) in the usual categorical sense. Furthermore, observe that our stratified map $f: X \longrightarrow Y$ is a stratified isomorphism (that is an isomorphism in the category Strat) if and only if it is both bijective on simplices and regular. In other words, the class of stratified isomorphisms coincides with the intersection $\mathcal{E} \cap \mathcal{I} \cap \mathcal{R}$ of the three classes defined above.

Observation 101 (stratified images and inverse images). If $f: X \longrightarrow Y$ is a stratified simplicial map then define

- the *image* of a stratified subset $X' \subseteq_s X$ under f to be the stratified subset given by $f(X') \stackrel{\text{def}}{=} (f(X'), f(tX'))$, and
- the *inverse image* of a stratified subset $Y' \subseteq_s Y$ under f to be the stratified subset given by $f^{-1}(Y') \stackrel{\text{def}}{=} (f^{-1}(Y'), f^{-1}(tY'))$.

We leave it up to the reader to verify these definitions do indeed specify well defined stratified simplicial subsets. Observe that:

- for any stratified map $f: X \longrightarrow Y$ the inverse image $f^{-1}(Y')$ of a regular subset $Y' \subseteq_r Y$ is a regular subset of X ,
- for any **regular** stratified map $f: X \longrightarrow Y$ the image $f(X')$ of a regular subset $X' \subseteq_r X$ is again a regular subset of Y .

Notation 102 (regular and entire stratified subsets). We say that $X \subseteq_s Y$ is:

- *regular*, and use the notation $X \subseteq_r Y$, if the associated inclusion map $X \xrightarrow{\subseteq_s} Y$ is regular or equivalently when $tX = X \cap tY$.

- *entire*, and use the notation $X \subseteq_e Y$, if the associated inclusion map $X \xrightarrow{\subseteq_s} Y$ is entire or equivalently when X and Y have the same underlying simplicial set.

We define S^* , the regular stratified subset *generated* by a subset $S \subseteq X$, to be the smallest regular stratified subset of X containing S and clearly $A^* = (A^*, tX \cap A^*)$.

Notice that the classes $\mathcal{R}$, $\mathcal{E}$ and $\mathcal{I}$ provide us with two distinct factorisation systems on Strat (in the sense of Freyd and Kelly [11]):

- $(\mathcal{E}, \mathcal{R} \cap \mathcal{I})$ which factorises a stratified map $f: X \longrightarrow Y$ as $X \xrightarrow{f_e} \text{im}_r(f) \xrightarrow{\subseteq_r} Y$, where $\text{im}_r(f) \subseteq_r Y$ is the *regular image* of f , which is the regular stratified subset of Y generated by $\{f(x) \in Y \mid x \in X\}$.
- $(\mathcal{E} \cap \mathcal{I}, \mathcal{R})$ which factorises a stratified map $f: X \longrightarrow Y$ as $X \xrightarrow{\subseteq_e} \text{coim}_e(f) \xrightarrow{f_r} Y$, where $X \subseteq_e \text{coim}_e(f)$ is the *entire coimage* of f , which has the same underlying simplicial set as X and thin simplices those $x \in X$ with $f(x)$ thin in Y .

Notation 103 (lifting and extension). Suppose that $f: X \longrightarrow Y$ is a stratified map:

- if Y' is a regular subset of Y then we say that f *lifts* to a stratified map with codomain Y' iff the simplex $f(x)$ is in Y' for all $x \in X$, in which case we can factor f through $Y' \subseteq_r Y$ and consider it to be a stratified map $f: X \longrightarrow Y'$.
- if X' is an entire superset of X (i.e. $X \subseteq_e X'$) then we say that f *extends* to a stratified map with domain X' iff the simplex $f(x)$ is thin in Y for all thin simplices x in X' , in which case we can factor it through $X \subseteq_e X'$ and consider it to be a stratified map $f: X' \longrightarrow Y$.

Definition 104. Let Simplex denote the full subcategory of Strat consisting of those stratified sets whose underlying simplicial sets is a standard simplex $\Delta[n]$ for some $n \in \mathbb{N}$.

We'll often adopt the alphabetic convention of assuming that the uppercase letter used to denote a given stratified set in Simplex matches the lower case letter used to denote the dimension of its underlying simplex. In other words, under this convention the stratified sets N, M, P and Q in Simplex would have underlying simplicial sets $\Delta[n]$, $\Delta[m]$, $\Delta[p]$ and $\Delta[q]$ respectively whereas N, N' and N_i in Simplex would all have underlying simplicial set $\Delta[n]$.

Notation 105 (some standard stratified simplicial sets). A few stratified simplicial sets of note are:

- (i) **The standard thin n -simplex** $\Delta[n]_t$, which is obtained from $\Delta[n]$ by making its only non-degenerate n -simplex $\text{id}_{[n]}: [n] \longrightarrow [n]$ thin.
- (ii) **The k^{th} standard admissible n -simplex** $\Delta_k^a[n]$ (for $n \geq 2, k = 1, \dots, n-1$) is obtained from $\Delta[n]$ by making all (non-degenerate) simplices $\alpha: [m] \longrightarrow [n]$ with $k-1, k, k+1 \in \text{im}(\alpha)$ thin. These are particularly important in characterising the nerves of n -categories.
- (iii) **The standard admissible $(n-1)$ -dimensional k -horn** $\Lambda_k^a[n]$ (for $n \geq 2, k = 1, \dots, n-1$) is obtained from $\Lambda_k[n]$ by making any simplex which is thin in $\Delta_k[n]$ thin in $\Lambda_k^a[n]$. In other words, $\Lambda_k^a[n]$ is the regular stratified subset of $\Delta_k[n]$ generated by the set of simplices $\{\delta_i^n \mid i = 0, \dots, k-1, k+1, \dots, n\}$.

The non-degenerate simplices of $\Delta[n]$ satisfying the condition given in example (ii) are said to be *k-divided*. It should be noted that this use of the term differs slightly from that used by Street in [31], a difference we discuss a little more in theorem 249.

Observation 106 (admissible simplices and horns in a stratified simplicial set). An n -simplex x in a stratified simplicial set X is said to be k -admissible if the corresponding simplicial map $\lceil x \rceil: \Delta[n] \longrightarrow X$ (under Yoneda's lemma) extends to a stratified simplicial map $\lceil x \rceil: \Delta_k^a[n] \longrightarrow X$.

By the definition of $\lceil x \rceil$, that is $\lceil x \rceil(\alpha) \stackrel{\text{def}}{=} x \cdot \alpha$, this holds if and only if $x \cdot \alpha$ is thin in X for all k -divided simplices $\alpha \in \Delta[n]$.

Similarly, we say that an $(n-1)$ -dimensional k -horn $\{x_i \in X_{n-1} \mid i = 0, 1, \dots, k-1, k+1, k+2, \dots, n\}$ in X is *admissible* if the corresponding simplicial map $\lceil \cdot \rceil: \Lambda_k[n] \longrightarrow X$ extends to a stratified map from $\Lambda_k^a[n]$ to X . By Yoneda's lemma, this condition holds if and only if

- x_i is $(k-1)$ -admissible in X for each $i < k-1$, and
- x_i is k -admissible in X for each $i > k+1$.

Observation 107 (duals of standard stratified simplicial sets). The dual operation on Δ gives rise to a canonical simplicial isomorphism:

$$\begin{array}{ccc} \Delta[n] & \xrightarrow{(-)^\circ} & \Delta[n]^\circ \\ \alpha & \longmapsto & \alpha^\circ \end{array}$$

This extends to a family of stratified simplicial isomorphisms between the standard stratified sets introduced in the last observation and their duals $\Delta[n]_t \cong \Delta[n]_t^\circ$ and $\Delta_{n-k}^a[n] \cong \Delta_k^a[n]^\circ$. Furthermore this latter isomorphism restricts to an isomorphism $\Lambda_{n-k}^a[n] \cong \Lambda_k^a[n]^\circ$ of admissible horns and their duals.

Observation 108 (Strat as an LFP-category). The small full subcategory $t\Delta$ on the set of objects $\{\Delta[n], \Delta[n]_t \mid n = 0, 1, \dots\}$ in Strat gives rise to a right adjoint functor

$$\text{Strat} \xleftarrow[\perp]{\quad} [t\Delta^{\text{op}}, \text{Set}] \quad (15)$$

by Kan's construction (observation 63).

Applying Yoneda's lemma, we may describe $t\Delta$ as an extension of Δ in the way originally proposed by Street in [31]. Under this description $t\Delta$ has two families of objects, the original family of finite ordinals $\{[n] \mid n \geq 0\}$ of Δ and a family of copies of the non-zero ordinals $\{[n]_t \mid n > 0\}$. In other words $[n]$ and $[n]_t$ are distinct copies of the finite ordinal with $n+1$ elements, where the former copy represents the standard n -simplex and the latter is a model for the standard thin n -simplex. In the sequel, we will sometimes also use the notation $[n]_?$ if we wish to refer to either of the objects $[n]$ or $[n]_t$. The arrows of $t\Delta$ are simply order preserving maps subject to a restriction which excludes all non-identity face operators (injective maps) whose domains are of the form $[n]_t$.

This category may be presented in terms of generators which include the elementary face and degeneracy operators of Δ , the elementary degeneracies $\zeta_i^n: [n+1]_t \longrightarrow [n]$ and the operator $\varrho^n: [n] \longrightarrow [n]_t$ that is the unique order isomorphism between the ordinal $[n]$ and its copy $[n]_t$. The relations that must hold between these operators are those of Δ plus the extra family of relations $\sigma_i^n = \zeta_i^n \circ \varrho^{n+1}$ (for $n \in \mathbb{N}$).

In line with our notation for stratified standard simplices, we will let $\Delta: t\Delta \longrightarrow \text{Strat}$ denote the Yoneda functor which embeds $t\Delta$ (under this alternative presentation) into Strat. In particular, this means that $\Delta(\varrho^n): \Delta[n] \longrightarrow \Delta[n]_t$ and $\Delta(\zeta_i^n): \Delta[n+1]_t \longrightarrow \Delta[n]$ denote the stratified maps whose underlying simplicial maps are the identity on $\Delta[n]$ and the map $\Delta(\sigma_i^n): \Delta[n+1] \longrightarrow \Delta[n]$ respectively.

Using a Yoneda argument we may demonstrate that the right adjoint functor in display (15) is fully faithful, or in other words that $t\Delta$ is embedded in $\mathbf{Strat}$ as a small *dense* subcategory. Consequently, we will identify $\mathbf{Strat}$ with the reflective full subcategory of $[t\Delta^{\text{op}}, \mathbf{Set}]$ which is the replete image of this right adjoint functor and use the notation $L_s: [t\Delta^{\text{op}}, \mathbf{Set}] \longrightarrow [t\Delta^{\text{op}}, \mathbf{Set}]$ for the corresponding reflector. Notice also that an object F of $[t\Delta^{\text{op}}, \mathbf{Set}]$ is in the full subcategory $\mathbf{Strat}$ if and only if the function $F(\varrho^n): F([n]_t) \longrightarrow F([n])$ is injective for each $n > 0$. This condition may equivalently be expressed as stating that the pullback of each $F(\varrho^n)$ along itself should be the identity on $F([n]_t)$ which in turn may be written as a perpendicularity condition with respect to certain maps between finitely presented objects in $[t\Delta^{\text{op}}, \mathbf{Set}]$. It follows, therefore, that $\mathbf{Strat}$ may be described as the LFP-category of algebras of an LE-theory $\mathbb{T}_{\mathbf{Strat}}$ with underlying category $t\Delta$ and that the associated reflector L_s is finitely accessible (see subsection 3.2).

Of course, now observation 59 may be applied to show that $\mathbf{Strat}$ is closed in $[t\Delta^{\text{op}}, \mathbf{Set}]$ under limits and filtered colimits, with all other colimits in $\mathbf{Strat}$ being constructed by forming them point-wise in $[t\Delta^{\text{op}}, \mathbf{Set}]$ and then reflecting them back into $\mathbf{Strat}$ using L_s . Furthermore, consulting definition 56 we also see that every stratified simplicial set X may be expressed in $\mathbf{Strat}$ as the canonical colimit $\text{colim}(X, \Delta)$ of standard simplices and standard thin simplices weighted by X (regarded as an object in $[t\Delta^{\text{op}}, \mathbf{Set}]$). Alternatively, as discussed in the text accompanying definition 56, we may describe this colimit in terms of a traditional conical colimit by constructing the Grothendieck category $\mathbb{G}(X) \stackrel{\text{def}}{=} t\Delta \downarrow X$, forming the diagram $D_X: \mathbb{G}(X) \longrightarrow \mathbf{Strat}$ by domain projection and using a Yoneda argument to show that the obvious cocone $i: D_X \longrightarrow X$ with components $i_f = f: D_X(f) = \Delta[n]_? \longrightarrow X$ induces a canonical isomorphism $X \cong \text{colim}_{\mathbb{G}(X)}(D_X)$.

If X is a stratified set then the category $\mathbb{G}(X)$ has a *cofinal* set of objects consisting of those stratified maps $\lceil x \rceil: \Delta[n]_? \longrightarrow X$ with $[n]_? \in t\Delta$ which correspond under Yoneda's lemma to simplices $x \in X$ which are non-degenerate. It follows that if X has only a finite set of non-degenerate simplices then it may be expressed as a **finite** colimit of standard (thin) simplices. However, we know that every standard (thin) simplex is finitely presentable and that any finite colimit of FP-stratified sets is also finitely presentable, so we may infer that any stratified set with only a finite set of non-degenerate simplices is finitely presentable. Indeed the reverse implication also holds, but we won't use it here and we leave its proof up to the reader.

Observation 109 (limits and colimits in $\mathbf{Strat}$). If $D: \mathbb{C} \longrightarrow \mathbf{Strat}$ is a diagram of stratified simplicial sets, where $\mathbb{C}$ is a small category, then the limit and colimit of D may be obtained via the following explicit constructions:

Limits First form the limit $\lim(U \circ D)$ in $\mathbf{Simp}$, as described in observation 31, then raise this to a stratified simplicial set by making thin those simplices $\{x_i \in D(i)\}_{i \in \text{obj}(\mathbb{C})}$ for which x_i is thin in $D(i)$ for each $i \in \text{obj}(\mathbb{C})$.

Colimits Again start by forming the colimit $\text{colim}(U \circ D)$ in $\mathbf{Simp}$, cf. observation 31, then raise this to $\mathbf{Strat}$ by making thin those simplices that can be written as an equivalence class $[x, i]$ with x thin in $D(i)$.

Given these descriptions, it is clear that many results about limits and colimits in $\mathbf{Simp}$ may be routinely raised to $\mathbf{Strat}$, simply by appealing to the result on underlying simplicial sets and then checking that stratifications are appropriately respected. In particular, it is immediate that the result of observation 32 also holds for unions of stratified subsets.

Observation 110 (skeleta and coskeleta of stratified sets). Fixing $n \in \mathbb{N}$, we say that a stratified set X is *n-skeletal* iff each simplex $x \in X$ with dimension $\dim(x) > n$ is degenerate. The full subcategory of *n-skeletal* stratified sets is coreflective in Strat, with coreflector $\|\cdot\|_n : \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ (cf. observation 49) which maps a stratified set X to the regular stratified subset $\|X\|_n \subseteq_r X$, called its *n-skeleton*, of those simplices $x \in X$ with $x = x' \cdot \alpha$ for some x' with $\dim(x') \leq n$. In other words, this is the unique largest *n-skeletal* regular subset of X .

This *n-skeleton* functor has a right adjoint $\text{Cosk}_n : \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}} \subseteq [t\Delta^{\text{op}}, \underline{\text{Set}}]$ formed by applying Kan's construction (observation 63) to the functor $\|\Delta(\cdot)\|_n : t\Delta \longrightarrow \underline{\text{Strat}}$. Equivalently, applying a Yoneda argument we see that $\text{Cosk}_n(X)$, the *n-coskeleton* of the stratified set Strat, may be described as the stratified set whose simplices, operator actions and stratification coincide with that of X at and below dimension n and which is completely determined at each dimension r above n by the fact that each one of its $(r-1)$ -dimensional cycles is the boundary of a unique thin r -simplex.

If we let $t\Delta_n$ denote the full subcategory of $t\Delta$ on those standard (thin) simplices of dimension $\leq n$ then it is an immediate consequence of the Eilenberg-Zilber lemma that $t\Delta_n$ is dense in the full subcategory of *n-skeletal* stratified sets in Strat. It follows that we may use the Grothendieck construction, just as we did in observation 108, to express the *n-skeleton* of a stratified set X as a canonical colimit of standard (thin) simplices of dimension $\leq n$. Adopting the notation $\mathbb{G}_n(X)$ for the comma category $t\Delta_n \downarrow X$ and D_X^n for the canonical diagram obtained by “codomain projection” then we find that each leg of the canonical cocone from this diagram into X restricts to a map into the regular subset $\|X\|_n$ and provides us with a colimiting cocone which induces a canonical isomorphism $\text{colim}(D_X^n) \cong \|X\|_n$.

Observation 111 (connected components of stratified sets). Under our identification of Simp with the subcategory of minimally stratified sets in Strat, we may extend the adjunction $\Pi_0 \dashv \text{dis} : \underline{\text{Set}} \longrightarrow \underline{\text{Simp}}$ of observation 33 to one between Set and the category of stratified sets Strat. If X is a stratified set then its set of connected components $\Pi_0(X)$ is simply formed by applying the simplicial connected components functor to the underlying simplicial set $U(X)$. As before, it is clear that $\text{dis} : \underline{\text{Set}} \longrightarrow \underline{\text{Strat}}$ is fully faithful and that a stratified set is in its replete image iff it is 0-skeletal. It follows that dis provides an equivalence between Set and the full subcategory of 0-skeletal stratified sets in Strat.

It follows that the category of 0-skeletal stratified sets is also a **reflective** full subcategory of Strat with corresponding reflector $L_0 = \text{dis} \circ \Pi_0$. A stratified map $f : X \longrightarrow Y$ is L_0 -invertible (cf. definition 50) iff the function $\Pi_0(f) : \Pi_0(X) \longrightarrow \Pi_0(Y)$ is an isomorphism, consequently we say that such a map is *bijective on components*. We also say that X is *connected* if $\Pi_0(X)$ is the singleton set. From the description of Π_0 above, it follows immediately that:

- The set of connected components of a stratified set is dependent only upon the structure of its underlying simplicial set.
- All of the standard simplices and horns are connected.
- The connected components functor Π_0 preserves products and it follows that products of connected sets are connected.

Furthermore, it is worth mentioning that any stratified map $f : X \longrightarrow Y$ whose domain and codomain are connected is bijective on components, since any function between the singleton sets $\Pi_0(X)$ and $\Pi_0(Y)$ must be a bijection.

6.2. Superstructures and Filtered Semi-Simplicial Sets.

Definition 112 (superstructures of stratified sets). For each $n \in \mathbb{N}$ there exists a pair of adjoint functors

$$\begin{array}{ccc} & \text{Th}_n & \\ \text{Strat} & \begin{array}{c} \xleftarrow{\quad} \\ \perp \\ \xrightarrow{\quad} \end{array} & \text{Strat} \\ & |\cdot|_n & \end{array}$$

which are defined as follows:

- $\text{Th}_n(\mathbf{X})$ is defined to be $(X, tX \cup \{x \in X \mid \dim(x) > n\})$, in other words it is the stratified simplicial set obtained by making thin all of the simplices of X with dimension $> n$. It is clear that any stratified map $f: X \longrightarrow Y$ extends to a stratified map from $\text{Th}_n(X)$ to $\text{Th}_n(Y)$, so we let $\text{Th}_n(f) \stackrel{\text{def}}{=} f$.
- $|X|_n$, called the n -dimensional superstructure of X is its regular stratified subset with underlying simplicial set given by:

$$\{x \in X \mid (\forall \alpha: [r] \longrightarrow [\dim(x)] \in \Delta) \ r > n \Rightarrow x \cdot \alpha \in tX\}$$

In other words, $|X|_n$ is the regular subset of those simplices of X for which each r -dimensional face with $r > n$ is thin. We may easily verify that a stratified map $f: X \longrightarrow Y$ has $|X|_n \subseteq_s f^{-1}(|Y|_n)$ and thus restricts to a stratified map $|f|_n: |X|_n \longrightarrow |Y|_n$.

We say that a stratified set X is n -trivial if and only if $X = |X|_n$ (or equivalently iff $X = \text{Th}_n(X)$) and we use the notation Strat_n to denote the full subcategory of these in Strat . In other words, X is n -trivial if and only if all of its simplices of dimension greater than n are thin.

Observation 113. The following properties of the superstructures of a stratified set X are worthy of note.

- (1) If $m \leq n$ then $|X|_m$ is a regular subset of $|X|_n$.
- (2) If $m < r$ and x is an m -simplex of X then all of the r -dimensional faces of x are degenerate, and thus thin, in X . It follows that $|X|_n$ contains **all** of the simplices of X with dimension $\leq n$.
- (3) If x is an n -simplex of X then the only non-degenerate n -dimensional face of x is x itself, so it follows that this is an element of $|X|_{n-1}$ if and only if it is thin in X .

In particular, the second of these observations immediately implies that X is equal to the union $\bigcup_{n \in \mathbb{N}} |X|_n$ of its superstructures.

Definition 114 (categories of filtered objects). The partially ordered set of natural numbers $\mathbb{N}$ may be considered to be a category in the usual way. That is to say we think of it as a category with objects non-negative integers and a unique arrow $i \longrightarrow j$ for each pair of integers with $i \leq j$. If $\mathcal{C}$ is any category then we call the functor category $[\mathbb{N}, \mathcal{C}]$ the *category of filtered objects* in $\mathcal{C}$.

Observation 115. The family of superstructures of X provides us with a filtered family of stratified sets, since we have $|X|_i \subseteq_r |X|_{i+1}$ for each $i \in \mathbb{N}$. Now, if we throw away stratifications and forget about the actions of degeneracy operators we may consider this to be a filtered family of semi-simplicial sets. It follows that we may gather this information

together to provide a single functor

$$\mathbf{Strat} \xrightarrow{|\cdot|} [\mathbb{N}, \mathbf{SSimp}]$$

which represents each stratified set as a filtered semi-simplicial set.

It is not, in general, the case that this functor is fully faithful, however it is possible to find a substantial full subcategory of $\mathbf{Strat}$ on which it is. In the remainder of this section we establish a characterisation of this subcategory and in section 8 we demonstrate that it contains all of those stratified sets which Street ([31] and [32]) identified as being candidates for those in the replete image of his ω -categorical nerve functor.

Definition 116. We say that a simplex x in a stratified set X is *pre-degenerate at k* (where $0 \leq k < \dim(x)$) if $x \cdot \alpha$ is thin in X for each face operator α for which $k, k+1 \in \text{im}(\alpha)$.

In particular, observation 29(c) and the fact that in a stratified set all degenerate simplices are thin implies that every simplex $x \in X$ which is degenerate at k is pre-degenerate at k . The converse is true for some stratified sets, including those that occur as the nerves of ω -categories.

Definition 117. If X is a stratified simplicial set then we say that it is *well tempered* iff any simplex $x \in X$ which is pre-degenerate at some k is in fact degenerate at k . In other words, X is well tempered if its collection of thin simplices is sufficient to detect those simplices which are degenerate.

We use the notation $\mathbf{Strat}_W$ to denote the full subcategory of $\mathbf{Strat}$ on those stratified sets which are well tempered.

Our primary motivation for introducing these definitions is:

Lemma 118. *Suppose that Y is a well tempered stratified set and $f: X \longrightarrow Y$ is a semi-simplicial map which preserves thinness then f is a stratified map.*

Proof. First recall that every simplicial operator factors as a face operator following a degeneracy operator and that this latter operator, in turn, factors as a composite of elementary degeneracy operators. However, we know that f preserves the actions of face operators (since it is a semi-simplicial map) so if we can show that it also preserves the action of elementary degeneracy operators then it preserves the actions of all operators and is thus a simplicial map.

Now the notion of pre-degeneracy is defined in terms of actions of **face** operators and thinness, thus it follows that a map like f which preserves both of these structures will also preserve pre-degeneracy. So suppose that we are given an n -simplex $x \in X$ and an elementary degeneracy operator $\sigma_k^n: [n+1] \longrightarrow [n]$, then we know that $x \cdot \sigma_k^n$ is pre-degenerate at k and so we may infer that $f(x \cdot \sigma_k^n) \in Y$ is also pre-degenerate at k . But Y is well tempered and so it follows that $f(x \cdot \sigma_k^n)$ is actually degenerate at k .

Finally we may demonstrate that f preserves the action of σ_k^n on x using the following simple calculation:

$$\begin{aligned} f(x \cdot \sigma_k^n) &= (f(x \cdot \sigma_k^n) \cdot \delta_k^{n+1}) \cdot \sigma_k^n && \text{since } f(x \cdot \sigma_k^n) \text{ is degenerate at } k, \\ &= f((x \cdot \sigma_k^n) \cdot \delta_k^{n+1}) \cdot \sigma_k^n && f \text{ preserves the action of face operators,} \\ &= f(x \cdot (\sigma_k^n \circ \delta_k^{n+1})) \cdot \sigma_k^n && \text{right action,} \\ &= f(x) \cdot \sigma_k^n && \delta_k^{n+1} \text{ is a right inverse of } \sigma_k^n. \end{aligned}$$

Therefore, quantifying over x and σ_k^n , we see that f preserves the actions of all elementary face operators and so, as we argued in the first paragraph, it follows that f is a simplicial map. However, we also assumed that f preserved thinness and it is thus a stratified map as required. $\square$

Lemma 119. *Restricting the domain of the superstructure functor of observation 115 to the category of well tempered stratified sets*

$$\text{Strat}_W \xrightarrow{|\cdot|_\bullet} [\mathbb{N}, \underline{\text{SSimp}}]$$

we obtain a functor which is fully faithful.

Proof. To show that $|\cdot|_\bullet$ is faithful, fix two stratified maps $f, g: X \longrightarrow Y$ and observe that, by definition, $|f|_\bullet = |g|_\bullet$ if and only if f and g agree when restricted to each superstructure $|X|_n$ of X . It follows that f and g agree on the union of these superstructures which, by observation 113, is equal to X itself and so they are indeed equal on the whole of X as required.

To show that $|\cdot|_\bullet$ is full, suppose that X and Y are well tempered stratified sets and consider a filtered map $f_\bullet: |X|_\bullet \longrightarrow |Y|_\bullet$ in $[\mathbb{N}, \underline{\text{SSimp}}]$. This is simply a family of semi-simplicial maps $f_i: |X|_i \longrightarrow |Y|_i$ for $i \in \mathbb{N}$ satisfying the naturality condition that if $i < j$ then f_i and f_j act identically on the subset $|X|_i \subseteq |X|_j$.

By observation 113(2) above it is clear that we can define a map $f: X \longrightarrow Y$ by $f(x) = f_{\dim(x)}(x)$ and, indeed, the naturality property of $f_\bullet$ actually implies that if n is an integer for which $x \in |X|_n$ then we have $f(x) = f_n(x)$. In particular, it follows that the restriction of f to $|X|_i$ is equal to f_i for each $i \in \mathbb{N}$, in other words we can infer that $|f|_\bullet = f_\bullet$ as required so long as we can prove that f is a stratified map. To prove this all we need do is demonstrate that this map:

- (a) **is semi-simplicial:** If x is an n -simplex in X and $\alpha: [m] \longrightarrow [n]$ is a face operator then we know that $m < n$ and so, by observation 113(2), we may infer that both x and $x \cdot \alpha$ are elements of $|X|_n$. However, we also know that f and f_n coincide on $|X|_n$ and that f_n is semi-simplicial, so it follows that $f(x \cdot \alpha) = f_n(x \cdot \alpha) = f_n(x) \cdot \alpha = f(x) \cdot \alpha$, that is to say f preserves the action of the face operator α on x . Now, quantifying over x and α , we see that f is semi-simplicial as required.
- (b) **preserves thinness:** If x is a thin n -simplex in X then we know, by observation 113(3), that it is an element of $|X|_{n-1}$. However, we also know that f_{n-1} and f agree on $|X|_{n-1}$ and so it follows that $f(x)$ is an element of the codomain $|Y|_{n-1}$ of f_{n-1} . Applying observation 113(3) again we may infer that $f(x)$ is thin in Y as required.

Finally, we may apply lemma 118 to demonstrate that f is a stratified map as required. $\square$

7. PRE-COMPLICIAL SETS

7.1. Introducing Pre-Complicial Sets.

Definition 120 (primitive t-extensions). For $n \geq 2$ and $k = 1, \dots, n-1$ let

- $\Delta_k^a[n]'$ denote the stratified set obtained from $\Delta_k^a[n]$ by making the $(n-1)$ -simplices δ_{k-1}^n and δ_{k+1}^n thin, and
- $\Delta_k^a[n]''$ denote the stratified set obtained from $\Delta_k^a[n]'$ by making the $(n-1)$ -simplex δ_k^n thin.

We say that the entire subset inclusion

$$\Delta_k^a[n]' \xrightarrow{\subseteq_e} \Delta_k^a[n]''$$

is a *primitive t-extension*.

Definition 121 (pre-complicial sets). A stratified set A is said to be a *pre-complicial set* if it is orthogonal to all primitive t-extensions. Let $\underline{\mathbf{Pcs}}$ denote the full subcategory of $\underline{\mathbf{Strat}}$ whose objects are the pre-complicial sets.

Of course, we know that each of the stratified sets $\Delta_k^a[n]'$ and $\Delta_k^a[n]''$ have only a finite set of non-degenerate simplices and are therefore finitely presentable in $\underline{\mathbf{Strat}}$ (cf. observation 108). It follows that the set of primitive t-extensions is an FP-regulus and so, by observation 59, that $\underline{\mathbf{Pcs}}$ is also an LFP-category which is reflective in $\underline{\mathbf{Strat}}$ (with associated idempotent monad (L_p, η^p)).

We let $\mathbb{T}_{\underline{\mathbf{Pcs}}}$ denote the LE-theory of pre-complicial sets, which we may construct from $\mathbb{T}_{\underline{\mathbf{Strat}}}$, the LE-theory of stratified sets, by adding the set of primitive t-extensions to its FP-regulus.

Observation 122 (describing pre-complicial sets explicitly). By Yoneda's lemma, a stratified map $\Delta_k^a[n] \longrightarrow A$ corresponds to an n -simplex $a \in A$ such that $a \cdot \alpha$ is thin for each k -divided operator α . We call such a simplex k -admissible.

So, a set $A \in \underline{\mathbf{Strat}}$ is pre-complicial iff for each k and each k -admissible simplex $a \in A$ if $a \cdot \delta_{k-1}$ and $a \cdot \delta_{k+1}$ are thin then $a \cdot \delta_k$ is thin as well.

Observation 123 (an explicit description of the reflector L_p). In the sequel it will be useful to have a concrete description of the reflector $L_p: \underline{\mathbf{Strat}} \longrightarrow \underline{\mathbf{Strat}}$ associated with $\underline{\mathbf{Pcs}}$. Starting with a stratified set X we inductively construct a sequence of subsets:

$$\begin{aligned} tX_0 &= tX \\ tX_{n+1} &= \left\{ x \in X \mid \left(\exists w \in X, k \in \mathbb{N} \right) \text{ s.t. } w \text{ is } k\text{-admissible in } (X, tX_n), \right. \\ &\quad \left. w \cdot \delta_{k-1}, w \cdot \delta_{k+1} \in tX_n \text{ and } w \cdot \delta_k = x \right\} \end{aligned}$$

Notice that a degenerate simplex $x \cdot \sigma_k$ is always k -admissible, from which it follows that $tX_0 \subseteq tX_1 \subseteq \dots \subseteq tX_n \subseteq \dots$, and define $L_p(X)$ to be the stratified set which has the same underlying simplicial set as X but has $tL_p(X) \stackrel{\text{def}}{=} \bigcup_{n=1}^{\infty} tX_n$ as its set of thin simplices. It is clear, by construction, that if $f: X \longrightarrow Y$ is a stratified map then $f(tX_n) \subset tY_n$ for each $n \geq 0$ and so f extends to a stratified map $f: L_p(X) \longrightarrow L_p(Y)$, making L_p into a functor. It is also easily shown that $L_p(X)$ is a pre-complicial set and that the family of inclusions $X \xrightarrow{\subseteq_r} L_p(X)$ constitutes the unit η^p for our idempotent monad.

Definition 124 (t-extensions). We call the L_p -invertible stratified maps *t-extensions*. For more on the properties of t-extensions please consult subsection 3.1. We will also use the term *t-almost* as a synonym for L_p -almost (cf. observation 53).

Observation 125 (duals of pre-complicial sets). We can extend the stratified isomorphisms of observation 107 to get two further families of stratified isomorphisms $\Delta_{n-k}^a[n]' \cong (\Delta_k^a[n]')^\circ$ and $\Delta_{n-k}^a[n]'' \cong (\Delta_k^a[n]'')^\circ$. Of course, the diagram

$$\begin{array}{ccc} \Delta_{n-k}^a[n]' & \xrightarrow{\subseteq_e} & \Delta_{n-k}^a[n]'' \\ \cong \downarrow & & \downarrow \cong \\ (\Delta_k^a[n]')^\circ & \xrightarrow{\subseteq_e} & (\Delta_k^a[n]'')^\circ \end{array}$$

commutes for each n and $1 \leq k < n$, in other words the dual of each primitive t-extension is again (isomorphic to) a primitive t-extension.

Now, since the dual operation on Strat is a (strict) involution, and in particular an equivalence of categories, it follows that a stratified set A is orthogonal to each primitive t-extension if and only if its dual A° is orthogonal to each primitive t-extension dual. However, as we have seen, every primitive t-extension dual is in fact (isomorphic to) a primitive t-extension, so it follows that A is pre-complicial if and only if A° is pre-complicial.

Equivalently, we have demonstrated that the class of t-extensions is closed under the action of applying the dual functor $(-)^\circ: \text{Strat} \longrightarrow \text{Strat}$, in other words $f: X \longrightarrow Y$ is a t-extension if and only if $f^\circ: X^\circ \longrightarrow Y^\circ$ is a t-extension.

Observation 126 (a simple explicit characterisation of t-extensions). It is clear from the construction of observation 123 that the underlying simplicial map of any t-extension is an isomorphism in Simp. In general, we will work with t-extensions which are inclusions $X \xrightarrow{\subseteq_e} Y$ associated with an entire subset $X \subseteq_e Y$ (cf. notation 102). Such an inclusion is a t-extension if and only if there exists a sequence of entire subsets $X = X^{(0)} \subseteq_e X^{(1)} \subseteq_e \dots \subseteq_e X^{(n)} \subseteq_e \dots$ of Y such that $Y = \bigcup_{i=0}^\infty X^{(i)}$ and for each $i \geq 0$ and each simplex $x \in tX^{(i+1)} \setminus tX^{(i)}$ there exists a $w \in Y$ and a $k \in \mathbb{N}$ satisfying the conditions

- w is k -admissible in $X^{(i)}$,
- $w \cdot \delta_{k-1}$ and $w \cdot \delta_{k+1}$ are both in thin in $X^{(i)}$, and
- $x = w \cdot \delta_k$

(in which case we say that w witnesses the extension of thinness to x).

7.2. Tensor Products of Pre-Complicial Sets.

Definition 127. Suppose that X and Y are stratified sets then we say that a partition $p, q \in \mathbb{N}$ of $r \in \mathbb{N}$ cleaves an r -simplex (x, y) of $X \times Y$ if either the p -simplex $x \cdot \mathbb{1}_1^{p,q}$ is thin in X or the q -simplex $y \cdot \mathbb{1}_2^{p,q}$ is thin in Y .

Definition 128 (the lax Gray-tensor of stratified sets). We may define a bifunctor

$$\text{Strat} \times \text{Strat} \xrightarrow{\otimes} \text{Strat}$$

called the *lax Gray-tensor* of stratified sets, by letting

$$X \otimes Y \stackrel{\text{def}}{=} (X \times Y, t(X \otimes Y))$$

for each pair of stratified sets X and Y , where:

$$t(X \otimes Y) \stackrel{\text{def}}{=} \left\{ (x, y) \in X \times Y \mid \begin{array}{l} \text{all partitions } p, q \in \mathbb{N} \text{ of } r = \dim(x, y) \\ \text{cleave } (x, y). \end{array} \right\} \quad (16)$$

The Gray-tensor $f \otimes g: X \otimes Y \longrightarrow X' \otimes Y'$ of a pair of stratified maps $f: X \longrightarrow X'$ and $g: Y \longrightarrow Y'$ is simply the stratified map whose underlying simplicial map is the product $f \times g: X \times Y \longrightarrow X' \times Y'$, in other words $(f \otimes g)(x, y) = (f(x), g(y))$.

Before proving that this bifunctor is well defined, we first establish the following simple lemma:

Lemma 129 (simplices in $X \otimes Y$ with degenerate ordinates). *Let X and Y be stratified sets and suppose that (x, y) is an r -simplex in $X \times Y$ and $p, q \in \mathbb{N}$ is an arbitrary partition of r . We have:*

- (a) *If x is degenerate at u then our partition cleaves (x, y) whenever $u < p$.*
- (b) *If y is degenerate at v then our partition cleaves (x, y) whenever $p \leq v$.*
- (c) *If there exists $u \leq v$ such that x is degenerate at u and y is degenerate at v then (x, y) is in $t(X \otimes Y)$.*

Proof.

(a) We have $\text{dom}(\Downarrow_1^{p,q}) = [p]$ so u and $u+1$ are both elements of the domain of this simplicial operator, by our assumption that $0 \leq u < p$, furthermore $\Downarrow_1^{p,q}(u) = u$ and $\Downarrow_1^{p,q}(u+1) = u+1$. But x is degenerate at u and it follows, by observation 29(c), that $x \cdot \Downarrow_1^{p,q}$ is degenerate, and thus thin, in X so our partition cleaves (x, y) as required.

(b) The argument to establish this point is dual to that of (a). We have $\text{dom}(\Downarrow_2^{p,q}) = [q] = [r-p]$ so $v' = v - p$ and $v' + 1$ are both elements of the domain of this simplicial operator, by our assumption that $p \leq v < r$, furthermore $\Downarrow_2^{p,q}(v') = v' + p = v$ and $\Downarrow_2^{p,q}(v') = v + 1$. But y is degenerate at v and it follows, by observation 29(c), that $y \cdot \Downarrow_2^{p,q}$ is degenerate, and thus thin, in X so our partition cleaves (x, y) as required.

(c) Under the condition given we may infer, by parts (a) and (b), that our partition cleaves (x, y) whenever $p > u$ or $p \leq v$. However, since $u \leq v$ also holds we know that one or other of these conditions will always apply. It follows that every partition will cleave (x, y) which is therefore an element of $t(X \otimes Y)$ as stated. $\square$

Proof. (that $\otimes$ is a well defined bifunctor as stated in definition 128) We need to demonstrate that:

1) **For any pair of stratified sets X and Y then $X \otimes Y$ is a well defined stratified set.** Clearly no 0-simplex of $X \times Y$ can be in $t(X \otimes Y)$ (since tX and tY contain no 0-simplices) therefore all we need show is that any degenerate simplex in $X \times Y$ is in $t(X \otimes Y)$. So suppose that (x, y) is a degenerate r -simplex in $X \times Y$ then, by observation 29(a) and the point-wise definition of the simplicial structure of $X \times Y$, we know that this holds if and only if there is some k such that both x and y are degenerate at k . So (x, y) satisfies the conditions of lemma 129(c) with $u = v = k$, by which result we know that it is an element of $t(X \otimes Y)$ as required.

2) **For any pair of stratified maps $f: X \longrightarrow X'$ and $g: Y \longrightarrow Y'$ the simplicial map $f \otimes g$ is a well defined stratified map from $X \otimes Y$ to $X' \otimes Y'$.** All we need to show is that $f \otimes g$ preserves thinness. So, suppose that the r -simplex (x, y) of $X \times Y$ is in $t(X \otimes Y)$ and that $p, q \in \mathbb{N}$ is an arbitrary partition of r , then we know, by defining equation (16), that this partition cleaves (x, y) so either

- $x \cdot \Downarrow_1^{p,q}$ is in tX , in which case, since f is a stratified map, it follows that $f(x) \cdot \Downarrow_1^{p,q} = f(x \cdot \Downarrow_1^{p,q})$ is in tX' , or

- $y \cdot \mathbb{1}_2^{p,q}$ is in tY , in which case, since g is a stratified map, it follows that $g(y) \cdot \mathbb{1}_2^{p,q} = f(y \cdot \mathbb{1}_2^{p,q})$ is in tY' .

In either case, our partition cleaves the simplex $(f \otimes g)(x, y) = (f(x), g(y))$ in $X' \times Y'$ and therefore, since this partition was chosen arbitrarily, it follows that $(f \otimes g)(x, y)$ is in $t(X' \otimes Y')$ as required.

3) $\otimes$ is functorial. This follows trivially from the functoriality of $\times$ as a bifunctor from $\mathbf{Simp} \times \mathbf{Simp}$ to $\mathbf{Simp}$. $\square$

Corollary 130 (of lemma 129). *In a tensor product of standard simplices $\Delta[n] \otimes \Delta[m]$, a simplex (α, β) is thin if and only if it satisfies the condition of lemma 129(c).*

Proof. The “if” direction is simply lemma 129(c). For the “only if” direction, suppose that the r -simplex (α, β) is thin in $\Delta[n] \otimes \Delta[m]$. Pick the partition p, q of r by selecting $p \in [r]$ to be the largest integer such that $\alpha(0) < \alpha(1) < \dots < \alpha(p)$ and letting $q = r - p$. Of course, by our choice of p , the p -simplex $\alpha \circ \mathbb{1}_1^{p,q}$ is non-degenerate and thus non-thin in $\Delta[n]$, since this has the minimal stratification, in which only degenerate simplices are thin. However we know that our partition must cleave (α, β) , by the definition of thinness for the tensor product $\Delta[n] \otimes \Delta[m]$, and thus $\beta \circ \mathbb{1}_2^{p,q}$ must be thin in $\Delta[m]$. It follows that this simplex must be degenerate, again by the minimality of the stratification of $\Delta[m]$, so there must exist some $0 \leq l < q$ such that $\beta \circ \mathbb{1}_2^{p,q}(l) = \beta \circ \mathbb{1}_2^{p,q}(l+1)$ equivalently, using the definition of $\mathbb{1}_2^{p,q}$, we see that there exists some $p \leq k < r$ (which is related to l by $k = l + p$) such that β is degenerate at k . Notice that this also implies that $p < r$ (since $0 < q$ and $p + q = r$) from which we may infer, from the maximality clause in the definition of p , that α is degenerate at p (i.e. $\alpha(p) = \alpha(p+1)$) otherwise p would not be maximal for the stated property). So we have found a pair $p \leq k$ satisfying the condition of lemma 129(c) as required. $\square$

Lemma 131. *Given stratified sets X, Y and Z the structural isomorphisms of the cartesian category $(\mathbf{Simp}, \times, \Delta[0])$ extend to stratified isomorphisms:*

$$X \otimes (Y \otimes Z) \cong (X \otimes Y) \otimes Z \quad (17)$$

$$X \otimes \Delta[0] \cong X \quad (18)$$

$$\Delta[0] \otimes X \cong X \quad (19)$$

Collectively these provide the structural isomorphisms which make the triple $(\mathbf{Strat}, \otimes, \Delta[0])$ into a monoidal category. Furthermore, the canonical isomorphism $(U(X) \times U(Y))^\circ \cong U(X)^\circ \times U(Y)^\circ$ in $\mathbf{Simp}$ extends to a stratified isomorphism:

$$(X \otimes Y)^\circ \cong Y^\circ \otimes X^\circ \quad (20)$$

Proof. The forgetful functor $U: \mathbf{Strat} \rightarrow \mathbf{Simp}$ is faithful and $\otimes$ is defined so that U commutes with it and $\times$, in other words the diagram of functors

$$\begin{array}{ccc} \mathbf{Strat} \times \mathbf{Strat} & \xrightarrow{\otimes} & \mathbf{Strat} \\ U \times U \downarrow & & \downarrow U \\ \mathbf{Simp} \times \mathbf{Simp} & \xrightarrow{\times} & \mathbf{Simp} \end{array}$$

commutes. So, assuming that we can show that the various structural isomorphisms associated with the cartesian structure on $\mathbf{Simp}$ extend as stated to stratified isomorphisms, it follows that they will trivially satisfy the various conditions required of the structural maps

of a monoidal category (Strat, $\otimes$, $\Delta[0]$), simply because they do so in (Simp, $\times$, $\Delta[0]$). We examine each of these isomorphisms in turn:

Associativity. An n -simplex $(x, (y, x))$ of $U(X) \times (U(Y) \times U(Z))$ is in the subset $t(X \otimes (Y \otimes Z))$ if and only if for all partitions p, u of n either

- $x \cdot \mathbb{1}_1^{p,u}$ is in tX , or
- $(y, z) \cdot \mathbb{1}_2^{p,u} = (y \cdot \mathbb{1}_2^{p,u}, z \cdot \mathbb{1}_2^{p,u})$ is in $t(Y \otimes Z)$.

Furthermore, $(y \cdot \mathbb{1}_2^{p,u}, z \cdot \mathbb{1}_2^{p,u})$ is in $t(Y \otimes Z)$ if and only if for all partitions q, r of u either

- $(y \cdot \mathbb{1}_2^{p,u}) \cdot \mathbb{1}_1^{q,r} = y \cdot (\mathbb{1}_2^{p,q+r} \circ \mathbb{1}_1^{q,r})$ is in tY , or
- $(z \cdot \mathbb{1}_2^{p,u}) \cdot \mathbb{1}_2^{q,r} = z \cdot (\mathbb{1}_2^{p,q+r} \circ \mathbb{1}_2^{q,r})$ is in tZ .

By a dual argument, the corresponding n -simplex $((x, y), z)$ in $(U(X) \times U(Y)) \times U(Z)$ is in $t((X \otimes Y) \otimes Z)$ iff for all p, q, r with $p + q + r = n$ we have:

- $x \cdot (\mathbb{1}_1^{p+q,r} \circ \mathbb{1}_1^{p,q})$ is in tX , or
- $y \cdot (\mathbb{1}_1^{p+q,r} \circ \mathbb{1}_2^{p,q})$ is in tY , or
- $z \cdot \mathbb{1}_2^{p+q,r}$ is in tZ .

Applying the partition identities laid out in display (1) of notation 5, we see that corresponding pairs of simplices from these characterisations are actually equal in X , Y and Z respectively and it follows that $(x, (y, z))$ is thin in $X \otimes (Y \otimes Z)$ if and only if $((x, y), z)$ is thin in $(X \otimes Y) \otimes Z$. In other words the associativity isomorphism of $\times$ on Simp preserves and reflects the stratifications associated with $\otimes$ and so extends to the stratified isomorphism of (17) as required.

Right and Left Identity. Consider an r -simplex (x, y) in $X \otimes \Delta[0]$ and a partition p, q of r . There are two cases:

- $q > 0$ in which case $y \cdot \mathbb{1}_2^{p,q}$ would be a simplex of dimension greater than 0 in the terminal simplicial set $\Delta[0]$. However, all simplices of $\Delta[0]$ of dimension greater than 0 are degenerate and therefore thin in $\Delta[0]$. So, it follows that our partition cleaves (x, y) .
- $q = 0$ or equivalently $p = r$, in this case $x \cdot \mathbb{1}_1^{p,q} = x$ and $y \cdot \mathbb{1}_2^{p,q}$ is the unique 0-simplex of $\Delta[0]$ and is therefore not thin. It follows that our partition would cleave (x, y) if and only if x is thin in X .

Quantifying over the partitions of r , it follows that (x, y) is thin in $X \otimes \Delta[0]$ if and only if x is thin in X . In other words, the right identity isomorphism (Simp, $\times$, $\Delta[0]$) extends to the stratified isomorphism of (18) and a dual argument demonstrates that the left identity isomorphism extends to the stratified isomorphism of (19), as required.

Dual Symmetry. Applying observation 6, it is easily seen that a partition p, q cleaves an r -simplex (y, x) in $Y^\circ \otimes X^\circ$ if and only if either

- $y * \mathbb{1}_1^{p,q} \stackrel{\text{def}}{=} y \cdot (\mathbb{1}_1^{p,q})^\circ$ is in tY° but then, by observation 6, $(\mathbb{1}_1^{p,q})^\circ = \mathbb{1}_2^{q,p}$ and so this condition holds iff $y \cdot \mathbb{1}_2^{q,p}$ is in tY , or
- $x * \mathbb{1}_2^{p,q} \stackrel{\text{def}}{=} x \cdot (\mathbb{1}_2^{p,q})^\circ$ is in tX° but then, by observation 6, $(\mathbb{1}_2^{p,q})^\circ = \mathbb{1}_1^{q,p}$ and so this condition holds iff $x \cdot \mathbb{1}_1^{q,p}$ is in tX

which says no more nor less than that q, p cleaves the simplex (x, y) of $X \otimes Y$. Quantifying this result over the partitions of r , we see that (y, x) is in $t(Y^\circ \otimes X^\circ)$ if and only if (x, y) is in $t(X \otimes Y)^\circ \stackrel{\text{def}}{=} t(X \otimes Y)$. In other words, the dual symmetry isomorphism on Simp extends to the stratified isomorphism of (20) as required. $\square$

Observation 132 (tensor products of regular maps). If the stratified maps $f: X \longrightarrow X'$ and $g: Y \longrightarrow Y'$ are both regular then so is their tensor product $f \otimes g: X \otimes Y \longrightarrow X' \otimes Y'$. It follows that a tensor product of regular subsets $X \subseteq_r X'$ and $Y \subseteq_r Y'$ is again a regular subset $X \otimes Y \subseteq_r X' \otimes Y'$.

Proof. Suppose that the r -simplex $(f(x), g(y))$ is thin in $X' \otimes Y'$ and that p, q is a partition of r . By the definition of $t(X' \otimes Y')$, this cleaves $(f(x), g(y))$, so either

- $f(x \cdot \mathbb{1}_1^{p,q}) = f(x) \cdot \mathbb{1}_1^{p,q}$ is in tX' , in which case $x \cdot \mathbb{1}_1^{p,q}$ is in tX since f is regular, or
- $g(y \cdot \mathbb{1}_2^{p,q}) = g(y) \cdot \mathbb{1}_2^{p,q}$ is in tY' , in which case $y \cdot \mathbb{1}_2^{p,q}$ is in tY since g regular.

In either case, we have shown that our partition also cleaves $(x, y) \in X \times Y$ and it follows, by quantifying over such partitions, that (x, y) is in $t(X \otimes Y)$ as required. $\square$

Recall that entire maps (resp. inclusions) are defined to be those stratified maps whose action on underlying simplicial sets is surjective (resp. injective). It therefore goes without saying, since $\otimes$ acts as the cartesian product on underlying sets, that these classes of stratified maps are also closed under (pre-)tensoring.

Lemma 133. *If X and Y are stratified sets and $n, m \geq 0$ are fixed integers then we have entire subset inclusions*

$$\begin{aligned} \text{Th}_{n+m}(X \otimes Y) &\subseteq_e \text{Th}_n(X) \otimes \text{Th}_m(Y) \quad \text{and} \\ \text{Th}_n(X) \otimes \text{Th}_m(Y) &\subseteq_e \text{Th}_{\min(n,m)}(X \otimes Y) \end{aligned} \tag{21}$$

furthermore the argument used to establish the second of these also shows that:

$$\begin{aligned} X \otimes \text{Th}_m(Y) &\subseteq_e \text{Th}_m(X \otimes Y) \quad \text{and} \\ \text{Th}_n(X) \otimes Y &\subseteq_e \text{Th}_n(X \otimes Y) \end{aligned}$$


In particular, applying the inclusions in display (21) it follows that $\text{Th}_0(X \otimes Y) = \text{Th}_0(X) \otimes \text{Th}_0(Y)$.

Proof. Of course $X \otimes Y \subseteq_e \text{Th}_n(X) \otimes \text{Th}_m(Y)$, so in order to prove the first of these inclusions all we need to show is that every simplex of $\text{Th}_n(X) \otimes \text{Th}_m(Y)$ of dimension $> n + m$ is thin. To this end, let $r > n + m$ and consider an r -simplex (x, y) in $\text{Th}_n(X) \otimes \text{Th}_m(Y)$. If p, q is a partition of r then, since $p + q = r > n + m$, we know that either $p > n$ or $q > m$. In the first case $x \cdot \mathbb{1}_1^{p,q}$ is of dimension $p > n$, and thus it is thin in $\text{Th}_n(X)$, and in the latter $y \cdot \mathbb{1}_2^{p,q}$ is of dimension $q > m$, which implies that it is thin in $\text{Th}_m(Y)$; either way we see that our partition cleaves (x, y) . It follows that (x, y) is thin in $\text{Th}_n(X) \otimes \text{Th}_m(Y)$, as required.

Now we also know that $X \otimes Y \subseteq_e \text{Th}_{\min(n,m)}(X \otimes Y)$, so in order to prove the second inclusion all we need to do is show that any r -simplex (x, y) which is thin in $\text{Th}_n(X) \otimes \text{Th}_m(Y)$ and not thin in $X \otimes Y$ is actually also thin in $\text{Th}_{\min(n,m)}(X \otimes Y)$. However, for any such simplex it is clear that there must exist some partition p, q of r for which either $x \cdot \mathbb{1}_1^{p,q}$ is thin in $\text{Th}_n(X)$ but not in X or $y \cdot \mathbb{1}_2^{p,q}$ is thin in $\text{Th}_m(Y)$ but not in Y . In the first of these cases we have $p > n$ and in the second we have $q > m$, but $r = p + q$ so in either case we have $r > \min(n, m)$. It follows, therefore, that any such simplex is thin in $\text{Th}_{\min(n,m)}(X \otimes Y)$ as required. Everything else follows trivially. $\square$

7.3. Pre-Tensors and Preservation of t-Extensions.

Observation 134 (some thin simplices in $X \otimes Y$). While $\otimes$ provides us with a rich and easily defined monoidal structure on Strat it is unfortunately deficient in one important respect -

FIGURE 3. A mediator simplex in $\Delta[n] \otimes \Delta[m]$.

it is not biclosed. However, the remainder of this section will demonstrate that it may be reflected to a monoidal structure on the full subcategory $\underline{\mathbf{Pcs}}$ which **is** biclosed.

To do this we need to analyse the thin simplices of a tensor product $X \otimes Y$ in some greater detail. In fact there are two quite distinct classes of simplices in such tensors which turn out to be of particular importance:

Mediator Simplices. An r -simplex (x, y) of $X \otimes Y$ is called a *mediator simplex* if there exists some $1 \leq k < r$ such that $x = x \cdot (\delta_{k-1} \circ \sigma_{k-1})$ and $y = y \cdot (\delta_k \circ \sigma_k)$. In this case we say that k *witnesses* the fact that (x, y) is a mediator simplex. Notice that if this simplex is non-degenerate then the mediation condition implies immediately that x cannot be degenerate at k and y cannot be degenerate at $k - 1$.

An equivalent way of stating this condition is to say that we may find decompositions $x = x' \cdot \alpha$ and $y = y' \cdot \beta$ (for some $x' \in X$, $y' \in Y$ and $\alpha, \beta \in \Delta$) which have $\alpha(k-1) = \alpha(k)$ and $\beta(k) = \beta(k+1)$.

To visualise this *mediation condition*, consider an r -simplex (α, β) in the tensor $\Delta[n] \otimes \Delta[m]$ recall, from observation 43, that we may consider such a simplex to be a single order preserving map $[r] \longrightarrow [n] \times [m]$. We can draw this as a path in $[n] \times [m]$, as depicted in figure 3, every step of which is a move toward the upper-right corner of that grid. Pictorially the mediation condition at the highlighted path point k states that our arrival there is achieved by a pure upward movement and our departure by a pure rightward one.

Most importantly, all mediator simplices are thin in a tensor $X \otimes Y$; a fact which follows directly from lemma 129(c) since the mediation condition is simply a special case of the condition given there.

Cylinders. An r -simplex (x, y) in $X \times Y$ is said to be a *cylinder* if there exists a partition p, q of r such that $x = x \cdot (\perp_1^{p,q} \circ \pi_1^{p,q})$ and $y = y \cdot (\perp_2^{p,q} \circ \pi_2^{p,q})$. In this case we say that our partition *witnesses* the fact that (x, y) is a cylinder.

The fact that $\perp_i^{p,q}$ is a right inverse of $\pi_i^{p,q}$ (for $i = 1, 2$, see notation 5) implies that our partition witnesses (x, y) as a cylinder if and only if there exists an p -simplex $x' \in X$ and an q -simplex $y' \in Y$ such that $x = x' \cdot \pi_1^{p,q}$ and $y = y' \cdot \pi_2^{p,q}$.

To understand what a cylinder “looks” like we can again consider the cylinders of $\Delta[n] \otimes \Delta[m]$. With reference to figure 4 we see that such a cylinder (α, β) corresponds to a path consisting of a (possibly empty) series of pure rightward moves followed by a (possibly empty)

series of pure upward moves. The projections of this simplex onto the x- and y-axes, as marked in the figure, correspond to the simplices $\alpha \circ \mathbb{I}_1^{p,q} \in \Delta[n]_p$ and $\beta \circ \mathbb{I}_2^{p,q} \in \Delta[m]_q$ respectively.

Now, suppose that the r -simplex (x, y) is witnessed as a cylinder by a partition p, q and suppose that s, t is any other partition. Observe that:

- $\pi_1^{p,q}(p+1) = \pi_1^{p,q}(p)$ and $x = (x \cdot \mathbb{I}_1^{p,q}) \cdot \pi_1^{p,q}$ which is therefore degenerate at p , so applying lemma 129(a) we see that s, t cleaves (x, y) whenever $p < s$.
- $\pi_2^{p,q}(p-1) = \pi_2^{p,q}(p)$ and $y = (y \cdot \mathbb{I}_2^{p,q}) \cdot \pi_2^{p,q}$ which is therefore degenerate at $p-1$, so applying lemma 129(b) with we see that s, t cleaves (x, y) whenever $s \leq p-1$.

It follows that (x, y) is in $t(X \otimes Y)$ if and only if p, q cleaves it.

Consequently, we say that an r -simplex (x, y) of $X \otimes Y$ is a *crushed cylinder* if it is witnessed as a cylinder by a partition which also cleaves it. Our last result simply establishes that a cylinder is thin in $X \otimes Y$ if and only if it is a crushed cylinder in there.

Definition 135 (the pre-tensor of stratified sets). We may define a bifunctor

$$\underline{\text{Strat}} \times \underline{\text{Strat}} \xrightarrow{\quad \boxtimes \quad} \underline{\text{Strat}}$$

called the **pre-tensor** of stratified sets, by letting

$$X \boxtimes Y \stackrel{\text{def}}{=} (X \times Y, t(X \boxtimes Y))$$

where a simplex (x, y) is in $t(X \boxtimes Y)$ if

- it is degenerate, or
- it is a mediator simplex, or
- it is a crushed cylinder.

Given a pair of stratified maps $f: X \longrightarrow X'$ and $g: Y \longrightarrow Y'$, the stratified map $f \boxtimes g$ is defined to have the product map $f \times g: X \times Y \longrightarrow X' \times Y'$ as its underlying simplicial map, in other words $(f \boxtimes g)(x, y) = (f(x), g(y))$.

Furthermore, by observation 134, we know that $X \boxtimes Y$ is an entire stratified subset of $X \otimes Y$ and so we have an entire inclusion map:

$$X \boxtimes Y \hookrightarrow X \otimes Y \quad \subseteq_e \quad (22)$$




FIGURE 4. A cylinder in $\Delta[n] \otimes \Delta[m]$.

The canonical isomorphism $(X \otimes Y)^\circ \cong Y^\circ \otimes X^\circ$ restricts along these inclusions to give an isomorphism $(X \boxtimes Y)^\circ \cong Y^\circ \boxtimes X^\circ$.

Proof. (that $\boxtimes$ is a well defined bifunctor as stated) We showed earlier that $t(X \otimes Y)$ contains all degenerate simplices (definition 128), all mediator simplices and all crushed cylinders (observation 134); it follows that $t(X \boxtimes Y)$ is a subset of $t(X \otimes Y)$. But $t(X \otimes Y)$ contains no 0-simplices, by the proof attached to definition 128, furthermore every degenerate simplex of $X \times Y$ is in $t(X \boxtimes Y)$, by definition, so it is certainly true that $(X \times Y, t(X \boxtimes Y))$ is a well defined stratified subset of $X \otimes Y$.

Inspecting the defining properties of mediator simplices and crushed cylinders, we see that both concepts are defined in terms of operations and properties which are preserved by the stratified maps f and g . It follows that if the r -simplex (x, y) is a mediator simplex witnessed by k (respectively a crushed cylinder witnessed by p, q) then so is $(f \boxtimes g)(x, y) = (f(x), g(y))$. From this fact, it follows that $f \boxtimes g$ preserves thinness as required. $\square$

Observation 136 (why introduce $\boxtimes$). While the tensor $\otimes$ possesses many properties that $\boxtimes$ lacks, for instance it is canonically part of a genuine monoidal structure on Strat, it is also deficient in one major respect - the left and right tensoring functors $X \otimes -$ and $- \otimes Y$ do not preserve colimits. Consequently, $\otimes$ cannot be the tensor of a biclosed monoidal structure on Strat.

This deficiency can be traced back to the very definition of the thin simplices in $X \otimes Y$, which relies upon a property that must hold for **every** partition of the dimension of a given simplex.

On the other hand, $\boxtimes$ does not suffer from this problem and consequently, as we shall show later on, the left and right pre-tensors $X \boxtimes -$ and $- \boxtimes Y$ **do** preserve the colimits of Strat. Furthermore, the more “local” explicit description of the thin simplices of $X \boxtimes Y$ often makes it easy to demonstrate that a given simplicial map $f: X \times Y \longrightarrow A$ extends to a stratified map $f: X \boxtimes Y \longrightarrow A$.

In this context, the next lemma provides the most cogent justification for introducing the pre-tensor $\boxtimes$. In essence it says that “pre-complicial sets do not recognise the difference between $X \otimes Y$ and $X \boxtimes Y$ ”.

Lemma 137 (mediator simplices witness thinness extension). *Suppose that the r -simplex (x, y) is a mediator simplex in $X \otimes Y$, as witnessed by some $1 \leq k < r$, then*

- (1) (x, y) is k -admissible in $X \boxtimes Y$, and
- (2) if $(x, y) \cdot \delta_k^r$ is thin in $X \otimes Y$ then so are the faces $(x, y) \cdot \delta_{k+1}^r$ and $(x, y) \cdot \delta_{k-1}^r$.

Proof. To establish part (1) we need to show that $(x, y) \cdot \mu$ is thin in $X \boxtimes Y$ for each face operator $\mu: [m] \longrightarrow [r]$ with $k-1, k$ and $k+1$ in $\text{im}(\mu)$. So, assume that μ is such a face operator and let $l \in [m]$ be the unique integer with $\mu(l) = k$, for which we know that $\mu(l-1) = k-1$ and $\mu(l+1) = k+1$. Now it is easily demonstrated that $\delta_{k-1}^r \circ \sigma_{k-1}^{r-1} \circ \mu = \mu \circ \delta_{l-1}^m \circ \sigma_{l-1}^{m-1}$ and $\delta_k^r \circ \sigma_k^{r-1} \circ \mu = \mu \circ \delta_l^m \circ \sigma_l^{m-1}$ simply by considering cases and applying these equations along with the mediation condition for (x, y) , we get

$$\begin{aligned}
 x \cdot \mu &= (x \cdot (\delta_{k-1}^r \circ \sigma_{k-1}^{r-1})) \cdot \mu = x \cdot (\delta_{k-1}^r \circ \sigma_{k-1}^{r-1} \circ \mu) \\
 &= x \cdot (\mu \circ \delta_{l-1}^m \circ \sigma_{l-1}^{m-1}) = (x \cdot \mu) \cdot (\delta_{l-1}^m \circ \sigma_{l-1}^{m-1}) \\
 y \cdot \mu &= (y \cdot (\delta_k^r \circ \sigma_k^{r-1})) \cdot \mu = y \cdot (\delta_k^r \circ \sigma_k^{r-1} \circ \mu) \\
 &= y \cdot (\mu \circ \delta_l^m \circ \sigma_l^{m-1}) = (y \cdot \mu) \cdot (\delta_l^m \circ \sigma_l^{m-1})
 \end{aligned}$$

or, in other words, l witnesses the fact that $(x, y) \cdot \mu = (x \cdot \mu, y \cdot \mu)$ is a mediator simplex. It follows, by the fact that all mediator simplices are thin in $X \boxtimes Y$, that $(x, y) \cdot \mu$ is thin there as required.

For part (2), it is only necessary to prove the stated result for $(x, y) \cdot \delta_{k+1}^r$, since the corresponding result for $(x, y) \cdot \delta_{k-1}^r$ may be obtained by applying the former result to the simplex (y, x) in the dual $(X \otimes Y)^\circ \cong Y^\circ \otimes X^\circ$. So we assume that $(x, y) \cdot \delta_k^r = (x \cdot \delta_k^r, y \cdot \delta_k^r)$ is thin in $X \otimes Y$ and demonstrate that the $(r-1)$ -simplex $(x, y) \cdot \delta_{k+1}^r = (x \cdot \delta_{k+1}^r, y \cdot \delta_{k+1}^r)$ is also thin there by considering an arbitrary partition p, q of $r-1$, for which there are two cases:

- $p \geq k$ applying the mediation condition we know that $x = x' \cdot \sigma_{k-1}^{r-1}$ for some $(r-1)$ -simplex x' so $x \cdot \delta_{k+1}^r = x' \cdot (\sigma_{k-1}^{r-1} \circ \delta_{k+1}^r) = x' \cdot (\delta_k^{r-1} \circ \sigma_{k-1}^{r-2}) = (x' \cdot \delta_k^{r-1}) \cdot \sigma_{k-1}^{r-2}$ where the penultimate equality is an application of the appropriate simplicial identity. It follows that $x \cdot \delta_{k+1}^r$ is degenerate at $k-1$ and so our assumption that $p \geq k$ allows us to apply lemma 129(a) to show that p, q cleaves $(x, y) \cdot \delta_{k+1}^r$.
- $p < k$ which implies that $\delta_k^r \circ \mathbb{1}_1^{p,q} = \mathbb{1}^{p,q+1} = \delta_{k+1}^r \circ \mathbb{1}_1^{p,q}$ and it follows that $(x \cdot \delta_{k+1}^r) \cdot \mathbb{1}_1^{p,q} = (x \cdot \delta_k^r) \cdot \mathbb{1}_1^{p,q}$. Furthermore, applying the mediation condition on y we see that $y \cdot \delta_{k+1}^r = (y \cdot (\delta_k^r \circ \sigma_k^{r-1})) \cdot \delta_{k+1}^r = y \cdot (\delta_k^r \circ (\sigma_k^{r-1} \circ \delta_{k+1}^r)) = y \cdot \delta_k^r$ where the last equality follows by applying the appropriate simplicial identity and so, in particular, $(y \cdot \delta_k^r) \cdot \mathbb{1}_2^{p,q} = (y \cdot \delta_{k+1}^r) \cdot \mathbb{1}_2^{p,q}$. From these two equalities, it clearly follows that our partition p, q cleaves $(x, y) \cdot \delta_{k+1}^r$ iff it cleaves $(x, y) \cdot \delta_k^r$, which it does by the assumption that this latter simplex is thin in $X \otimes Y$.

So we have shown that every partition of $r-1$ cleaves $(x, y) \cdot \delta_{k+1}^r$ and thus that it is thin in $X \otimes Y$ as required. $\square$

Lemma 138. *If the r -simplex $(x, y) = (x' \cdot \alpha, y' \cdot \beta)$ is thin in $X \otimes Y$ and not thin in $X \boxtimes Y$ then there exists a $0 < k \leq r$ such that $\alpha(k-1) < \alpha(k)$ and $\beta(k-1) < \beta(k)$.*

Proof. Define two integers p and q as follows

- let p be the largest integer in $[r]$ such that $\alpha(0) < \alpha(1) < \dots < \alpha(p)$ and $\beta(0) = \beta(1) = \dots = \beta(p)$, and
- let q be the largest integer in $[r-p]$ such that $\alpha(p) = \alpha(p+1) = \dots = \alpha(p+q)$ and $\beta(p) < \beta(p+1) < \dots < \beta(p+q)$,

and suppose, for a contradiction, that p, q is a partition of r (that is $p+q=r$). Were this the case then the defining properties of p and q would imply that $\alpha = \alpha \circ \mathbb{1}_1^{p,q} \circ \pi_1^{p,q}$ and $\beta = \beta \circ \mathbb{1}_2^{p,q} \circ \pi_2^{p,q}$. It follows that $x \cdot (\mathbb{1}_1^{p,q} \circ \pi_1^{p,q}) = (x' \cdot \alpha) \cdot (\mathbb{1}_1^{p,q} \circ \pi_1^{p,q}) = x' \cdot (\alpha \circ \mathbb{1}_1^{p,q} \circ \pi_1^{p,q}) = x' \cdot \alpha = x$ and $y \cdot (\mathbb{1}_2^{p,q} \circ \pi_2^{p,q}) = (y' \cdot \beta) \cdot (\mathbb{1}_2^{p,q} \circ \pi_2^{p,q}) = y' \cdot (\beta \circ \mathbb{1}_2^{p,q} \circ \pi_2^{p,q}) = y' \cdot \beta = y$, and so our partition witnesses the fact that (x, y) is a cylinder. Of course we know, from our analysis in observation 134, that a cylinder is thin in $X \otimes Y$ if and only if it is a crushed cylinder. But, all crushed cylinders are thin in $X \boxtimes Y$ (by definition) which contradicts our original assumption that (x, y) is **not** thin in $X \boxtimes Y$, so it follows that $p+q < r$.

Now, let $k = p+q+1 \in [r]$ for which we may show that:

- $\alpha(k-1) < \alpha(k)$. *Proof.* Suppose, for a contradiction, that $\alpha(k-1) = \alpha(k)$; then either
 - $\beta(k-1) = \beta(k)$, in which case (x, y) would be degenerate, which would contradict the assumption that it is not thin in $X \boxtimes Y$, or

- $\beta(k-1) < \beta(k)$, which would contradict the maximality clause in the definition of q given above.

In either case we get a contradiction and it follows that $\alpha(k-1) < \alpha(k)$ as required.

- $\beta(k-1) < \beta(k)$. *Proof.* Suppose, for a contradiction, that $\beta(k-1) = \beta(k)$, then we already know that $\alpha(k-1) < \alpha(k)$ and so either:

- $q = 0$, in which case $k-1 = p$ and we get a contradiction of the maximality clause in the definition of p given above.
- $q > 0$, from which we may infer that $\alpha(k-2) = \alpha(k-1)$ which, when combined with the assumption that $\beta(k-1) = \beta(k)$, would imply that $k-1$ witnesses that (x, y) is a mediator simplex, again contradicting the assumption that it is not thin in $X \boxtimes Y$.

In either case we get a contradiction and it follows that $\beta(k-1) < \beta(k)$ as required. $\square$

Lemma 139. *For each pair of stratified sets X and Y the entire inclusion map of display (22) is a t -extension.*

Proof. Our intention is to prove this lemma by applying observation 126. To this end, we define a function $\Phi: \Delta[n] \times \Delta[m] \longrightarrow \mathbb{N}$ by

$$\Phi(\alpha, \beta) \stackrel{\text{def}}{=} \sum_{i=0}^{\dim(\alpha, \beta)} \alpha(i) + (m - \beta(i))$$

and an increasing sequence of subsets $tA^{(0)} \subseteq tA^{(1)} \subseteq \dots \subseteq tA^{(n)} \subseteq \dots$ of $t(X \otimes Y)$ by

$$tA^{(n)} \stackrel{\text{def}}{=} t(X \boxtimes Y) \cup \left\{ (x, y) \in t(X \otimes Y) \mid \begin{array}{l} \exists x' \in X, y' \in Y, \alpha, \beta \in \Delta \text{ such that} \\ x = x' \cdot \alpha \wedge y = y' \cdot \beta \wedge \Phi(\alpha, \beta) \leq n \end{array} \right\}$$

and let $A^{(n)}$ be the stratified set $(X \times Y, tA^{(n)})$. The following observations are of note:

$tA^{(0)} = t(X \boxtimes Y)$. *Proof.* Suppose that the r -simplex $(x, y) \in t(X \otimes Y)$ possesses decompositions $x = x' \cdot \alpha$ and $y = y' \cdot \beta$ such that $\Phi(\alpha, \beta) \leq 0$. Consulting the definition of Φ , we see that for this latter condition to hold each $\alpha(i) = 0$ and that each $\beta(i) = m$ where $\text{cod}(\beta) = [m]$. Furthermore r must be greater than 0, since $t(X \otimes Y)$ does not contain any 0-simplices, therefore $0, 1 \in [r]$, $\alpha(0) = \alpha(1)$ and $\beta(0) = \beta(1)$ from which we may infer that $(x, y) = (x' \cdot \alpha, y' \cdot \beta)$ is degenerate and so it is an element of $t(X \boxtimes Y)$. It follows, by the definition of $tA^{(0)}$, that $tA^{(0)} \subseteq t(X \boxtimes Y)$ as required.

$\bigcup_{n=0}^{\infty} tA^{(n)} = t(X \otimes Y)$. *Proof.* Given an r -simplex $(x, y) \in t(X \otimes Y)$ we know that $x = x \cdot \text{id}_{[r]}$, $y = y \cdot \text{id}_{[r]}$ and $\Phi(\text{id}_{[r]}, \text{id}_{[r]}) = r^2$; it follows, by the defining property of $tA^{(r^2)}$, that $(x, y) \in tA^{(r^2)}$. In other words, every element of $t(X \otimes Y)$ is a member of those $tA^{(n)}$ for which n is “sufficiently large” and therefore $t(X \otimes Y) \subseteq \bigcup_{n=0}^{\infty} tA^{(n)}$ as required.

Now we are in position to verify the postulates of observation 126, so suppose that (x, y) is an r -simplex in $tA^{(n+1)} \setminus tA^{(n)}$. We know, by the definition of $tA^{(n+1)}$, that we may find decompositions $x = x' \cdot \alpha$ and $y = y' \cdot \beta$ such that $\Phi(\alpha, \beta) \leq n+1$, furthermore $tA^{(n+1)} \subseteq t(X \otimes Y)$ and $t(X \boxtimes Y) \subseteq tA^{(n)}$ so (x, y) is thin in $X \otimes Y$ and it is **not** thin in $X \boxtimes Y$, and it follows that we may apply lemma 138 to find a $0 < k \leq r$ with $\alpha(k-1) < \alpha(k)$ and $\beta(k-1) < \beta(k)$. So, consider the $(r+1)$ -simplex $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r)$ and observe that:

- k witnesses that $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r)$ is a mediator simplex, in particular it is k -admissible in $X \boxtimes Y$, by lemma 137, and $t(X \boxtimes Y) \subseteq tA^{(n)}$ so it is also k -admissible in $A^{(n)}$.

- $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r) \cdot \delta_k^{r+1} = (x \cdot (\sigma_{k-1}^r \circ \delta_k^{r+1}), y \cdot (\sigma_k^r \circ \delta_k^{r+1})) = (x, y)$. However, we selected this so that it was thin in $X \otimes Y$ and we may apply lemma 137(2) to show that the faces $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r) \cdot \delta_{k+1}^{r+1}$ and $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r) \cdot \delta_{k-1}^{r+1}$ are also thin in there.
- $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r) \cdot \delta_{k-1}^{r+1} = (x \cdot (\sigma_{k-1}^r \circ \delta_{k-1}^{r+1}), y \cdot (\sigma_k^r \circ \delta_{k-1}^{r+1})) = (x, y \cdot (\sigma_k^r \circ \delta_{k-1}^{r+1}))$ and applying our decompositions of x and y this is equal to $(x' \cdot \alpha, y' \cdot (\beta \circ \sigma_k^r \circ \delta_{k-1}^{r+1}))$. However, notice that $\beta \circ \sigma_k^r \circ \delta_{k-1}^{r+1}$ and β only differ at $k-1$ where $\beta \circ \sigma_k^r \circ \delta_{k-1}^{r+1}(k-1) = \beta(k)$, so it follows that

$$\begin{aligned} \Phi(\alpha, \beta \circ \sigma_k^r \circ \delta_{k-1}^{r+1}) &= \Phi(\alpha, \beta) + \beta(k-1) - \beta(k) && \text{by the definition of } \Phi, \\ &< \Phi(\alpha, \beta) - 1 && \text{since } \beta(k-1) < \beta(k), \\ &\leq n \end{aligned}$$

and so $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r) \cdot \delta_{k-1}^{r+1}$, which we've already shown to be thin in $X \otimes Y$, is an element of $tA^{(n)}$ (by definition).

- dually $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r) \cdot \delta_{k+1}^{r+1} = (x' \cdot (\alpha \circ \sigma_{k-1}^r \circ \delta_{k+1}^{r+1}), y' \cdot \beta)$, but $\alpha \circ \sigma_{k-1}^r \circ \delta_{k+1}^{r+1}$ and α only differ at k where $\alpha \circ \sigma_{k-1}^r \circ \delta_{k+1}^{r+1}(k) = \alpha(k-1)$. It follows that

$$\begin{aligned} \Phi(\alpha \circ \sigma_{k-1}^r \circ \delta_{k+1}^{r+1}, \beta) &= \Phi(\alpha, \beta) - \alpha(k) + \alpha(k-1) && \text{by the definition of } \Phi, \\ &\leq \Phi(\alpha, \beta) - 1 && \text{since } \alpha(k-1) < \alpha(k), \\ &\leq n \end{aligned}$$

and so $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r) \cdot \delta_{k+1}^{r+1}$, which we've already shown to be thin in $X \otimes Y$, is also an element of $tA^{(n)}$ (by definition).

In summary, if (x, y) is an r -simplex of $tA^{(n+1)} \setminus tA^{(n)}$ then the $r+1$ -simplex $(x \cdot \sigma_{k-1}^r, y \cdot \sigma_k^r)$ which we constructed witnesses the extension of thinness to (x, y) , as required by observation 126.

This completes the proof that the sequence $\{A^{(n)}\}_{i=0}^\infty$ satisfies all of the conditions of observation 126, from which it follows that the stratified map $X \boxtimes Y \xrightarrow{\subseteq_e} X \otimes Y$ is a t -extension as required. $\square$

Lemma 140. *If $X' \subseteq_e X$ and $Y' \subseteq_e Y$ are entire stratified subsets then we have a commutative square*

$$\begin{array}{ccc} (X \boxtimes Y') \cup (X' \boxtimes Y) & \xrightarrow{\subseteq_e} & X \boxtimes Y \\ \downarrow \subseteq_e & & \downarrow \subseteq_e \\ (X \otimes Y') \cup (X' \otimes Y) & \xrightarrow{\subseteq_e} & X \otimes Y \end{array} \quad (23)$$

of entire inclusions in which the upper horizontal is a stratified isomorphism and the lower horizontal is a t -extension.

Proof. By lemma 139 the vertical inclusions in this square are t -extensions so, by observation 52(5), it follows that its lower horizontal is a t -extension if and only if its upper horizontal is a t -extension. So in order to establish both of the stated results it is enough to demonstrate that this latter map is an isomorphism, however since it is an entire inclusion all that remains is to demonstrate that it is also regular.

To show that this is the case, it is clear that all we need do is demonstrate that any crushed cylinder (x, y) in $X \boxtimes Y$ is a crushed cylinder in one of $X \boxtimes Y'$ or $X' \boxtimes Y$. So suppose that

(x, y) is an r -simplex and that it is witnessed as a crushed cylinder by the partition p, q of r then, by definition, we have two cases

- $x \cdot \mathbb{I}_1^{p,q}$ is thin in X , so our partition also witnesses (x, y) as a crushed cylinder in $X \boxtimes Y'$, or
- $y \cdot \mathbb{I}_2^{p,q}$ is thin in Y , so our partition also witnesses (x, y) as a crushed cylinder in $X' \boxtimes Y$

as required. $\square$

Observation 141. Of course, lemma 139 implies that if A is a pre-complicial set then we may show that a simplicial map $f: X \times Y \longrightarrow A$ extends to a stratified map $f: X \otimes Y \longrightarrow A$ simply by verifying that f maps all **mediator simplices** and **crushed cylinders** in $X \otimes Y$ to thin simplices in A .

7.4. Some Other Preservation Properties.

Lemma 142. *For each stratified set Y the functor*

$$\underline{\text{Strat}} \xrightarrow{- \boxtimes Y} \underline{\text{Strat}}$$

preserves all small colimits.

Proof. Given a diagram $D: \mathbb{C} \longrightarrow \underline{\text{Strat}}$ our aim is to show that the canonical stratified map

$$\text{colim}(D(-) \boxtimes Y) \xrightarrow{c_{D,Y}} \text{colim}(D) \boxtimes Y \quad (24)$$

is a stratified isomorphism. In fact, we know that the functor $U: \underline{\text{Strat}} \longrightarrow \underline{\text{Simp}}$ preserves colimits and that, for a fixed stratified set Y , we have a commutative square of functors:

$$\begin{array}{ccc} \underline{\text{Strat}} & \xrightarrow{- \boxtimes Y} & \underline{\text{Strat}} \\ U \downarrow & \cong & \downarrow U \\ \underline{\text{Simp}} & \xrightarrow{- \times Y} & \underline{\text{Simp}} \end{array}$$

So the underlying simplicial map of $c_{D,Y}$ is simply the corresponding comparison map

$$\text{colim}(U \circ D(-) \times Y) \longrightarrow \text{colim}(U \circ D) \times Y$$

in $\underline{\text{Simp}}$, which is an isomorphism since $\underline{\text{Simp}}$ is a cartesian closed category (from which it follows that $- \times Y$ preserve colimits). We can infer, using observation 100, that all we need to do is demonstrate that $c_{D,Y}$ is regular.

Under the concrete presentation of the colimits in $\underline{\text{Strat}}$ given in observations 31 and 109, it is clear that this map bears an explicit description under which it carries the equivalence class $[i, (z, y)]$ in $\text{colim}(D(-) \boxtimes Y)$ to the pair $([i, z], y)$ in $\text{colim}(D) \boxtimes Y$. So suppose that the non-degenerate r -simplex $([i, z], y)$ is thin in $Y \boxtimes \text{colim}(D)$, our aim will be to find another $j \in \mathbb{C}$ and $w \in D(j)$ such that (w, y) is thin in $D(j) \boxtimes Y$ and $[j, w] = [i, z]$. By definition 135 there are two possibilities:

- **$([i, z], y)$ is a mediator simplex as witnessed by some k** , in other words $[i, z] = [i, z] \cdot (\delta_{k-1}^r \circ \sigma_{k-1}^{r-1})$ and $y = y \cdot (\delta_k^r \circ \sigma_k^{r-1})$. Notice that $[i, z] \cdot (\delta_{k-1}^r \circ \sigma_{k-1}^{r-1}) = [i, z \cdot (\delta_{k-1}^r \circ \sigma_{k-1}^{r-1})]$.

$(\delta_{k-1}^r \circ \sigma_{k-1}^{r-1})]$, so let $w = z \cdot (\delta_{k-1}^r \circ \sigma_{k-1}^{r-1})$, $j = i$ and observe that the integer k also witnesses the r -simplex (w, y) in $D(j) \boxtimes Y$ as a mediator simplex.

- **$([i, z], y)$ is a crushed cylinder as witnessed by some partition p, q of r** , in other words $[i, z] = [i, z] \cdot (\perp_1^{p,q} \circ \pi_1^{p,q})$, $y = y \cdot (\perp_2^{p,q} \circ \pi_2^{p,q})$ and either of $[i, z] \cdot \perp_1^{p,q}$ or $y \cdot \perp_2^{p,q}$ is thin. If $[i, z] \cdot \perp_1^{p,q}$ is thin then we can find an $j \in \mathbb{C}$ and a $v \in D(j)$ such that v is thin and $[j, v] = [i, z] \cdot \perp_1^{p,q}$, otherwise, simply let $j = i$ and $v = z \cdot \perp_1^{p,q}$. Setting $w = v \cdot \pi_1^{p,q}$, it is easy to show that the r -simplex (w, y) is a crushed cylinder in $D(j) \boxtimes Y$ and that $[i, z] = [j, w]$.

In either case (w, y) is thin in $D(j) \boxtimes Y$, which implies that $[j, (w, y)]$ is thin in $\text{colim}(D(-) \boxtimes Y)$. Furthermore $c_{D,Y}[j, (w, y)] = ([j, w], y) = ([i, z], y)$, so the unique simplex mapping to $([i, z], y)$ under $c_{D,Y}$ is thin in $\text{colim}(D(-) \boxtimes Y)$; in other words $c_{D,Y}$ is regular as required. $\square$

Lemma 143 (tensor products of primitive t-extensions). *If Y is a stratified set then in the commutative square*

$$\begin{array}{ccc} \Delta_k^a[n]' \boxtimes Y & \xrightarrow{\subseteq_e} & \Delta_k^a[n]'' \boxtimes Y \\ \subseteq_e \downarrow & & \downarrow \subseteq_e \\ \Delta_k^a[n]' \otimes Y & \xrightarrow{\subseteq_e} & \Delta_k^a[n]'' \otimes Y \end{array} \quad (25)$$

the horizontal inclusions, which may be constructed by tensoring and pre-tensoring the primitive t-extension $\subseteq_e: \Delta_k^a[n]' \hookrightarrow \Delta_k^a[n]''$ by Y , are both t-extensions.

Proof. Firstly observe that the vertical inclusions in the square are both t-extensions, by lemma 139, and therefore applying observation 52(5) we see that its lower horizontal is a t-extension if and only if its upper horizontal is a t-extension. We may factor the lower of these two into a pair of entire inclusions

$$\Delta_k^a[n]' \otimes Y \xrightarrow{\subseteq_e} (\Delta_k^a[n]' \otimes Y) \cup (\Delta_k^a[n]'' \boxtimes Y) \xrightarrow{\subseteq_e} \Delta_k^a[n]'' \otimes Y \quad (26)$$

and show that their composite is a t-extension by showing that each of its factors is and applying the composition result of observation 52(5).

To show that the right hand inclusion here is a t-extension, we factor the t-extension of lemma 139 as

$$\Delta_k^a[n]'' \boxtimes Y \xrightarrow{\subseteq_e} (\Delta_k^a[n]' \otimes Y) \cup (\Delta_k^a[n]'' \boxtimes Y) \xrightarrow{\subseteq_e} \Delta_k^a[n]'' \otimes Y$$

and appeal to the right cancellation result given in observation 52(5), which we may apply in this case since all entire subset inclusions are epimorphisms in Strat.

Finally we show that the left hand inclusion in display (26) is also a t-extension by demonstrating that we may apply observation 126 to the “single step” sequence $\Delta_k^a[n]' \otimes Y \subseteq_e (\Delta_k^a[n]' \otimes Y) \cup (\Delta_k^a[n]'' \boxtimes Y)$. By the definition of the pre-tensor product $\boxtimes$ we know that if an r -simplex (α, y) is thin in $\Delta_k^a[n]'' \boxtimes Y$ and not thin in $\Delta_k^a[n]' \otimes Y$ then it must be a cylinder witnessed by a partition s, t for which $y' \stackrel{\text{def}}{=} y \cdot \perp_2^{s,t}$ is not thin in Y and $\alpha \circ \perp_1^{s,t}$ is thin in $\Delta_k^a[n]''$ but not in $\Delta_k^a[n]'$. This latter condition implies that $\alpha \circ \perp_1^{s,t} = \delta_k^n$ and so we have $s = n - 1$ and $t = r - n + 1$ (in particular $r \geq n - 1$). Consequently consider the $(r + 1)$ -simplex $(\pi_1^{n,t}, y' \cdot \pi_2^{n,t})$ and observe that this witnesses the extension of thinness from $\Delta_k^a[n]' \otimes Y$ to the simplex (α, y) as verified in the following points:

- (i) If $i \in [n]$ then a simple calculation with simplicial operators demonstrates that:

$$\begin{aligned} (\pi_1^{n,t}, y' \cdot \pi_2^{n,t}) \cdot \delta_i^{r+1} &= (\pi_1^{n,t} \circ \delta_i^{r+1}, y' \cdot (\pi_2^{n,t} \circ \delta_i^{r+1})) \\ &= \begin{cases} (\delta_i^n \circ \pi_1^{n-1,t}, y' \cdot \pi_2^{n-1,t}) & \text{when } i < n, \\ (\pi_1^{n,t-1}, y' \cdot \pi_2^{n-1,t}) & \text{when } i = n. \end{cases} \end{aligned} \quad (27)$$

In particular, the assumption that s, t witnesses our simplex (α, y) as a cylinder implies that $\alpha = \delta_k^n \circ \pi_1^{s,t}$ and $y = y' \cdot \pi_2^{s,t}$, so it follows, from the calculation above, that $(\pi_1^{n,t}, y' \cdot \pi_2^{n,t}) \cdot \delta_k^{r+1} = (\alpha, y)$ (since $k < n$).

- (ii) We know that $k-1 < n$ so applying equation (27) we see that the face $(\pi_1^{n,t}, y' \cdot \pi_2^{n,t}) \cdot \delta_{k-1}^{r+1}$ is a cylinder which is thin in $\Delta_k^a[n]' \otimes Y$ since δ_{k-1}^n is thin in $\Delta_k^a[n]'$.
- (iii) If $k+1 < n$ then we can apply the same argument to show that the face $(\pi_1^{n,t}, y' \cdot \pi_2^{n,t}) \cdot \delta_{k+1}^{r+1}$ is thin in $\Delta_k^a[n]' \otimes Y$, leaving us the special case $k = n-1$ in which equation (27) identifies this face as the simplex $(\pi_1^{n,t-1}, y' \cdot \pi_2^{n-1,t})$. To show that this face is thin in $\Delta_k^a[n]' \otimes Y$ simply observe that
- $\pi_1^{n,t-1} \circ \mathbb{I}^{n-1,t} = \delta_n^n$ and $\pi_1^{n,t-1} \circ \mathbb{I}^{n,t-1} = \text{id}_{[n]}$ are both thin in $\Delta_k^a[n]'$ so the partitions $n-1, t$ and $n, t-1$ both cleave $(\pi_1^{n,t-1}, y' \cdot \pi_2^{n-1,t})$, and
 - $\pi_1^{n,t-1}$ is degenerate at n and $y' \cdot \pi_2^{n-1,t}$ is degenerate at $n-2$ so we may apply observation 129 to show that all other partitions also cleave that simplex, thereby completing the demonstration that it is thin as stated.
- (iv) If $\mu: [m] \longrightarrow [r+1]$ is a k -divided face operator and $l \in [m]$ is the unique integer for which $\mu(l) = k$ (and thus $\mu(l-1) = k-1$ and $\mu(l+1) = k+1$) and consider the face $(\pi_1^{n,t}, y' \cdot \pi_2^{n,t}) \cdot \mu = (\pi_1^{n,t} \circ \mu, y' \cdot (\pi_2^{n,t} \circ \mu))$. We show that this is thin in $\Delta_k^a[n]' \otimes Y$ by selecting an arbitrary partition p, q of m and considering cases:
- $p \leq l$ in which case we have $\mu \circ \mathbb{I}_2^{p,q}(l-p) = \mu(l) = k$ and $\mu \circ \mathbb{I}_2^{p,q}(l-p+1) = \mu(l+1) = k+1$, however $k < k+1 \leq n$ so $\pi_2^{n,t}(k) = \pi_2^{n,t}(k+1) = 0$ and it follows that $(y' \cdot (\pi_2^{n,t} \circ \mu)) \cdot \mathbb{I}_2^{p,q} = y' \cdot (\pi_2^{n,t} \circ \mu \circ \mathbb{I}_2^{p,q})$ is degenerate at $l-p$ and is thus thin in Y , or
 - $p \geq l+1$ which implies that $l-1, l$ and $l+1$ are all elements in the domain of $\pi_1^{n,t} \circ \mu \circ \mathbb{I}_1^{p,q}$ which maps them to $k-1, k$ and $k+1$ respectively, so it follows that this operator is k -divided and thus thin in $\Delta_k^a[n]$.

In either case we've shown that p, q cleaves the simplex $(\pi_1^{n,t}, y' \cdot \pi_2^{n,t}) \cdot \mu$ so quantifying over all such partitions we see that this face is thin in $\Delta_k^a[n] \otimes Y \subseteq_e \Delta_k^a[n]' \otimes Y$. $\square$

7.5. A Monoidal Biclosed Structure on Pre-Complicial sets.

Corollary 144 (of lemma 142). *The functor $- \boxtimes Y: \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ admits a right adjoint, which we'll denote by $* \Leftarrow Y: \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$. Dually, for each stratified set X the functor $X \boxtimes -: \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ has a right adjoint $X \Rightarrow *: \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ given by $(X \Rightarrow *) \stackrel{\text{def}}{=} ((*)^\circ \Leftarrow X^\circ)^\circ$.*

Proof. We know, from observation 108, that the small full subcategory $t\Delta$ is dense in $\underline{\text{Strat}}$. Applying theorem 5.33 of Kelly [19], it follows that any (small) colimit preserving functor with codomain $\underline{\text{Strat}}$ has a right adjoint. In particular, in view of lemma 142 this result applies to $- \boxtimes Y$ giving us stated right adjoint $* \Leftarrow Y$.

The dual result follows directly from basic properties of the strictly involutive dual $(-)^\circ$ and the canonical isomorphisms $(X \boxtimes Y)^\circ \cong Y^\circ \boxtimes X^\circ$. $\square$

Observation 145 (presenting $A \Leftarrow Y$ and $X \Rightarrow A$ explicitly). The result of the last corollary fully characterises $A \Leftarrow Y$, however it will sometimes be useful to assume an explicit representation of this stratified set. Using Yoneda's lemma, we know that an n -simplex in $A \Leftarrow Y$ corresponds to a stratified map $\Delta[n] \longrightarrow A \Leftarrow Y$ which in turn corresponds to a stratified map

$$\Delta[n] \boxtimes Y \xrightarrow{f} A \quad (28)$$

via the adjunction between the functors $- \boxtimes Y$ and $* \Leftarrow Y$ of the last corollary. It follows that we may identify n -simplices in $A \Leftarrow Y$ and stratified maps f of the form shown in display (28) so, using the usual naturality properties of adjunctions, we may derive the following explicit presentation of $A \Leftarrow Y$:

- **n -simplices** are stratified maps of the form given in figure (28),
- **simplicial action** of an operator $\alpha: [m] \longrightarrow [n]$ on an n -simplex f given by $f \cdot \alpha \stackrel{\text{def}}{=} f \circ (\Delta(\alpha) \boxtimes Y)$,
- **thin simplices** those n -simplices $f: \Delta[n] \boxtimes Y \longrightarrow A$ which extend to a stratified map $f: \Delta[n]_t \boxtimes Y \longrightarrow A$.

Dually, we may represent the n -simplices of $X \Rightarrow A$ as stratified maps $g: X \boxtimes \Delta[n]_? \longrightarrow A$ under the simplicial action given by $g \cdot \alpha \stackrel{\text{def}}{=} g \circ (X \boxtimes \Delta(\alpha))$.

Observation 146. Of course, if A is a pre-complicial set then we know, by lemma 139, that stratified maps of the form given in figure (28) extend to maps of the form:

$$\Delta[n] \otimes Y \xrightarrow{f} A \quad (29)$$

It follows that, in this case, it is equally valid to identify the elements of $A \Leftarrow Y$ with stratified maps of the form given in figure (29). The remainder of the explicit description of $A \Leftarrow Y$ given in the last observation carries over to elements represented in this form simply by substituting $\otimes$ for $\boxtimes$ wherever it occurs.

Lemma 147. Suppose that Y and A are stratified sets and that $Y' \subseteq_e Y$ is an entire subset then we may canonically identify $A \Leftarrow Y$ with a regular stratified subset of $A \Leftarrow Y'$. Furthermore, a stratified map $g: Y' \longrightarrow X$ extends to a stratified map with domain Y iff for each A the corresponding map $A \Leftarrow g: A \Leftarrow X \longrightarrow A \Leftarrow Y'$ lifts to a stratified map with codomain $A \Leftarrow Y$.

Proof. Applying the contravariant functor $A \Leftarrow *$ to the entire inclusion $Y' \xrightarrow{\subseteq_e} Y$ gives rise to a stratified map

$$A \Leftarrow Y \xrightarrow{A \Leftarrow (\subseteq_e)} A \Leftarrow Y' \quad (30)$$

which, in terms of the explicit presentation given above, acts on an n -simplex f of $A \Leftarrow Y'$ to restrict its domain from $\Delta[n] \boxtimes Y$ to its entire stratified subset $\Delta[n] \boxtimes Y'$. This action is clearly injective, since these two stratified sets share the same underlying simplicial set.

This result certainly allows us to identify $A \Leftarrow Y$ with a stratified subset of $A \Leftarrow Y'$, but we also need to show that the inclusion in (30) is regular. So suppose that $f: \Delta[n] \boxtimes Y \longrightarrow A$ is an n -simplex in $A \Leftarrow Y$ which maps to a thin simplex in $A \Leftarrow Y'$. We know that this simply means that the restriction of f to $\Delta[n] \boxtimes Y'$ extends to a stratified map with domain $\Delta[n]_t \boxtimes Y'$. Clearly it follows that f extends to $(\Delta[n] \boxtimes Y) \cup (\Delta[n]_t \boxtimes Y')$ and, by lemma 140, this latter stratified set is equal to $\Delta[n]_t \boxtimes Y$ thus f is thin in $A \Leftarrow Y$ as required.

The remainder of the observation is a matter of simple abstract nonsense and is left as an exercise. $\square$

Theorem 148. *For each stratified set Y the right tensor functor $- \otimes Y : \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ has partial right adjoint $* \Leftarrow Y : \underline{\text{Pcs}} \longrightarrow \underline{\text{Pcs}}$ (cf. observation 54). Dually, the left tensor functor $X \otimes - : \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ associated with a stratified set X has partial right adjoint $X \Rightarrow * : \underline{\text{Pcs}} \longrightarrow \underline{\text{Pcs}}$. It follows, by observation 54(1), that if $f : Z \longrightarrow Z'$ is a t -extension then so are the maps:*

$$\begin{array}{ccc} Z \otimes Y & \xrightarrow{f \otimes Y} & Z' \otimes Y \\ X \otimes Z & \xrightarrow{X \otimes f} & X \otimes Z' \end{array}$$

Furthermore, by Day's reflection theorem (theorem 55), we may reflect the monoidal structure on $\underline{\text{Strat}}$ to a monoidal biclosed structure on $\underline{\text{Pcs}}$ with tensor product $A \otimes_p B \stackrel{\text{def}}{=} L_p(A \otimes B)$ (where L_p is reflector associated with $\underline{\text{Pcs}}$) and left and right closures $A \Rightarrow *$ and $* \Leftarrow B$ respectively.

Proof. Fix a stratified set Y . We know (by definition) that the set of primitive t -extensions is adequate to detect pre-complicial sets and that the functor $- \otimes Y : \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ maps primitive t -extensions to t -extensions (by lemma 143). We may infer that it is enough to show that $* \Leftarrow Y : \underline{\text{Pcs}} \longrightarrow \underline{\text{Strat}}$ is a partial right adjoint to $- \otimes Y$, since if that is the case then we may apply observation 54(2) to demonstrate that $* \Leftarrow A$ maps pre-complicial sets to pre-complicial sets. However, this latter result is a direct consequence of corollary 144 via lemma 139, as demonstrated by the following sequence of natural bijections

$$\begin{array}{ccc} X \longrightarrow A \Leftarrow Y & & \text{corollary 144,} \\ \hline X \boxtimes Y \longrightarrow A & \text{the } t\text{-extension of lemma 139 is } \perp & \\ \hline X \otimes Y \longrightarrow A & \text{to the pre-complicial set } A, & \end{array} \quad (31)$$

The remainder of this theorem, that is the corresponding result for $\Rightarrow$, is an immediate dual of the above result. $\square$

Notation 149. When making calculations with the tensor $\otimes$ and the partial closures $* \Leftarrow Y$ and $X \Rightarrow *$ we will tend to simplify matters by following traditional conventions with regard to associativities and the identity object. In particular:

- We often write unbracketed n -fold tensors $X_1 \otimes X_2 \otimes \dots \otimes X_n$ and assume that such expressions are interpreted via the consistent application of a rule, such as right associativity, under which they might be bracketed into binary tensors.
- Wherever possible, we will reduce tensor and closure expressions which involve the identity object $\Delta[0]$ to eliminate them wherever possible. In other words, we will tend to identify expressions such as $X \otimes \Delta[0]$ with X , $\Delta[0] \otimes Y$ with Y , $A \Leftarrow \Delta[0]$ with A and $\Delta[0] \Rightarrow B$ with B .

These rules tacitly assume that whenever we manipulate expressions involving composites of arrows between objects which have been specified in this way, we will be prepared to silently introduce canonical associativity and identity isomorphisms wherever necessary to make sense of those composites. That this is possible to do in general is a consequence of the coherence theorem for monoidal categories and we refer the reader to Mac Lane's book [22] for more on that topic.

7.6. Superstructures of Pre-Complcial Sets.

Lemma 150. *If A is a pre-complicial set then so is its n -dimensional superstructure $|A|_n$. Equivalently, if the stratified map $f: X \longrightarrow Y$ is a t -extension then so is the associated stratified map $\text{Th}_n(f): \text{Th}_n(X) \longrightarrow \text{Th}_n(Y)$.*

Proof. Since $\text{Th}_n \dashv |\cdot|_n$ we know, by observation 54, that the formulations given in the statement of this lemma are equivalent and that they both follow if we can demonstrate that each map obtained by applying Th_n to a primitive t -extension is again a t -extension. So consider the primitive t -extension $\Delta_k^a[m]' \xrightarrow{\subseteq_e} \Delta_k^a[m]''$ and observe that we have two cases:

- $n \geq m - 1$ In this case, all of the simplices of $\Delta_k^a[m]'$ and $\Delta_k^a[m]''$ of dimension $> n$ are already thin, therefore $\text{Th}_n(\Delta_k^a[m]') = \Delta_k^a[m]'$, $\text{Th}_n(\Delta_k^a[m]'') = \Delta_k^a[m]''$ and, consequently, Th_n maps $\Delta_k^a[m]' \xrightarrow{\subseteq_e} \Delta_k^a[m]''$ to itself.
- $n < m - 1$ We know that $\Delta_k^a[m]'$ and $\Delta_k^a[m]''$ differ only in as much as the $(m - 1)$ -simplex δ_k^m is not thin in the former and is thin in the latter. However, since $n < m - 1$ we know that δ_k^m is thin in both of $\text{Th}_n(\Delta_k^a[m]')$ and $\text{Th}_n(\Delta_k^a[m]'')$ and thus that these two stratified sets are identical. It follows that Th_n maps $\subseteq_e: \Delta_k^a[m]' \longrightarrow \Delta_k^a[m]''$ to the identity on $\text{Th}_n(\Delta_k^a[m]') = \text{Th}_n(\Delta_k^a[m]'')$.

In either case, the resulting map is a t -extension as required. $\square$

Lemma 151. *If A is an n -trivial pre-complicial set then $X \Rightarrow A$ and $A \Leftarrow Y$ are also n -trivial for any stratified sets $X, Y \in \underline{\text{Strat}}$.*

Proof. The results for $X \Rightarrow A$ and $A \Leftarrow Y$ are dual so we only consider the former. By definition, $\underline{\text{Strat}}_n$ is a reflective full subcategory of $\underline{\text{Strat}}$ with reflector Th_n , so it follows that a stratified set is n -trivial if and only if it is orthogonal to each entire inclusion $Z \xrightarrow{\subseteq_e} \text{Th}_n(Z)$. Now, we know that $X \Rightarrow *$ is a partial right adjoint to $X \otimes -$ on the category of pre-complicial sets, so taking adjoint transposes we see that $X \Rightarrow A$ is orthogonal to $Z \xrightarrow{\subseteq_e} \text{Th}_n(Z)$ if and only if A is orthogonal to $X \otimes Z \xrightarrow{\subseteq_e} X \otimes \text{Th}_n(Z)$. However, we have a sequence of entire inclusions

$$\begin{aligned} \text{Th}_n(X \otimes Z) &\subseteq_e \text{Th}_n(X \otimes \text{Th}_n(Z)) && \text{applying } \text{Th}_n \text{ to } X \otimes Z \subseteq_e X \otimes \text{Th}_n(Z), \\ &\subseteq_e \text{Th}_n(\text{Th}_n(X \otimes Z)) && \text{applying } \text{Th}_n \text{ to } X \otimes \text{Th}_n(Z) \subseteq_e \text{Th}_n(X \otimes Z) \\ & && \text{of lemma 133,} \\ &\subseteq_e \text{Th}_n(X \otimes Z) && \text{Th}_n \text{ is idempotent.} \end{aligned}$$

from which we see that $\text{Th}_n(X \otimes Z) = \text{Th}_n(X \otimes \text{Th}_n(Z))$ and thus that the inclusion of the last sentence is Th_n -invertible. It follows that the n -trivial set A is perpendicular to that inclusion for each $Z \in \underline{\text{Strat}}$ and thus that $X \Rightarrow A$ is n -trivial as required. $\square$

Observation 152. Let $\underline{\text{Pcs}}_n = \underline{\text{Pcs}} \cap \underline{\text{Strat}}_n$ denote the full subcategory of n -trivial pre-complicial sets in $\underline{\text{Pcs}}$. A stratified set is in $\underline{\text{Pcs}}_n$ iff it is orthogonal to the maps in the FP-regulus which is the union of the set of primitive t -extensions and the set $\{\Delta[m] \xrightarrow{\subseteq_e} \Delta[m]_t \mid m \in \mathbb{N} \text{ and } m > n\}$. It follows that $\underline{\text{Pcs}}_n$ is an LFP-category and that it is a reflective full subcategory of $\underline{\text{Strat}}$.

Now, lemma 151 implies that $X \Rightarrow *$ (resp. $* \Leftarrow Y$) restricts to a partial right adjoint to $X \otimes -$ (resp. $- \otimes Y$) on $\underline{\text{Pcs}}_n$, so it follows that we may apply Day's reflection theorem, just as in theorem 148, to reflect the monoidal structure of $\underline{\text{Strat}}$ to a monoidal biclosed structure on $\underline{\text{Pcs}}_n$.

8. COMPLICIAL SETS

8.1. Introducing Complicial Sets.

Definition 153 (primitive f-extensions). The class of *primitive f-extensions* is the union of the following classes of stratified maps

- a) the primitive t-extensions,
- b) the inclusions

$$\Lambda_k^a[n] \xrightarrow{\subseteq_r} \Delta_k^a[n]$$

(cf. notation 105) for $n \geq 2$ and $k = 1, 2, \dots, n-1$ which we call *admissible horn extensions*), and

- c) the unique surjection:

$$\Delta[1]_t \xrightarrow{d} \Delta[0]$$

Definition 154 (complicial set). A stratified set X is said to be a *complicial set* if it is orthogonal to each primitive f-extension. Let $\underline{\mathbf{Cs}}$ denote the full subcategory of $\underline{\mathbf{Strat}}$ whose objects are the complicial sets. Of course, since every primitive t-extension is a primitive f-extension it follows that every complicial set is pre-complicial or, in other words, that $\underline{\mathbf{Cs}}$ is a full subcategory of $\underline{\mathbf{Pcs}}$.

As in definition 121, we know that the stratified sets $\Delta_k^a[n]', \Delta_k^a[n]'', \Delta_k^a[n], \Lambda_k^a[n], \Delta[1]$ and $\Delta[0]$ are all finitely presentable in $\underline{\mathbf{Strat}}$ (cf observation 108). It follows, therefore, that the set of primitive f-extensions is also an FP-regulus and so, by observation 59, we see that $\underline{\mathbf{Cs}}$ is an LFP-category which is reflective in $\underline{\mathbf{Strat}}$ (with associated idempotent monad (L_c, η^c)).

We let $\mathbb{T}_{\underline{\mathbf{Cs}}}$ denote the LE-theory of complicial sets, which we construct from $\mathbb{T}_{\underline{\mathbf{Strat}}}$, the LE-theory of stratified sets, by adding the set of primitive f-extensions to its FP-regulus.

Observation 155 (describing complicial sets explicitly). More explicitly, by Yoneda's lemma a pre-complicial set A satisfies these conditions if and only if

- (a) for each $(n-1)$ -dimensional k -horn ($n \geq 2$ and $0 < k < n$) $\{x_i \mid i \neq k\}$ in X which is admissible (in the sense of observation 106) there is a unique thin n -simplex $x \in tX$, called a *thin filler*, such that $x \cdot \delta_i^n = x_i$ for $i \neq k$, and
- (b) each thin 1-simplex of X is degenerate.

Definition 156 (f-extensions). We call the L_c -invertible stratified maps *f-extensions*. For more on the properties of f-extensions please consult subsection 3.1. We will also use the term *f-almost* as a synonym for L_c -almost (cf. observation 53).

Observation 157 (duals of complicial sets). We can extend the stratified isomorphisms of observation 107 to get a further family of stratified isomorphisms:

$$\Delta_{n-k}^a[n] \xrightarrow{\cong} (\Delta_k^a[n])^\circ \tag{32}$$

Furthermore, it is a matter of a simple calculation to demonstrate that the stratified subset $(\Lambda_k^a[n])^\circ \subseteq_r (\Delta_k^a[n])^\circ$ corresponds to the subset $\Lambda_{n-k}^a[n] \subseteq_r \Delta_{n-k}^a[n]$ under this isomorphism.

In other words, we have a commutative diagram

$$\begin{array}{ccc} \Lambda_{n-k}^a[n] & \xrightarrow{\subseteq_r} & \Delta_{n-k}^a[n] \\ \cong \downarrow & & \downarrow \cong \\ (\Lambda_k^a[n])^\circ & \xrightarrow{\subseteq_r} & (\Delta_k^a[n])^\circ \end{array}$$

for each $n \geq 2$ and $1 \leq k < n$. So the dual of each admissible horn extension is again (isomorphic to) an admissible horn extension.

So, by an argument identical to that of observation 125, we see that A is a complicial set if and only if A° is. Equivalently, $f: X \longrightarrow Y$ is an f-extension if and only if $f^\circ: X^\circ \longrightarrow Y^\circ$ is.

8.2. Pasting Squares and Filling Lemmas. The following lemmas are mostly a direct consequence of the properties of f-extensions listed in observation 52. They provide us with higher level tools for constructing a whole range of complex f-extensions and, as such, they will be used repeatedly in the sequel.

Observation 158 (pasting squares). Suppose that $f: X \hookrightarrow Z$ and $g: Y \hookrightarrow Z$ are stratified inclusions and that $Z = f(X) \cup g(Y)$ then the pullback

$$\begin{array}{ccc} X \times_f Y & \xrightarrow{\pi_X} & X \\ \pi_Y \downarrow \lrcorner & & \downarrow f \\ Y & \xrightarrow{g} & Z \end{array}$$

is also a pushout in Strat; such a square is known as a *pasting square*. It follows, by observation 52(8) that if the upper horizontal map π_X is an f-extension then so is the lower one g . Usually we'll apply this result in one of the following cases:

- (a) f is an arbitrary stratified inclusion and g is the inclusion associated with a subset $Y \subseteq_s Z$, in which case our pasting square reduces to

$$\begin{array}{ccc} f^{-1}(Y) & \xrightarrow{\subseteq_s} & X \\ f \downarrow \lrcorner & & \downarrow f \\ Y & \xrightarrow{\subseteq_s} & f(X) \cup Y \end{array}$$

and we infer that $Y \xrightarrow{\subseteq_s} f(X) \cup Y$ is an f-extension from the fact that $f^{-1}(Y) \xrightarrow{\subseteq_s} X$ is an f-extension (by observation 52(8)).

- (b) f is the inclusion associated with $X \subseteq_s Z$ and g is the inclusion associated with a subset $Y \subseteq_s Z$, in which case our pasting square reduces to

$$\begin{array}{ccc} X \cap Y & \xrightarrow{\subseteq_s} & X \\ \subseteq_s \downarrow \lrcorner & & \downarrow \subseteq_s \\ Y & \xrightarrow{\subseteq_s} & X \cup Y \end{array}$$


and we infer that $Y \xrightarrow{\subseteq_s} X \cup Y$ is an f-extension from the fact that $X \cap Y \xrightarrow{\subseteq_s} X$ is an f-extension (by observation 52(8)).

Lemma 159 (pasting lemma for f-extensions). *Suppose that $f: X \longrightarrow Y$ is a stratified map and that we have stratified subsets $X^{(i)} \subseteq_s X$ and $Y^{(i)} \subseteq_s Y$ with $X = \bigcup_{i=1}^r X^{(i)}$, $Y = \bigcup_{i=1}^r Y^{(i)}$ and $X^{(i)} \subseteq f^{-1}(Y^{(i)})$ for $i = 1, \dots, r$ such that f restricts to*

- (a) *an f-extension $f^{(i)}: X^{(i)} \longrightarrow Y^{(i)}$ for each i and*
- (b) *an f-extension $f^{(i,j)}: X^{(i)} \cap X^{(j)} \longrightarrow Y^{(i)} \cap Y^{(j)}$ for each $i < j$*

then $f: X \longrightarrow Y$ itself is also an f-extension.

Proof. From observations 32 and 109 we know that the unions $\bigcup_l X^{(l)}$ and $\bigcup_l Y^{(l)}$ may be expressed as wide pushouts as depicted in display (3). Furthermore, the various restrictions of the statement fit together into commuting diagrams of the form



for each $i < j$. These demonstrate that $f: X \longrightarrow Y$ is the unique map induced by the restrictions at the top of our diagram, each one of which is an f-extension (by assumption), between the wide pushouts at the bottom. Thus the postulated result follows from the fact that the class of f-extensions is closed under colimits (see observation 52). $\square$

Corollary 160. *Suppose that X is a stratified subset of $Y \in \underline{\text{Strat}}$ and that $Y^{(i)}$ ($i = 1, \dots, r$) is a family of stratified subsets of Y such that*

- (a) *the inclusion $X \xrightarrow{\subseteq_s} X \cup Y^{(i)}$ is an f-extension for each i and*
- (b) *the inclusion $X \xrightarrow{\subseteq_s} X \cup (Y^{(i)} \cap Y^{(j)})$ is an f-extension for each $i < j$*

then the inclusion $X \xrightarrow{\subseteq_s} X \cup (\bigcup_{i=1}^r Y^{(i)})$ is also an f-extension.

Proof. Apply lemma 159 to the inclusion map $X \xrightarrow{\subseteq_s} Y$ and the families of subsets $\{X\}_{i=1}^r$ and $\{X \cup Y^{(i)}\}_{i=1}^r$, for which the conditions given in that lemma simply reduce to those given in the statement of this corollary. $\square$

Lemma 161 (minor filling lemma). *If X is a regular stratified subset of $\Delta_k^a[n]$ and*

- (a) δ_{k-1}^n and δ_{k+1}^n are elements of X and
- (b) *if μ is a k -divided r -simplex in $\Delta_k^a[n]$, $l \in [r]$ is the unique integer such that $\mu(l) = k$ and $\mu \circ \delta_l^r$ is an element of the subset X then μ is also an element of X ,*

then the inclusion map $X \xrightarrow{\subseteq_r} \Delta_k^a[n]$ is an f-extension.

Proof. By induction on n , using lemma 159:

n=2: The only stratified subset satisfying the conditions of the lemma is $\Lambda_1^a[2] \subseteq_r \Delta_1^a[2]$ which is a primitive f-extension.

the inductive case: Suppose that if $m < n$ then the result is true for all subsets of $\Delta_l^a[m]$ ($l = 1, \dots, m-1$) satisfying the conditions of the lemma.

Assume that $X \subseteq_r \Delta_k^a[n]$ satisfies the conditions of the lemma. If $\text{id}_{[n]} \in X$ then $X = \Delta_k^a[n]$ and there is nothing to prove, so assume that $\text{id}_{[n]} \notin X$ from which it follows, by condition (b), that $\delta_k^n \notin X$. In other words, X is a stratified subset of $\Delta_k^a[n]$.

Suppose now that μ is a non-degenerate k -divided m -simplex in $\Delta_k^a[n]$, that is to say a face operator $\mu: [m] \longrightarrow [n]$ with $k-1, k, k+1 \in \text{im}(\mu)$ and let $l \in [m]$ be the unique number such that $\mu(l) = k$, for which we also know that $k-1 = \mu(l-1)$ and $k+1 = \mu(l+1)$. Clearly an element ν of $\Delta[m]$ is l -divided if and only if $\Delta(\mu)(\nu) = \mu \circ \nu \in \Delta[n]$ is k -divided. It follows that the simplicial map $\Delta(\mu): \Delta[m] \longrightarrow \Delta[n]$ extends to a regular stratified map from $\Delta_l^a[m]$ to $\Delta_k^a[n]$. So consider the inverse image $\Delta(\mu)^{-1}(X) \subseteq_r \Delta_l^a[m]$ of $X \subseteq_r \Delta_k^a[n]$, which we know to be a regular stratified subset by observation 101. It is also easily established that this inverse image satisfies the conditions (a) and (b) in the statement of this lemma:

- (i) Given that μ is k -divided with $\mu(l) = k$ and that $l-1 \notin \text{im}(\delta_{l-1}^m)$, we may infer that $k-1 \notin \text{im}(\mu \circ \delta_{l-1}^m)$ and, consequently, that $\mu \circ \delta_{l-1}^m = \Delta(\mu)(\delta_{l-1}^m)$ is an $(m-1)$ -dimensional face of δ_{k-1}^n in $\Delta_k^a[n]$. However, by assumption (a) for X , we know that δ_{k-1}^n is in X as is each of its faces, including in particular $\Delta(\mu)(\delta_{l-1}^m)$. It follows that δ_{l-1}^m is in the inverse image $\Delta(\mu)^{-1}(X)$ as required. An identical argument demonstrates that δ_{l+1}^m is also in $\Delta(\mu)^{-1}(X)$.
- (ii) Suppose that ν is an l -divided r -simplex in $\Delta_l^a[m]$, that $j \in [r]$ has $\nu(j) = l$ and that $\nu \circ \delta_j^r$ is in $\Delta^{-1}(X)$. By definition, this latter condition holds iff $\Delta(\mu)(\nu \circ \delta_j^r) = \mu \circ (\nu \circ \delta_j^r) = (\mu \circ \nu) \circ \delta_j^r$ is an element of X . Furthermore, we know that μ is k -divided with $\mu(l) = k$ and that ν is l -divided with $\nu(j) = l$ so it follows that $\mu \circ \nu$ is k -divided with $\mu \circ \nu(j) = k$. Now apply condition (b) for the subset $X \subseteq_r \Delta_k^a[n]$ to demonstrate that if $(\mu \circ \nu) \circ \delta_j^r$ is an element of X then so is $\mu \circ \nu = \Delta(\mu)(\nu)$. From this we may infer that ν is an element of $\Delta(\mu)^{-1}(X)$ as required.

It follows, by our inductive hypothesis, that the inclusion $\Delta(\mu)^{-1}(X) \xrightarrow{\subseteq_r} \Delta_l^a[m]$ is an f-extension and since it features in the pasting square

$$\begin{array}{ccc} \Delta(\mu)^{-1}(X) & \xrightarrow{\subseteq_r} & \Delta_l^a[m] \\ \Delta(\mu) \downarrow & & \downarrow \Delta(\mu) \\ X & \xrightarrow[\subseteq_r]{} & \Delta(\mu)(\Delta_l^a[m]) \cup X \end{array}$$

we may infer that the inclusion $X \xrightarrow{\subseteq_r} \Delta(\mu)(\Delta_l^a[m]) \cup X$ is also an f-extension (by observation 158(a)). Furthermore, since we know that $\Delta(\mu): \Delta_l^a[m] \longrightarrow \Delta_k^a[n]$ is regular, we may apply observation 101 to show that the image $\Delta(\mu)(\Delta_l^a[m])$ is a regular subset of $\Delta_k^a[n]$ and indeed that it must be $\{\mu\}^* \subseteq_r \Delta_k^a[n]$, the regular stratified subset generated by the m -simplex $\mu \in \Delta_k^a[n]$. To summarise, we have established that the inclusion map $X \xrightarrow{\subseteq_r} X \cup \{\mu\}^*$ is an f-extension for each k -divided operator $\mu: [m] \longrightarrow [n]$.

Now, for each $i \neq k-1, k, k+1$ we know that δ_i^n is k -divided, so let $Y^{(i)} \stackrel{\text{def}}{=} \{\delta_i^n\}^*$ and the result of the last paragraph implies that $X \xrightarrow{\subseteq_r} X \cup Y^{(i)}$ is an f-extension. Furthermore, if $i < j$ ($i, j \neq k-1, k, k+1$) then we have $Y^{(i)} \cap Y^{(j)} = \{\delta_i^n\}^* \cap \{\delta_j^n\}^* = \{\delta_j^n \circ \delta_i^{n-1}\}^*$

(cf. notation 36) and $\delta_j^n \circ \delta_i^{n-1}$ is also k -divided thus the result of the last paragraph implies that $X \xrightarrow{\subseteq_r} X \cup (Y^{(i)} \cap Y^{(j)})$ is an f-extension as well. In other words, the family $\{Y^{(i)}\}_{i \neq k-1, k, k+1}$ satisfies the conditions of corollary 160, from which it follows that the inclusion map $X \xrightarrow{\subseteq_r} X \cup \left(\bigcup_{i \neq k-1, k, k+1} Y^{(i)} \right)$ is also an f-extension.

However, the faces δ_{k-1}^n and δ_{k+1}^n are elements of $X \subseteq_r \Lambda_k^a[n]$ (by condition (a) in the statement of this lemma) and the faces δ_i^n ($i \neq k-1, k, k+1$) are the remaining non-degenerate $(n-1)$ -simplices of $\Lambda_k^a[n]$, from which it follows that $X \cup \left(\bigcup_{i \neq k-1, k, k+1} Y^{(i)} \right) = \Lambda_k^a[n]$. So the result of the last paragraph simply demonstrated that the inclusion $X \xrightarrow{\subseteq_r} \Lambda_k^a[n]$ is an f-extension, which we now compose with the primitive f-extension $\Lambda_k^a[n] \xrightarrow{\subseteq_r} \Delta_k^a[n]$ in order to prove that the resulting inclusion $X \xrightarrow{\subseteq_r} \Delta_k^a[n]$ is also an f-extension as stated. $\square$

The following variant of this result will prove particularly useful:

Corollary 162. *Suppose that N is a stratified set in the subcategory Simplex of definition 104 and that $X \subseteq_r N$ is a regular subset which satisfies the conditions of lemma 161 and these also satisfy*

- (a) $\Delta_k^a[n] \subseteq N$, in other words every k -divided simplex of $\Delta[n]$ is in tN , and
- (b) whenever μ is a k -divided r -simplex in $\Delta[n]$, l is the unique integer with $\mu(l) = k$ and $\mu \circ \delta_l^r \in tN \setminus X$ then both of the faces $\mu \circ \delta_{l-1}^r$ and $\mu \circ \delta_{l+1}^r$ are also in tN .

then the inclusion $X \xrightarrow{\subseteq_r} N$ is an f-extension.

Proof. Consider the stratified subsets $X \subseteq_r N$ and $\Delta_k^a[n] \subseteq_e N$. Since X is a regular subset of N it follows that $X \cap \Delta_k^a[n]$ is a regular subset of $\Delta_k^a[n]$, furthermore $X \cap \Delta_k^a[n]$ satisfies the conditions of lemma 161 since it has the same underlying simplicial set as X which does so by assumption. Consequently, we may apply that lemma to demonstrate that the inclusion $X \cap \Delta_k^a[n] \xrightarrow{\subseteq_r} \Delta_k^a[n]$ is an f-extension and, by observation 158(b), we know that this map features in a pasting square

$$\begin{array}{ccc} X \cap \Delta_k^a[n] & \xrightarrow{\subseteq_r} & \Delta_k^a[n] \\ \downarrow & & \downarrow \\ X & \xrightarrow{\subseteq_r} & X \cup \Delta_k^a[n] \end{array}$$

from which we may conclude that the inclusion $X \xrightarrow{\subseteq_r} X \cup \Delta_k^a[n]$ is also an f-extension.

We can also show that the inclusion $X \cup \Delta_k^a[n] \xrightarrow{\subseteq_r} N$ is a t-extension. To this end, suppose that ν is thin in N and not thin in $X \cup \Delta_k^a[n]$. Then ν is necessarily non-degenerate, since otherwise it would be thin in $X \cup \Delta_k^a[n]$.

Notice that if a simplex ν of $\Delta[n]$ has $k-1 \notin \text{im}(\nu)$ (resp. $k+1 \notin \text{im}(\nu)$) then it is a face of δ_{k-1}^n (resp. δ_{k+1}^n) which is an element of X , by assumption (a) of lemma 161. But the stratified subset X is closed in N under the right action of Δ , so if it contains the simplex δ_{k-1}^n (resp. δ_{k+1}^n) then it must also contain all of its faces and it follows that $\nu \in X$. Furthermore X is a **regular** subset of N , so in such a case ν is thin in X iff it is thin in N .

So suppose, for a contradiction, that $k-1$ (resp. $k+1$) is not an element of $\text{im}(\nu)$, then we could infer (by the result of the last paragraph) that ν is in X and that it thin there, since we assumed that it was thin in N . However, this would contradict our assumption that ν was **not** thin in $X \cup \Delta_k^a[n]$.

It follows that both $k-1$ and $k+1$ are elements of $\text{im}(\nu)$. However, k is not in $\text{im}(\nu)$, because if it were ν would be k -divided and thus thin in $X \cup \Delta_k^a[n]$. So let μ be the unique

non-degenerate simplex of $\Delta[n]$ which has $\text{im}(\mu) = \text{im}(\nu) \cup \{k\}$ and let l be the unique integer such that $\mu(l) = k$.

By construction μ is k -divided, in particular it is k -admissible in $X \cup \Delta_k^a[n]$, and $\mu \circ \delta_l^n = \nu$. We assumed that ν was thin in N therefore, by assumption (b) of this corollary, we can infer that both of $\mu \circ \delta_{l-1}^n$ and $\mu \circ \delta_{l+1}^n$ are also thin in N . But $k-1$ (resp. $k+1$) is not an element of $\text{im}(\mu \circ \delta_{l-1}^n)$ (resp. $\text{im}(\mu \circ \delta_{l+1}^n)$) so, by our observation of a few paragraphs ago, both of these simplices are in X and they are thin in there, thus they are thin in $X \cup \Delta_k^a[n]$.

But this is precisely what we need in order to show that $X \cup \Delta_k^a[n] \hookrightarrow N$ is a t-extension by invoking observation 126, and all t-extensions are f-extensions (cf. observation 52(4)).

So both of $X \hookrightarrow X \cup \Delta_k^a[n]$ and $X \cup \Delta_k^a[n] \hookrightarrow N$ are f-extensions and it follows that their composite $X \hookrightarrow N$ is also an f-extension as required. $\square$

8.3. Tensor Products and Complicial Sets. First we show that complicial sets are all well tempered as indicated in section 6.2. Then we push on to prove a range of particularly important results with respect to complicial sets and their interaction with the biclosed monoidal structure introduced in section 7.

Lemma 163. *Every complicial set is well tempered (cf. definition 117). In other words if A is a complicial set then every simplex $a \in A$ which is pre-degenerate at k is degenerate at k .*

Proof. By induction on $n = \dim(a)$:

$n = 1$: This is simply (equivalent to) the axiom that states that a complicial set is orthogonal to $d: \Delta[1]_t \longrightarrow \Delta[0]$, as expressed in the form given in observation 155(b).

inductive case: Suppose that the given result holds for every r -simplex $a \in A$ with $1 \geq r < n$ and each $k \in [r]$.

Let $a \in A$ be an n -simplex which is pre-degenerate at k , assume for the moment that $k > 0$ (we'll consider the case $k = 0$ later on) and consider the $(n-1)$ -dimensional faces $a \cdot \delta_i^n$ of a :

- **$i < k$** Let $\alpha: [r] \longrightarrow [n-1]$ be a face operator with $k-1, k \in \text{im}(\alpha)$ then, since $i \leq k-1$, we know that $k, k+1 \in \text{im}(\delta_i^n \circ \alpha)$; it follows, since a is pre-degenerate at k , that $(a \cdot \delta_i^n) \cdot \alpha = a \cdot (\delta_i^n \circ \alpha)$ is thin. Consequently $a \cdot \delta_i^n$ is pre-degenerate at $k-1$ and so, by our inductive hypotheses, it is degenerate at $k-1$; it follows that $a \cdot \delta_i^n = ((a \cdot \delta_i^n) \cdot \delta_k^{n-1}) \cdot \sigma_{k-1}^{n-2} = a \cdot (\delta_i^n \circ \delta_k^{n-1} \circ \sigma_{k-1}^{n-2})$.

Applying the simplicial identities of observation 4 (twice), under the assumption that $i < k$, we see that $\delta_i^n \circ \delta_k^{n-1} \circ \sigma_{k-1}^{n-2} = \delta_{k+1}^n \circ \delta_i^{n-1} \circ \sigma_{k-1}^{n-2} = \delta_{k+1}^n \circ \sigma_k^{n-1} \circ \delta_i^n$ and so the equation at the end of the last paragraph becomes $a \cdot \delta_i^n = a \cdot (\delta_{k+1}^n \circ \sigma_k^{n-1} \circ \delta_i^n) = ((a \cdot \delta_{k+1}^n) \cdot \sigma_k^{n-1}) \cdot \delta_i^n$.

- **$i > k+1$** Let $\alpha: [r] \longrightarrow [n-1]$ be a face operator with $k, k+1 \in \text{im}(\alpha)$ then, since $i > k+1$, we know that $k, k+1 \in \text{im}(\delta_i^n \circ \alpha)$; it follows, since a is pre-degenerate at k , that $(a \cdot \delta_i^n) \cdot \alpha = a \cdot (\delta_i^n \circ \alpha)$ is thin. Consequently $a \cdot \delta_i^n$ is pre-degenerate at k and so, by our inductive hypotheses, it is degenerate at k ; it follows that $a \cdot \delta_i^n = ((a \cdot \delta_i^n) \cdot \delta_{k+1}^{n-1}) \cdot \sigma_k^{n-2} = a \cdot (\delta_i^n \circ \delta_{k+1}^{n-1} \circ \sigma_k^{n-2})$.

Applying the simplicial identities of observation 4 (twice), under the assumption that $i > k+1$, we see that $\delta_i^n \circ \delta_{k+1}^{n-1} \circ \sigma_k^{n-2} = \delta_{k+1}^n \circ \delta_{i-1}^{n-1} \circ \sigma_k^{n-2} = \delta_{k+1}^n \circ \sigma_k^{n-1} \circ \delta_i^n$ and so the equation at the end of the last paragraph becomes $a \cdot \delta_i^n = a \cdot (\delta_{k+1}^n \circ \sigma_k^{n-1} \circ \delta_i^n) = ((a \cdot \delta_{k+1}^n) \cdot \sigma_k^{n-1}) \cdot \delta_i^n$.

- $i = k + 1$ Since $\sigma_k^{n-1} \circ \delta_{k+1}^n = \text{id}_{[n-1]}$ we have $((a \cdot \delta_{k+1}^n) \cdot \sigma_k^{n-1}) \cdot \delta_{k+1}^n = (a \cdot \delta_{k+1}^n) \cdot (\sigma_k^{n-1} \circ \delta_{k+1}^n) = (a \cdot \delta_{k+1}^n) \cdot \text{id}_{[n-1]} = a \cdot \delta_{k+1}^n$.

Now we are in a position to compare the simplices a and $(a \cdot \delta_{k+1}^n) \cdot \sigma_k^{n-1}$; the equations derived above simply state that the corresponding horns $\{a \cdot \delta_i^n \mid i \neq k\}$ and $\{((a \cdot \delta_{k+1}^n) \cdot \sigma_k^{n-1}) \cdot \delta_i^n \mid i \neq k\}$ are in fact identical. Furthermore, since every k -divided operator α has $k, k+1 \in \text{im}(\alpha)$ and a is pre-degenerate at k , we may infer that a is k -admissible. In summary, a and $(a \cdot \delta_{k+1}^n) \cdot \sigma_k^{n-1}$ are both thin fillers of the admissible horn $\{a \cdot \delta_i^n \mid i \neq k\}$.

However we know that thin fillers of admissible horns in A are unique, since it is a complicial set, therefore $a = (a \cdot \delta_{k+1}^n) \cdot \sigma_k^{n-1}$ and thus a is degenerate at k .

All that remains is to prove that the result also holds when $k = 0$. To do so, observe that an n -simplex $a \in A$ is degenerate (resp. pre-degenerate) at k if and only if it is degenerate (resp. pre-degenerate) at $(n - 1 - k)$ when considered as a simplex in the dual A° . So if our simplex a is pre-degenerate at 0 as an element of A then it is pre-degenerate at $(n - 1)$ when considered as an element of the dual complicial set A° . Applying the result we've already proved, we may infer that a is degenerate at $(n - 1)$ in A° and thus is degenerate at 0 in A as required. $\square$

Corollary 164. *For each stratified set Y the stratified map*

$$\Delta[1]_t \otimes Y \xrightarrow{d \otimes Y} Y \quad (33)$$

is an f -extension.

Observation 165. Before we proceed with this proof, we introduce a little notation for the elements of $\Delta[1]$. For $k \in [r + 1]$ let $\rho_k^r : [r] \longrightarrow [1]$ be the simplicial operator given by

$$\rho_k^r(i) = \begin{cases} 0 & \text{if } i < k, \\ 1 & \text{if } i \geq k. \end{cases}$$

for which the following, easily demonstrated, identities will be of use in this proof:

$$\begin{aligned} \rho_{k+1}^{r+1} \circ \delta_k^{r+1} &= \rho_k^r \\ \rho_k^{r+1} \circ \delta_k^{r+1} &= \rho_k^r \end{aligned} \quad (34)$$

Proof. We know that the map $d : \Delta[1]_t \longrightarrow \Delta[0]$ has $\Delta(\varepsilon_0^1) : \Delta[0] \longrightarrow \Delta[1]_t$ as a right inverse so, by the functoriality of $\otimes$, we also know that $\Delta(\varepsilon_0^1) \otimes Y$ is a right inverse of $d \otimes Y$. We'll actually prove that $\Delta(\varepsilon_0^1) \otimes Y$ is an f -extension and then appeal to observation 52(6) to prove that $d \otimes Y$ is also an f -extension.

Since $d \otimes Y$ is left inverse to $\Delta(\varepsilon_0^1) \otimes Y$, we may extend any stratified map $f : Y \longrightarrow A$ along $\Delta(\varepsilon_0^1) \otimes Y$ by simply composing it with $d \otimes Y$. To complete our proof we must demonstrate that if A is a complicial set then this extension is unique or, in other words, that if $g, g' : \Delta[1]_t \otimes Y \longrightarrow A$ are any pair of stratified maps with $g \circ (\Delta(\varepsilon_0^1) \otimes Y) = g' \circ (\Delta(\varepsilon_0^1) \otimes Y)$ then we have $g = g'$.

Observe that in order to verify this latter condition it suffices to show that for any stratified $g : \Delta[1]_t \otimes Y \longrightarrow A$ with A complicial we have $g(\rho_k^r, y) = g(\rho_{k+1}^r, y)$ for each $k \in [r]$; this follows because any simplicial operator $\rho : [r] \longrightarrow [1]$ is of the form ρ_i^r for some $i \in [r + 1]$ so the stated equality (for each k) would give $g(\rho, y) = g(\rho_i^r, y) = g(\rho_{i-1}^r, y) = g(\rho_{i-2}^r, y) = \dots = g(\rho_0^r, y)$. However $\rho_0^r = \varepsilon_0^1 \circ \eta^r$ so we have $(\rho_0^r, y) = (\Delta(\varepsilon_0^1) \otimes Y)(y)$ and it follows that if g and g' are as in the previous paragraph we would have $g(\rho, y) = (g \circ (\Delta(\varepsilon_0^1) \otimes Y))(y) = (g' \circ (\Delta(\varepsilon_0^1) \otimes Y))(y) = g'(\rho, y)$ as required.

To prove the equality $g(\rho_k^r, y) = g(\rho_{k+1}^r, y)$ (for a given $k \in [r]$) consider the $(r+1)$ -simplex $(\rho_{k+1}^{r+1}, y \cdot \sigma_k^r)$ in $\Delta[1]_t \otimes Y$ which, by the equalities in (34), has:

$$\begin{aligned} (\rho_{k+1}^{r+1}, y \cdot \sigma_k^r) \cdot \delta_k^{r+1} &= (\rho_{k+1}^{r+1} \circ \delta_k^{r+1}, y \cdot (\sigma_k^r \circ \delta_k^{r+1})) = (\rho_k^r, y) \\ (\rho_{k+1}^{r+1}, y \cdot \sigma_k^r) \cdot \delta_{k+1}^{r+1} &= (\rho_{k+1}^{r+1} \circ \delta_{k+1}^{r+1}, y \cdot (\sigma_k^r \circ \delta_{k+1}^{r+1})) = (\rho_{k+1}^r, y) \end{aligned} \quad (35)$$

Now observe that we have $(d \otimes Y)(\rho_{k+1}^{r+1}, y \cdot \sigma_k^r) = y \cdot \sigma_k^r$ which is degenerate, and thus pre-degenerate, at k in Y . However we know that d is regular from which we may infer, by observation 132, that $d \otimes Y$ is also regular and it follows that $(\rho_{k+1}^{r+1}, y \cdot \sigma_k^r)$ is pre-degenerate at k in $\Delta[1]_t \otimes Y$ (as observed in definition 116). Finally, we see that $g(\rho_{k+1}^{r+1}, y \cdot \sigma_k^r)$ is also pre-degenerate at k in A and so, since A is complicial, we may apply lemma 163 to show that this simplex is, in fact, degenerate at k . Now we may infer our desired equality since:

$$\begin{aligned} g(\rho_k^r, y) &= g(\rho_{k+1}^{r+1}, y \cdot \sigma_k^r) \cdot \delta_k^{r+1} \quad \text{from first equality in (35)} \\ &= g(\rho_{k+1}^{r+1}, y \cdot \sigma_k^r) \cdot \delta_{k+1}^{r+1} \quad \text{since } g(\rho_{k+1}^{r+1}, y \cdot \sigma_k^r) \text{ is degenerate at } k. \\ &= g(\rho_{k+1}^r, y) \quad \text{from second equality in (35)} \end{aligned} \quad \square$$

Now we present the primary combinatorial result of this section, and indeed of this work, that being lemma 169. Firstly however, we introduce some notation and prove an associated technical lemma.

Notation 166. Suppose that N and M are stratified sets in Simplex and that $\odot$ represents any of the bifunctors $\times$, $\otimes$ or $\boxtimes$ on Strat then in the sequel we adopt the following notation for various useful regular subsets of these stratified sets and their tensor products:

- (i) As usual, let ∂N and ∂M denote the “boundaries” of N and M , that is their regular subsets consisting of those operators which are **not** degeneracy operators.
- (ii) Furthermore let $\partial(N \odot M)$ be the boundary of the tensor product $N \odot M$, which is its regular subset given by the Leibniz formula:

$$\partial(N \odot M) \stackrel{\text{def}}{=} (\partial N \odot M) \cup (N \odot \partial M)$$

In other words, a simplex (α, β) of $N \odot M$ is in $\partial(N \odot M)$ if and only if either α or β is **not** a degeneracy operator.

- (iii) Let $H(N \odot M)$ be the regular stratified subset of $N \odot M$ on those simplices which do **not** have $(n, 0)$ as a vertex. Recall that every simplex of $N \odot M$ is a face of some shuffle and that the $\leftarrow$ -minimal shuffle $(\alpha_1, \beta_1) = (\pi_1^{n,m}, \pi_2^{n,m})$ is the only one which has $(n, 0)$ as a vertex. Therefore $H(N \odot M)$ is precisely the regular stratified subset of $N \odot M$ generated by the set of shuffles $\{(\alpha_i, \beta_i) : 1 < i \leq \#(n, m)\}$ and

$$H(N \odot M) \cup \{(\alpha_1, \beta_1)\}^* = N \odot M$$

- (iv) Let $\partial H(N \odot M)$ denote the “boundary” of $H(N \odot M)$

$$\partial H(N \odot M) \stackrel{\text{def}}{=} H(N \odot M) \cap \partial(N \odot M)$$

in other words, the regular stratified subset of $H(N \odot M)$ consisting of those simplices (α, β) for which either α or β is **not** a degeneracy operator.

- (v) Given a natural number i with $1 \leq i \leq \#(n, m)$ let $H_i(N \odot M)$ denote the regular stratified subset given by

$$H_i(N \odot M) = \partial H(N \odot M) \cup \{(\alpha_{i'}, \beta_{i'}) \mid i' > i\}^*$$

So $H_{\#(n,m)}(N \odot M) = \partial H(N \odot M)$, $H_1(N \odot M) = H(N \odot M)$ and $H_{i-1}(N \odot M)$ is the regular stratified subset of $H(N \odot M)$ obtained by adjoining the shuffle (α_i, β_i) to $H_i(N \odot M)$ (in other words $H_{i-1}(N \odot M) = H_i(N \odot M) \cup \{(\alpha_i, \beta_i)\}^* \subseteq_r H(N \odot M)$).

Lemma 167. *Fix $n, m \in \mathbb{N}$ and suppose that (α, β) is a non-degenerate r -simplex of $\Delta[n] \times \Delta[m]$ which is a mediator as witnessed by some k with $0 < k < r$, furthermore assume that $(\alpha, \beta) \cdot \delta_k^r$ is in $H_{i-1}(\Delta[n] \times \Delta[m])$ then*

- (a) (α, β) is in $H_{i-1}(\Delta[n] \times \Delta[m])$ and
- (b) $(\alpha, \beta) \cdot \delta_{k-1}^r$ and $(\alpha, \beta) \cdot \delta_{k+1}^r$ are both in $H_i(\Delta[n] \times \Delta[m])$.

Proof. Let w be the largest integer for which the simple $(\alpha, \beta) \cdot \delta_k^r$ is a face of the shuffle (α_w, β_w) . By observation 48 we know that the associated operator $\gamma_w: [n-1] \longrightarrow [m]$ is given by the formula in display (6) of the proof of lemma 45(6), which in the case of the particular value $\gamma_w(\alpha(k-1))$ reduces to:

$$\gamma_w(\alpha(k-1)) = \min\{\beta \circ \delta_k^r(l) \mid l \in [r-1] \wedge \alpha \circ \delta_k^r(l) > \alpha(k-1)\}$$

Now since (α, β) is *non-degenerate* and satisfies the mediation condition at k we know that the (in)equalities

$$\alpha(k-1) = \alpha(k) < \alpha(k+1) \text{ and } \beta(k-1) < \beta(k) = \beta(k+1) \quad (36)$$

hold. In particular, we see that $\alpha \circ \delta_k^r(k) = \alpha(k+1) > \alpha(k-1)$ and $\alpha \circ \delta_k^r(k-1) = \alpha(k-1) \not> \alpha(k-1)$ so it clearly follows that $\gamma_w(\alpha(k-1)) = \beta(k+1)$. Applying these (in)equalities again, this time to β , we see that $\gamma_w(\alpha(k-1)) = \beta(k+1) > \beta(k-1) \geq 0$ so γ_w is a non-zero operator and therefore the corresponding shuffle (α_w, β_w) is not $(\pi_1^{n,m}, \pi_2^{n,m})$ and is thus an element of $H(\Delta[n] \times \Delta[m])$.

Applying lemma 45(5) we have $(\alpha, \beta) \cdot \delta_k^r = (\alpha_w, \beta_w) \cdot (\alpha \circ \delta_k^r + \beta \circ \delta_k^r) = ((\alpha_w, \beta_w) \cdot (\alpha + \beta)) \cdot \delta_k^r$, where the final equality is a consequence of the fact that $+: \Delta[n] \times \Delta[m] \longrightarrow \Delta[n+m]$ is a simplicial map. It follows that (α, β) and the face $(\alpha_w, \beta_w) \cdot (\alpha + \beta)$ agree at vertices $j \in [r]$ with $j \neq k$, and so they are equal if and only if they also agree at vertex k . However, by lemma 45(5), this is the case if and only if the inequalities $\gamma_w(\alpha(k)-1) \leq \beta(k) \leq \gamma_w(\alpha(k))$ of display (4) hold, which is certainly the case since we may apply the equalities of the last paragraph to show that $\gamma_w(\alpha(k)) = \gamma_w(\alpha(k-1)) = \beta(k+1) = \beta(k)$.

We may now dispense with the trivial case, wherein $(\alpha, \beta) \cdot \delta_k^r = (\alpha \circ \delta_k^r, \beta \circ \delta_k^r)$ is in the boundary $\partial H(\Delta[n] \times \Delta[m]) \subseteq H_{i-1}(\Delta[n] \times \Delta[m])$ which implies that one of $\alpha \circ \delta_k^r$ or $\beta \circ \delta_k^r$ is not surjective. However, consulting display 36 we know that $\alpha(k-1) = \alpha(k)$ and $\beta(k) = \beta(k+1)$ and it follows easily that $\text{im}(\alpha) = \text{im}(\alpha \circ \delta_k^r)$ and $\text{im}(\beta) = \text{im}(\beta \circ \delta_k^r)$, consequently either α or β is also non-surjective and thus (α, β) is in $\partial(\Delta[n] \times \Delta[m])$. Furthermore, as verified in the last paragraph, we know that (α, β) is a face of the shuffle $(\alpha_w, \beta_w) \in H(\Delta[n] \times \Delta[m])$ and it is thus an element of $\partial H(\Delta[n] \times \Delta[m]) \subseteq H_i(\Delta[n] \times \Delta[m])$ as are its faces $(\alpha, \beta) \cdot \delta_{k-1}^r$ and $(\alpha, \beta) \cdot \delta_{k+1}^r$ as required.

Otherwise, if $(\alpha, \beta) \cdot \delta_k^r$ is not in the boundary $\partial H(\Delta[n] \times \Delta[m])$ then, since it is an element of $H_{i-1}(\Delta[n] \times \Delta[m])$, there must exist an integer $u > i-1$ such that it is a face of the shuffle (α_u, β_u) . Now, by the maximality of w it follows that $w \geq u > i-1$ and therefore that (α_w, β_w) is in $H_{i-1}(\Delta[n] \times \Delta[m])$. However, we've already shown that (α, β) is a face of the shuffle (α_w, β_w) so it follows that it is an element of $H_{i-1}(\Delta[n] \times \Delta[m])$ as postulated in point (a) of the statement.

Now consider the face $(\alpha, \beta) \cdot \delta_{k-1}^r$ and observe that the inequality $\beta(k-1) < \beta(k)$ of display (36) implies that either:

- $\beta(k-1) \notin \text{im}(\beta \circ \delta_{k-1}^r)$ and thus that $(\alpha, \beta) \cdot \delta_{k-1}^r = (\alpha \circ \delta_{k-1}^r, \beta \circ \delta_{k-1}^r)$ is an element of the boundary $\partial H(\Delta[n] \times \Delta[m]) \subseteq H_i(\Delta[n] \times \Delta[m])$ as required by clause (b) of the statement, or
- $k-2 \geq 0$ and $\beta(k-2) = \beta(k-1)$, in which case we let $l = k-1$ and discharge our remaining obligation in the last paragraph of this proof.

Arguing dually for the face $(\alpha, \beta) \cdot \delta_{k+1}^r$ we see that the inequality $\alpha(k) < \alpha(k+1)$ of display (36) implies that either:

- $\alpha(k+1) \notin \text{im}(\alpha \circ \delta_{k+1}^r)$ and thus that $(\alpha, \beta) \cdot \delta_{k+1}^r = (\alpha \circ \delta_{k+1}^r, \beta \circ \delta_{k+1}^r)$ is an element of the boundary $\partial H(\Delta[n] \times \Delta[m]) \subseteq H_i(\Delta[n] \times \Delta[m])$ as required by clause (b) of the statement, or
- $k+2 \leq r$ and $\alpha(k+1) = \alpha(k+2)$, in which case we let $l = k+1$ and discharge our remaining obligation in the last paragraph of this proof.

So in the cases which remain outstanding we have an $l \in [r]$ (with $0 < l < r$) at which the (in)equalities

$$\alpha(l-1) < \alpha(l) = \alpha(l+1) \text{ and } \beta(l-1) = \beta(l) < \beta(l+1) \quad (37)$$

hold and in order to establish clause (b) of the statement we need to prove that the face $(\alpha, \beta) \cdot \delta_l^r$ is an element of $H_i(\Delta[n] \times \Delta[m])$. To that end let v be the largest integer for which $(\alpha, \beta) \cdot \delta_l^r$ is a face of the shuffle (α_v, β_v) . Notice that we've already shown that (α, β) is a face of the shuffle (α_w, β_w) so it follows that $(\alpha, \beta) \cdot \delta_l^r$ is also a face of that shuffle and therefore we may apply the maximality of v to show that $v \geq w$. Indeed we may show that γ_v and γ_w are actually distinct operators and therefore that the inequality of the last sentence is actually strict. To do this start by arguing just as in the first paragraph of this proof to show that $\gamma_v(\alpha(l-1)) = \beta(l+1)$. To compare this with the value $\gamma_w(\alpha(l-1))$ we start by observing that $\gamma_w(\alpha(l-1)) \leq \gamma_w(\alpha(l) - 1)$, since the first inequality of display (37) implies that $\alpha(l-1) \leq \alpha(l) - 1$ and γ_w is order preserving, then we use the fact that (α, β) is a face of (α_w, β_w) to establish that the inequality $\gamma_w(\alpha(l) - 1) \leq \beta(l)$ also holds as in display (4) of lemma 45(5). Combining these various (in)equalities with the second inequality of display (37) we see that

$$\gamma_w(\alpha(l-1)) \leq \gamma_w(\alpha(l) - 1) \leq \beta(l) < \beta(l+1) = \gamma_v(\alpha(l-1))$$

and in particular that γ_w and γ_v are distinct as postulated. So we've succeeded in showing that $v > w$ and we already know that $w > i-1$ so it clearly follows that $v > i$ and therefore that (α_v, β_v) is an element of $H_i(\Delta[n] \times \Delta[m])$ (by definition), as is its face $(\alpha, \beta) \cdot \delta_l^r$ as required. $\square$

Lemma 168 (Major filling lemma). *For fixed $n, m \in \mathbb{N}$ and each $i > 1$ the inclusion map*

$$H_i(\Delta[n] \otimes \Delta[m]) \xhookrightarrow{\subseteq_r} H_{i-1}(\Delta[n] \otimes \Delta[m]) \quad (38)$$

is an f -extension. It follows that for each $N, M \in \underline{\text{Simplex}}$ the horizontal inclusions in the following commutative square

$$\begin{array}{ccc} \partial H(N \boxtimes M) & \xhookrightarrow{\subseteq_r} & H(N \boxtimes M) \\ \subseteq_e \downarrow & & \downarrow \subseteq_e \\ \partial H(N \otimes M) & \xhookrightarrow{\subseteq_r} & H(N \otimes M) \end{array} \quad (39)$$

are both f -extensions.

Proof. Assuming, for the next few paragraphs, that the first part of the statement holds, it follows that the inclusion

$$\partial H(\Delta[n] \otimes \Delta[m]) \xhookrightarrow{\subseteq_r} H(\Delta[n] \otimes \Delta[m]) \quad (40)$$

is also an f -extension since we know, from notation 166, that we may it may be expressed as the composite of the sequence consisting of the inclusions of display (38).

Notice also that in the proof of lemma 139 each simplex used to witness an extension of thinness is a mediator, so it follows by lemma 167(a) that this witness is in $H_i(N \otimes M)$ whenever the face to which it extends thinness is in there. Consequently we may easily adapt that proof, by intersecting each of the subsets $A^{(j)} \cap N \otimes M$ constructed there with $H_i(N \otimes M)$, to demonstrate that each inclusion

$$H_i(\Delta[n] \boxtimes \Delta[m]) \xhookrightarrow{\subseteq_r} H_i(\Delta[n] \otimes \Delta[m])$$


is a t -extension (and thus an f -extension). Taking the special cases $i = 1, \#(n, m)$ we see that both of the vertical maps in display (39) are f -extensions and consequently, applying the composition and cancellation results of observation 52(5), it follows that the upper horizontal of that square is an f -extension iff its lower horizontal is. In particular, it follows that we may infer that the upper horizontal map in the square of display (41) below is an f -extension from the fact that the map in display (40) is an f -extension.

Furthermore, by the definition of $\boxtimes$, it is the case that any simplex which is thin in $N \boxtimes M$ and is not thin in $\Delta[n] \boxtimes \Delta[m]$ must be a crushed cylinder in $N \boxtimes M$. However the only one of these which is not in $\partial(N \boxtimes M)$ is $(\pi_1^{n,m}, \pi_2^{n,m})$ and this is not itself in $H(N \boxtimes M)$, so it follows that $H(N \boxtimes M) = \partial H(N \boxtimes M) \cup H(\Delta[n] \boxtimes \Delta[m])$. Of course, we also know that $\partial H(\Delta[n] \boxtimes \Delta[m]) = \partial H(N \boxtimes M) \cap H(\Delta[n] \boxtimes \Delta[m])$ so we get a pasting square

$$\begin{array}{ccc} \partial H(\Delta[n] \boxtimes \Delta[m]) & \xhookrightarrow{\subseteq_r} & H(\Delta[n] \boxtimes \Delta[m]) \\ \subseteq_e \downarrow & & \downarrow \subseteq_e \\ \partial H(N \boxtimes M) & \xhookrightarrow{\subseteq_r} & H(M \boxtimes M) \end{array} \quad (41)$$

as in observation 158(b). Using that result we may infer that the lower horizontal of this square is an f -extension, as postulated, from the fact we have already demonstrated that its upper horizontal is an f -extension.

It remains to prove that the inclusion in display (38) an f -extension. To that end, start by observing that the shuffle $(\alpha_i, \beta_i) \in H_{i-1}(\Delta[n] \otimes \Delta[m])$ gives rise to a stratified inclusion

FIGURE 5. The shuffle (α_i, β_i) in $\Delta[n] \otimes \Delta[m]$.

$\ulcorner(\alpha_i, \beta_i)\urcorner: \Delta[n+m] \hookrightarrow H_{i-1}(\Delta[n] \otimes \Delta[m])$, by Yoneda's lemma, which we may extend to a regular map on its entire coimage $\Delta[n+m] \subseteq_e P \stackrel{\text{def}}{=} \text{coim}_e(\ulcorner(\alpha_i, \beta_i)\urcorner)$ as in notation 102. This provides us with a stratified isomorphism between P and the regular stratified subset $\{(\alpha_i, \beta_i)\}^* \subseteq_r H_{i-1}(\Delta[n] \otimes \Delta[m])$ generated by the shuffle (α_i, β_i) . Furthermore, by observation 158(a) we know that if we define $X \subseteq_r P$ to be the inverse image of the regular subset $H_i(\Delta[n] \otimes \Delta[m]) \subseteq_r H_{i-1}(\Delta[n] \otimes \Delta[m])$ under $\ulcorner(\alpha_i, \beta_i)\urcorner$ then we get a pasting square

$$\begin{array}{ccc}
 X & \xhookrightarrow{\subseteq_r} & P \\
 \ulcorner(\alpha_i, \beta_i)\urcorner \downarrow & & \downarrow \ulcorner(\alpha_i, \beta_i)\urcorner \\
 H_i(\Delta[n] \otimes \Delta[m]) & \xhookrightarrow[\subseteq_r]{} & H_{i-1}(\Delta[n] \otimes \Delta[m])
 \end{array} \tag{42}$$

in Strat. More explicitly, an r -simplex (operator) $\mu: [r] \longrightarrow [n+m]$ is thin in P iff $(\alpha_i, \beta_i) \cdot \mu$ is thin in $\Delta[n] \otimes \Delta[m]$ and it is in the regular subset $X \subseteq_r P$ iff $(\alpha_i, \beta_i) \cdot \mu$ is in $H_i(\Delta[n] \otimes \Delta[m])$.

It is our intention to apply corollary 162 to prove that the upper horizontal inclusion in display (42) is an f-extension, but to do so we must first select a suitable integer $0 < k < n+m$. By the definition of the linear order $\triangleleft$, it is clear that the $\triangleleft$ -minimal operator $\gamma_1: [n-1] \longrightarrow [m]$ identically 0 (that is $\gamma_1(j) = 0$ for all $j \in [n-1]$). Since $i > 1$, it follows that γ_i is *not* identically 0, in particular there is some $s \in [n-1]$ such that $\gamma_i(j) = 0$ for $j < s$ and $\gamma_i(s) = t$ for some $t > 0$. So letting $k = s + t$ and applying lemma 45 we see that the corresponding shuffle (α_i, β_i) has

$$\alpha_i(j) = \begin{cases} j & \text{for } j < s, \\ s & \text{for } s \leq j \leq k, \\ s+1 & \text{for } j = k+1, \\ \dots & \end{cases} \quad \beta_i(j) = \begin{cases} 0 & \text{for } j < s, \\ j-s & \text{for } s \leq j \leq k, \\ t & \text{for } j = k+1, \\ \dots & \end{cases} \tag{43}$$

(see figure 5). Now we may check each of the conditions of corollary 162 in turn:

corollary 162(a) Consulting display (43) it is clear that (α_i, β_i) satisfies the meditation condition at k , so applying lemma 137(1) we know that it is k -admissible in $\Delta[n] \otimes \Delta[m]$. So if $\mu: [r] \longrightarrow [n+m]$ is k -divided then the face $(\alpha_i, \beta_i) \cdot \mu$ is thin in $\Delta[n] \otimes \Delta[m]$ and it follows that μ itself is thin in P and consequently $\Delta_k^a[n+m] \subseteq_e P$ as required.

lemma 161(a) The shuffle (α_i, β_i) is a simplex of $H_{i-1}(\Delta[n] \otimes \Delta[m])$ and satisfies the mediation condition at k , so we may apply lemma 167(b) to show that the faces $(\alpha_i, \beta_i) \cdot \delta_{k-1}^{n+m}$ and $(\alpha_i, \beta_i) \cdot \delta_{k+1}^{n+m}$ are both elements of $H_i(\Delta[n] \otimes \Delta[m])$ and thus that the operators δ_{k-1}^{n+m} and δ_{k+1}^{n+m} are in the regular subset $X \subseteq_r P$ as required.

lemma 161(b) Suppose that $\mu: [r] \longrightarrow [n+m]$ is a k -divided r -simplex in P and l is the unique integer with $\mu(l) = k$ then the face $(\alpha_i, \beta_i) \cdot \mu$ is a mediator simplex as witnessed by l so we may apply lemma 167(a) to show that $(\alpha_i, \beta_i) \cdot \mu$ is in $H_i(\Delta[n] \otimes \Delta[m])$ whenever $((\alpha_i, \beta_i) \cdot \mu) \cdot \delta_l^r = (\alpha_i, \beta_i) \cdot (\mu \circ \delta_l^r)$ is in there and thus we see that μ is in the subset $X \subseteq_r P$ whenever $\mu \circ \delta_l^r$ is in there as required.

corollary 161(b) If μ and l are as in the last paragraph and $\mu \circ \delta_l^r$ is thin in P then the face $((\alpha_i, \beta_i) \cdot \mu) \cdot \delta_l^r = (\alpha_i, \beta_i) \cdot (\mu \circ \delta_l^r)$ is thin in $\Delta[n] \otimes \Delta[m]$ and we may apply lemma 137(2) to the mediator simplex $(\alpha_i, \beta_i) \cdot \mu$ to show that $((\alpha_i, \beta_i) \cdot \mu) \cdot \delta_{l-1}^r = (\alpha_i, \beta_i) \cdot (\mu \circ \delta_{l-1}^r)$ and $((\alpha_i, \beta_i) \cdot \mu) \cdot \delta_{l+1}^r = (\alpha_i, \beta_i) \cdot (\mu \circ \delta_{l+1}^r)$ are both thin in there and thus that the simplices $\mu \circ \delta_{l-1}^r$ and $\mu \circ \delta_{l+1}^r$ are both thin in P as required.

So applying corollary 162 as foreshadowed, we see that the upper horizontal inclusion in display (42) is an f-extension and consequently, by applying observation 158(a), we find that the map of display (38), which appears as the lower horizontal in our pasting square (42), is an f-extension as required. $\square$

Lemma 169. *For each stratified set Y the horizontal arrows in the commutative square*

$$\begin{array}{ccc} \Lambda_k^a[n] \boxtimes Y & \xrightarrow{\subseteq_r \boxtimes Y} & \Delta_k^a[n] \boxtimes Y \\ \subseteq_r \downarrow & & \downarrow \subseteq_r \\ \Lambda_k^a[n] \otimes Y & \xrightarrow{\subseteq_r \otimes Y} & \Delta_k^a[n] \otimes Y \end{array} \quad (44)$$

are both f-extensions. Combining this result with lemma 143 and corollary 164 we see that for each complicial set A the stratified set $A \Leftarrow Y$ is also complicial.

Proof. First, it is worth commenting that the equivalence of the clauses in the statement follows by applying observation 54 parts (1) and (2) to the partial right adjoint of theorem 148. Furthermore, since the vertical maps in display (44) are both t-extensions (and thus f-extensions) by lemma 139 we may apply the composition and cancellation results of observation 52(5) to show that its upper horizontal is a f-extension iff its lower horizontal is.

Recall, from observation 108, that we may represent an arbitrary stratified set Y as a weighted colimit $\text{colim}(Y, \Delta)$ of standard (possibly thin) simplices. We know that the functors $\Lambda_k^a[n] \boxtimes -$ and $\Delta_k^a[n] \boxtimes -$ preserve this colimit, by lemma 142, from which it follows that the inclusion $\subseteq_r \boxtimes Y: \Lambda_k^a[n] \boxtimes Y \hookrightarrow \Delta_k^a[n] \boxtimes Y$ is isomorphic to the induced map $\text{colim}(Y, \subseteq_r \boxtimes \Delta(-)): \text{colim}(Y, \Lambda_k^a[n] \boxtimes \Delta(-)) \hookrightarrow \text{colim}(Y, \Delta_k^a[n] \boxtimes \Delta(-))$. So applying observation 52(7) we see that this latter map is an f-extension if $\subseteq_r \boxtimes \Delta[m]_?$ is an f-extension for each $[m]_? \in t\Delta$.

In other words, it is sufficient for us to establish the result given in the statement of this lemma for each standard simplex $Y = \Delta[m]_?$, which we do by induction on the dimension m of that simplex:

Base case. In this case the stratified set Y in display (44) is $\Delta[0]$ which is the identity for $\boxtimes$, thus there is nothing to prove.

Inductive case. Assume the induction hypothesis that the result given in the statement of the lemma holds for any standard (thin) simplex $Y = \Delta[p]_?$ of dimension $p < m$. In fact, arguing as above we know that this hypothesis immediately implies that the result of the statement holds for any $(m-1)$ -skeletal stratified set Y , since any such may be expressed as a canonical colimit of standard simplices of dimension $< m$ as discussed in observation 110.

In order to prove that the inclusion $\subseteq_r \otimes \Delta[m]_?: \Lambda_k^a[n] \otimes \Delta[m]_? \hookrightarrow \Delta_k^a[n] \otimes \Delta[m]_?$ is also an f-extension, first note that we may apply observation 132 to the regular inclusion $\Lambda_k^a[n] \subseteq_r \Delta_k^a[n]$ to show that $\Lambda_k^a[n] \otimes \Delta[m]_?$ is a regular subset of $\Delta_k^a[n] \otimes \Delta[m]_?$ for which $\subseteq_r \otimes \Delta[m]_?$ is the inclusion map. Furthermore it is clear that a simplex (α, β) is in this regular subset iff $\text{im}(\alpha) \cup \{k\} \neq [n]$. Our approach to showing that this regular inclusion is an f-extension will be to decompose it into a sequence of regular subsets

$$\Lambda_k^a[n] \otimes \Delta[m]_? \subseteq_r \underset{(1)}{U} \subseteq_r \underset{(2)}{V} \subseteq_r \underset{(3)}{W} \subseteq_r \underset{(4)}{\Delta_k^a[n] \otimes \Delta[m]_?} \quad (45)$$

given by

$$\begin{aligned} (\alpha, \beta) \in U &\iff \text{im}(\alpha) \cup \{k\} \neq [n] \text{ or } \text{im}(\beta) \neq [m] \\ (\alpha, \beta) \in V &\iff \text{im}(\alpha) \cup \{k\} \neq [n] \text{ or } \text{im}(\beta) \neq [m] \text{ or} \\ &\quad (k \notin \text{im}(\alpha) \text{ and } (n, 0) \notin \text{im}(\alpha, \beta)) \\ (\alpha, \beta) \in W &\iff \text{im}(\alpha) \cup \{k\} \neq [n] \text{ or } \text{im}(\beta) \neq [m] \text{ or } (n, 0) \notin \text{im}(\alpha, \beta) \end{aligned} \quad (46)$$

and to show that each one of the inclusions in this decomposition is an f-extension, from which it then follows, by observation 52(5), that their composite is also an f-extension as required. To complete our argument, the proofs in the following numbered paragraphs establish that the correspondingly numbered inclusions in display (45) are f-extensions as stated:

- (1) By observation 132, the regular subset $\partial\Delta[m] \subseteq_r \Delta[m]_?$ give rise to a regular subset $\Delta_k^a[n] \otimes \partial\Delta[m]$ of $\Delta_k^a[n] \otimes \Delta[m]_?$ and it is easily demonstrated that

$$\begin{aligned} \Lambda_k^a[n] \otimes \partial\Delta[m] &= (\Lambda_k^a[n] \otimes \Delta[m]_?) \cap (\Delta_k^a[n] \otimes \partial\Delta[m]) \text{ and} \\ U &= (\Lambda_k^a[n] \otimes \Delta[m]_?) \cup (\Delta_k^a[n] \otimes \partial\Delta[m]) \end{aligned}$$

so we get a pasting square

$$\begin{array}{ccc} \Lambda_k^a[n] \otimes \partial\Delta[m] & \xrightarrow{\subseteq_r} & \Delta_k^a[n] \otimes \partial\Delta[m] \\ \subseteq_r \downarrow & & \downarrow \subseteq_r \\ \Lambda_k^a[n] \otimes \Delta[m]_? & \xrightarrow{\subseteq_r} & U \end{array} \quad (47)$$

in Strat. However, $\partial\Delta[m]$ is clearly $(m-1)$ -skeletal, since all of its simplices of dimension $> m-1$ are degenerate and thus, as discussed above, our induction hypothesis implies that the upper horizontal map in this diagram is an f-extension. It follows, by applying observation 158(b) that the lower horizontal inclusion in this diagram is also an f-extension as required.

- (2) The stratified map $\Delta(\delta_k^n): \Delta[n-1] \longrightarrow \Delta_k^a[n]$ is both regular and an inclusion, the former fact following from the observation that elements in the image of $\Delta(\delta_k^n)$ do not have k as a vertex and thus, by the definition of $\Delta_k^a[n]$, are thin if and only if they

are degenerate. It follows, by observation 132, that the same properties hold for the stratified map:

$$\Delta[n-1] \otimes \Delta[m]_? \xrightarrow{\Delta(\delta_k^n) \otimes \Delta[m]_?} \Delta_k^a[n] \otimes \Delta[m]_?$$

Notice that the image of the subset $H(\Delta[n-1] \otimes \Delta[m]_?)$ under this map is precisely the regular subset of $\Delta_k^a[n] \otimes \Delta[m]_?$ of those simplices (α, β) such that $k \notin \text{im}(\alpha)$ and $(n, 0) \notin \text{im}(\alpha, \beta)$. It is therefore clear that the union of that image and the regular subset U is precisely the subset V of display (46). Furthermore, it is also easily seen that the inverse image of $U \subseteq_r V$ under the restricted regular inclusion $\Delta(\delta_k^n) \otimes \Delta[m]_?: H(\Delta[n-1] \otimes \Delta[m]_?) \hookrightarrow V$ is simply the regular subset $\partial H(\Delta[n-1] \otimes \Delta[m]_?) \subseteq_r H(\Delta[n-1] \otimes \Delta[m]_?)$. Consequently we have a pasting square

$$\begin{array}{ccc} \partial H(\Delta[n-1] \otimes \Delta[m]_?) & \xhookrightarrow{\subseteq_r} & H(\Delta[n-1] \otimes \Delta[m]_?) \\ \Delta(\delta_k^n) \otimes \Delta[m]_? \downarrow & & \downarrow \Delta(\delta_k^n) \otimes \Delta[m]_? \\ U & \xhookrightarrow[\subseteq_r]{} & V \end{array}$$

in Strat. Now, by lemma 168 we know that the upper horizontal map in this diagram is an f-extension and, consequently, can apply observation 158(a) to infer that the lower horizontal map is an f-extension as well.

- (3) Consider the regular subsets $H(\Delta_k^a[n] \otimes \Delta[m]_?)$ and V of $\Delta_k^a[n] \otimes \Delta[m]_?$ and observe that it is easily demonstrated, directly from the definitions given in display (46), that we have

$$\begin{aligned} \partial H(\Delta_k^a[n] \otimes \Delta[m]_?) &= V \cap H(\Delta_k^a[n] \otimes \Delta[m]_?) \quad \text{and} \\ W &= V \cup H(\Delta_k^a[n] \otimes \Delta[m]_?) \end{aligned}$$

and thus that we have a pasting square

$$\begin{array}{ccc} \partial H(\Delta_k^a[n] \otimes \Delta[m]_?) & \xhookrightarrow{\subseteq_r} & H(\Delta_k^a[n] \otimes \Delta[m]_?) \\ \subseteq_r \downarrow & & \downarrow \subseteq_r \\ V & \xhookrightarrow[\subseteq_r]{} & W \end{array}$$

in Strat. The upper horizontal map in this diagram is an f-extension by lemma 168, so we may apply observation 158(b) to infer that the lower horizontal inclusion is an f-extension as well.

- (4) From the description given in display (46) it is clear that only two non-degenerate simplices of $\Delta_k^a[n] \otimes \Delta[m]_?$ are **not** in its regular subset W , those being the maximal shuffle $(\pi_1^{n,m}, \pi_2^{n,m})$ and its k^{th} face $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_k^{n+m}$. The shuffle $(\pi_1^{n,m}, \pi_2^{n,m}) \in \Delta_k^a[n] \otimes \Delta[m]_?$ gives rise to a stratified inclusion $i: \Delta[n+m] \hookrightarrow \Delta_k^a[n] \otimes \Delta[m]_?$, by Yoneda's lemma, which we may extend to a regular map on its entire coimage $\Delta[n+m] \subseteq_e P \stackrel{\text{def}}{=} \text{coim}_e(i)$ as in notation 102. This provides us with a stratified isomorphism between P and the regular stratified subset $\{(\pi_1^{n,m}, \pi_2^{n,m})\}^* \subseteq_r \Delta_k^a[n] \otimes \Delta[m]_?$ generated by our shuffle. Furthermore, by observation 158(a) we know that if

we define $X \subseteq_r P$ to be the inverse image of the regular subset $W \subseteq_r \Delta_k^a[n] \otimes \Delta[m]_?$ under i then we get a pasting square

$$\begin{array}{ccc} X & \xrightarrow{\subseteq_r} & P \\ i \downarrow & & \downarrow i \\ W & \xrightarrow[\subseteq_r]{} & \Delta_k^a[n] \otimes \Delta[m]_? \end{array} \quad (48)$$

in Strat. More explicitly, an r -simplex (operator) $\mu: [r] \longrightarrow [n+m]$ is thin in P iff the face $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \mu$ is thin in $\Delta_k^a[n] \otimes \Delta[m]_?$ and it is in the regular subset $X \subseteq_r P$ iff $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \mu$ is in W .

Since $(\pi_1^{n,m}, \pi_2^{n,m})$ and its face $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_k^{n+m}$ are the only simplices of $\Delta_k^a[n] \otimes \Delta[m]$ which are not in W , it follows that the regular subset $X \subseteq_r P$ has the horn $\Lambda_k[n+m]$ as its underlying simplicial set. Furthermore, arguing as in item (iv) of the proof of lemma 143, we know that the shuffle $(\pi_1^{n,m}, \pi_2^{n,m})$ is k -admissible in $\Delta_k^a[n] \otimes \Delta[m]_?$ so if $\mu: [r] \longrightarrow [n+m]$ is a k -divided simplex (operator) then $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \mu$ is thin in $\Delta_k^a[n] \otimes \Delta[m]_?$ and so μ is thin in P and it follows that $\Delta_k^a[n+m] \subseteq_e P$.

These observations immediately serve to establish the conditions of lemma 161 and condition (a) of corollary 162 for $X \subseteq_r P$. So to complete the verification of the remaining condition (b) of that corollary all we need do is demonstrate that if δ_k^{n+m} is thin in P then so are δ_{k-1}^{n+m} and δ_{k+1}^{n+m} , or equivalently that if $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_k^{n+m}$ is thin in $\Delta_k^a[n] \otimes \Delta[m]_?$ then so are $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_{k-1}^{n+m}$ and $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_{k+1}^{n+m}$. However the face $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_k^{n+m}$ is equal to the cylinder $(\delta_k^n \circ \pi_1^{n-1,m}, \pi_2^{n-1,m})$, since $k < n$, and this is thin in $\Delta_k^a[n] \otimes \Delta[m]_?$ iff $\text{id}_{[m]}$ is thin in $\Delta[m]_?$ (since δ_k^n is not thin in $\Delta_k^a[n]$) which must therefore be equal to $\Delta[m]_t$. Of course, $\text{Th}_{m-1}(\Delta[m]) = \Delta[m]_t$ and $\text{Th}_{n-1}(\Delta[n]) \subseteq_e \Delta_k^a[n]$ so we may apply lemma 133 to show that $\text{Th}_{n+m-2}(\Delta[n] \otimes \Delta[m]) \subseteq_e \text{Th}_{n-1}(\Delta[n]) \otimes \text{Th}_{m-1}(\Delta[m]) \subseteq_e \Delta_k^a[n] \otimes \Delta[m]_t$ and thus that all $(n+m-1)$ -dimensional simplices of $\Delta_k^a[n] \otimes \Delta[m]_t$, including the faces $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_{k-1}^{n+m}$ and $(\pi_1^{n,m}, \pi_2^{n,m}) \cdot \delta_{k+1}^{n+m}$, are all thin as required.

Finally applying corollary 162 we may infer that the horizontal inclusion at the top of display (48) is an f-extension so, by observation 158(a), it follows that the horizontal inclusion at the bottom of that pasting square is also an f-extension as required. $\square$

To conclude this section we present an important result which parallels theorem 148, but applies to the category of complicial sets $\underline{\mathbf{Cs}}$ rather than the larger category of pre-complicial sets $\underline{\mathbf{Pcs}}$.

Theorem 170. *For any complicial set A and any stratified sets X and Y , the stratified sets $A \Leftarrow Y$ and $X \Rightarrow A$ are both complicial. It follows that if $f: Z \longrightarrow Z'$ is an f-extension then so are the maps*

$$\begin{array}{ccc} Z \otimes Y & \xrightarrow{f \otimes Y} & Z' \otimes Y \\ X \otimes Z & \xrightarrow{X \otimes f} & X \otimes Z' \end{array}$$

Furthermore, by Day's reflection theorem (theorem 55), we may reflect the monoidal structure on Strat to a monoidal bi-closed structure on $\underline{\mathbf{Cs}}$ with tensor product $A \otimes_c B \stackrel{\text{def}}{=} \text{L}_c(A \otimes B)$

(where L_c is reflector associated with $\underline{\mathbf{Pcs}}$) and left and right closures $A \Rightarrow *$ and $* \Leftarrow B$ respectively.

Proof. Almost identical to that of theorem 148, except that we start by observing that the functor $- \otimes Y : \underline{\mathbf{Strat}} \longrightarrow \underline{\mathbf{Strat}}$ carries the primitive f-extensions $d : \Delta[1]_t \longrightarrow \Delta[0]$ and $\subseteq_r : \Lambda_k^a[n] \hookrightarrow \Delta_k^a[n]$ to stratified maps which are also f-extensions by applying lemmas 164 and 169 respectively. The remainder of the argument presented there carries through unaltered, aside from the substitution of the words “complicial” and “f-extension” for “pre-complicial” and “t-extension”. $\square$

8.4. Superstructures of Complicial Sets. Of course, we are motivated to think of the category of complicial sets $\underline{\mathbf{Cs}}$ as being analogous to the category of ω -categories $\omega\text{-}\underline{\mathbf{Cat}}$. On doing so, the functor $|\cdot|_n : \underline{\mathbf{Strat}} \longrightarrow \underline{\mathbf{Strat}}$ takes on a particular importance since it allows us to identify a subcategory of $\underline{\mathbf{Cs}}$ which corresponds to the full subcategory $n\text{-}\underline{\mathbf{Cat}}$ of n -categories in $\omega\text{-}\underline{\mathbf{Cat}}$. In pursuing this line of inquiry we’ll need the following lemma:

Lemma 171. *If A is a complicial set then so is its n -dimensional superstructure $|A|_n$. Equivalently, if the stratified map $f : X \longrightarrow Y$ is an f-extension then so is the stratified map $\text{Th}_n(f) : \text{Th}_n(X) \longrightarrow \text{Th}_n(Y)$.*

Proof. Extending the argument used to prove lemma 150, all we need do is prove that each map obtained by applying Th_n to a primitive f-extension is again an f-extension. We consider each of the classes of primitive f-extensions in turn:

$d : \Delta[1]_t \longrightarrow \Delta[0]$ All of the simplices in $\Delta[0]$ and $\Delta[1]_t$ of dimension > 0 are already thin, therefore $\text{Th}_n(\Delta[0]) = \Delta[0]$ and $\text{Th}_n(\Delta[1]_t) = \Delta[1]_t$ for any $n \in \mathbb{N}$. It follows that the functor Th_n maps $d : \Delta[1]_t \longrightarrow \Delta[0]$ to itself.

$\Lambda_k^a[m] \hookrightarrow_r \Delta_k^a[m]$ Again, we have two cases:

- **$n \geq m - 1$** In this case, all of the simplices of $\Lambda_k^a[m]$ and $\Delta_k^a[m]$ of dimension $> n$ are already thin, therefore $\text{Th}_n(\Lambda_k^a[m]) = \Lambda_k^a[m]$, $\text{Th}_n(\Delta_k^a[m]) = \Delta_k^a[m]$ and, consequently, Th_n maps $\Lambda_k^a[m] \hookrightarrow_r \Delta_k^a[m]$ to itself.
- **$n < m - 1$** This is a simple matter of applying corollary 162 to the regular subset $\text{Th}_n(\Lambda_k^a[m]) \subseteq_r \text{Th}_n(\Delta_k^a[m])$. To do so observe that the conditions of lemma 161 follow trivially for any stratified subset whose underlying simplicial set is a k -horn and that condition (a) of corollary 162 simply asks for the standard inclusion $\Delta_k^a[m] \subseteq_e \text{Th}_n(\Delta_k^a[m])$. Finally, the only simplex of $\text{Th}_n(\Delta_k^a[m])$ which satisfies the postulates of corollary 162 condition (b) is the m -simplex $\text{id}_{[m]}$ of which all $(m - 1)$ -dimensional faces, including δ_{k-1}^m , δ_k^m and δ_{k+1}^m , are thin as required.

In either case the resulting map is an f-extension as required. $\square$

Definition 172. We define an n -complicial set to be a complicial set A which is n -trivial and let $\underline{\mathbf{Cs}}_n = \underline{\mathbf{Cs}} \cap \underline{\mathbf{Strat}}_n$ denote the full subcategory of $\underline{\mathbf{Cs}}$ whose objects are these n -complicial sets.

Applying lemma 151 and arguing just as in observation 152, we know that $X \Rightarrow A$ and $A \Leftarrow Y$ are n -complicial whenever A is and that, just as in theorem 170, we may reflect the monoidal structure of $\underline{\mathbf{Strat}}$ to a monoidal biclosed structure on the reflective full subcategory $\underline{\mathbf{Cs}}_n$.

Lemma 173. *If A is a complicial set then its n -skeleton $\|A\|_0$ and 0-superstructure $|A|_0$ coincide. Furthermore, any 0-skeletal stratified set is 0-complicial and, consequently, the category $\underline{\mathbf{Cs}}_0$ is actually the full subcategory of 0-skeletal stratified sets in Strat (and is therefore equivalent to Set).*

Proof. Any simplex of dimension > 0 in the 0-skeleton $\|A\|_0$ is degenerate and thus thin in A , so it follows that $\|A\|_0 \subseteq_r |A|_0$. Conversely, if a is a simplex in $|A|_0$ of dimension > 0 then it is pre-degenerate at 0 (see definition 116), since every one of its faces of dimension > 0 must be thin for it to be in $|A|_0$, and it follows, by lemma 163, that a is actually degenerate at 0. In other words, we see (by definition) that $|A|_0$ is 0-skeletal and thus it must be a regular subset of $\|A\|_0$, so the first of our results follows.

From observation 111 we know that all of the simplices and horns which constitute the domains and codomains of the primitive f-extensions are connected and therefore that each of these is bijective on components. However, by the same observation, a stratified set is 0-skeletal iff it is orthogonal to every stratified map which is bijective on components and it follows that every 0-skeletal stratified set is orthogonal to every primitive f-extension. In other words, every 0-skeletal stratified set is complicial furthermore, by definition, any simplex of a 0-skeletal set of dimension > 0 is degenerate and thus thin, thereby demonstrating that any such stratified set is actually 0-complicial. Therefore, we may conclude that $\underline{\mathbf{Cs}}_0$ coincides with the full subcategory of 0-skeletal sets in Strat and thus (from observation 111) that the functor $\text{dis}: \underline{\mathbf{Set}} \longrightarrow \underline{\mathbf{Strat}}$ provides us with the required equivalence between Set and $\underline{\mathbf{Cs}}_0$. $\square$

9. THE PATH CATEGORY CONSTRUCTION

9.1. The Complicial Category of Prisms.

Notation 174 (complicial categories). We will often use the terms *complicial category* and *complicial functor* to refer to the objects and arrows of $\underline{\text{Cat}}(\underline{\text{Cs}})$ (respectively). An n -arrow (resp. n -object) of a complicial category $\mathbb{C}$ is simply defined to be an n -simplex in its complicial set of arrows C (resp. objects $\mathbb{C}_o$).

When manipulating complicial categories we will, in general, assume that all standard categorical notions and constructions have been internalised to $\underline{\text{Cs}}$ in the canonical way. When referring to these we may not stress the use of the “complicial” qualifier if it is clear that it may be inferred from the context.

We will also tend to freely apply qualifiers from our theory of stratified and complicial sets to complicial categories where the meaning is clear. For instance, if we say that $\mathbb{D}$ is a regular subcategory of the complicial category $\mathbb{C}$, it should be clear that we mean that the former is a complicial subcategory of the latter for which $\mathbb{D}_a$ is, in fact, a regular subset of $\mathbb{C}_a$.

Lemma 175 (the category of prisms in a complicial set). *If $\mathbb{T}_{\underline{\text{Cat}}}$ is the LE-theory of categories discussed in observation 70 then the functor $\Delta_{\mathbb{P}} = \text{Th}_1 \circ \Delta: \Delta \longrightarrow \underline{\text{Strat}}$ is a finitely presented f -almost $\mathbb{T}_{\underline{\text{Cat}}}$ -coalgebra. Consequently we may apply the internal version of Kan’s construction, observation 64, to build a right adjoint, finitely accessible functor*

$$\underline{\text{Cs}} \xrightarrow{\mathbb{P} \stackrel{\text{def}}{=} K_{\Delta_{\mathbb{P}}}^l} \underline{\text{Cat}}(\underline{\text{Cs}}) = \mathbb{T}_{\underline{\text{Cat}}}\text{-Alg}(\underline{\text{Cs}})$$

which carries a complicial set A to a complicial category $\mathbb{P}(A)$ which we call its **category of prisms**.

Proof. Following the argument given in observation 66, we may apply Yoneda’s lemma to show that the stratified map $\text{colim}(\subseteq_s, \Delta): \text{colim}(\Lambda_k[n], \Delta) \longrightarrow \text{colim}(\Delta[n], \Delta)$ is actually isomorphic to the inclusion $\Lambda_k[n] \xrightarrow{\subseteq_r} \Delta[n]$ under the usual identification of $\underline{\text{Simp}}$ as the category of minimally stratified sets in $\underline{\text{Strat}}$. Furthermore, we know that Th_1 is a left adjoint functor and thus that it preserves colimits, so it follows we have a family of isomorphisms $\text{colim}(X, \text{Th}_1 \circ \Delta) \cong \text{Th}_1(\text{colim}(X, \Delta)) \cong \text{Th}_1(X)$ which is natural in $X \in \underline{\text{Simp}}$, and thus that $\text{colim}(\subseteq_s, \text{Th}_1 \circ \Delta): \text{colim}(\Lambda_k[n], \text{Th}_1 \circ \Delta) \longrightarrow \text{colim}(\Delta[n], \text{Th}_1 \circ \Delta)$ is isomorphic to the inclusion $\text{Th}_1(\Lambda_k[n]) \xrightarrow{\subseteq_r} \text{Th}_1(\Delta[n])$. Applying this observation to the colimit characterisation of L -almost coalgebras from observation 63, it follows that $\text{Th}_1 \circ \Delta$ is an f -almost $\mathbb{T}_{\underline{\text{Cat}}}$ -coalgebra if and only if each of the latter inclusions is an f -extension. However, we know that $\text{Th}_1(\Delta[n]) = \text{Th}_1(\Delta_k^a[n])$ and $\text{Th}_1(\Lambda_k[n]) = \text{Th}_1(\Lambda_k^a[n])$ so these inclusions may equally well be obtained by applying Th_1 to the primitive f -extensions of definition 153 and thus, applying lemma 171, we see that they too are f -extensions as required. $\square$

Observation 176 (handedness conventions). Notice that we could equally well have chosen to define $\mathbb{P}$ to be the functor $K_{\Delta_{\mathbb{P}}}^r: \underline{\text{Cs}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ obtained using the right handed, rather than left handed, version of Kan’s construction (as discussed in observation 64). However, we have chosen to use the left handed construction in this section because it is compatible with subsequent arguments involving the decalage construction on $\underline{\text{Cs}}$.

Where necessary we might differentiate these two possibilities using superscripts $\mathbb{P}^l \stackrel{\text{def}}{=} K_{\Delta_{\mathbb{P}}}^l$ and $\mathbb{P}^r \stackrel{\text{def}}{=} K_{\Delta_{\mathbb{P}}}^r$ and adopt this superscript convention to indicate the handedness convention for the other category and double category constructions introduced in this section. However,

for the remainder of this section and the next we elide these superscripts and assume that all constructions will be made using the left closure. In the final section we shall be a little more careful, since there it turns out that we will need to deploy the right handed dual.

Of course every result proved here for these left handed constructions also applies, dually, for the right handed version. Indeed, the dual construction on stratified simplicial sets allows us to directly relate these constructions. For instance we may show that if A is a complicial set then $\mathbb{P}^r(A^\circ)$ is actually canonically isomorphic to the complicial category obtained by applying the duality functor $(-)^{\circ}: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Cs}}$ point-wise to $\mathbb{P}^l(A)$ and then applying the usual categorical dual.

Observation 177 (prism categories under the microscope). Observation 66 demonstrates that the prism category functor bears a more explicit description as the external Kan functor $K_{\Delta_{\mathbb{P}} \otimes \Delta}$ from $\underline{\mathbf{Cs}}$ to $\mathbb{T}_{\mathbf{Cat}} \otimes \mathbb{T}_{\underline{\mathbf{Cs}}}\text{-}\underline{\mathbf{Alg}} \cong \mathbf{Cat}(\underline{\mathbf{Cs}})$. Combining this with the explicit description of the relationship between $\mathbb{T}_{\mathbf{Cat}}$ -algebras and categories given in observation 70 we gain the following explicit information about the structure of $\mathbb{P}(A)$ for a complicial set A :

- (1) The complicial set of arrows of $\mathbb{P}(A)$ is $K_{\Delta_{\mathbb{P}}}^l(A)_1 = \Delta[1] \Rightarrow A$, alternatively using the explicit presentation $K_{\Delta_{\mathbb{P}} \otimes \Delta}(A)$ we see that the n -arrows of $\mathbb{P}(A)$ may be identified with stratified maps of the form:

$$\Delta[1] \otimes \Delta[n] \xrightarrow{p} A$$

Furthermore the action of a simplicial operator $\alpha: [m] \longrightarrow [n]$ on an n -arrow expressed in this form is given by $p \cdot \alpha = p \circ (\Delta[1] \otimes \Delta(\alpha))$.

- (2) Such an n -arrow in $\mathbb{P}(A)$ is thin if and only if it extends to a stratified map:

$$\Delta[1] \otimes \Delta[n]_t \xrightarrow{p} A$$

but we know, by observation 141, that to demonstrate this all we need to do is show that p maps the mediator simplices and crushed cylinders of $\Delta[1] \otimes \Delta[n]_t$ to thin simplices in A . However, we already know that this is true for all mediator simplices, since these are thin in $\Delta[1] \otimes \Delta[n]$, and it is an easy matter to demonstrate that the only non-degenerate crushed cylinders of $\Delta[1] \otimes \Delta[n]_t$ are the two n -simplices $(\varepsilon_0^1 \circ \eta^n, \text{id}_{[n]})$ and $(\varepsilon_1^1 \circ \eta^n, \text{id}_{[n]})$ and the $(n+1)$ -simplex $(\pi_1^{(1,n)}, \pi_2^{(1,n)})$. It follows that p is a thin n -arrow iff it maps these three simplices to thin simplices in A .

- (3) The complicial set of objects of $\mathbb{P}(A)$ is (isomorphic to) A itself and the following diagram displays its various source, target and identity maps:

$$\begin{array}{ccc} & \xrightarrow{s \stackrel{\text{def}}{=} \Delta(\varepsilon_0^1) \Rightarrow A} & \\ A & \xleftarrow{i \stackrel{\text{def}}{=} \Delta(\eta^1) \Rightarrow A} \Delta[1] \Rightarrow A & \\ & \xrightarrow{t \stackrel{\text{def}}{=} \Delta(\varepsilon_1^1) \Rightarrow A} & \end{array}$$

More explicitly, if $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ is an n -arrow of $\mathbb{P}(A)$, presented as in (1), then we have $s(p) \stackrel{\text{def}}{=} p(\varepsilon_0^1 \circ \eta^n, \text{id}_{[n]})$ and $t(p) \stackrel{\text{def}}{=} p(\varepsilon_1^1 \circ \eta^n, \text{id}_{[n]})$ in A . In the other direction, if $a \in A_n$ then the identity n -arrow $i(a): \Delta[1] \otimes \Delta[n] \longrightarrow A$ maps an r -simplex $(\alpha, \beta) \in \Delta[1] \otimes \Delta[n]$ to $i(a)(\alpha, \beta) \stackrel{\text{def}}{=} a \cdot \beta$,

In order to explicitly describe the corresponding complicial subset of identities within $\Delta[1] \Rightarrow A$, notice that the stratified map $\Delta(\eta^1): \Delta[1] \longrightarrow \Delta[0]$ extends to one

with domain $\Delta[1]_t$ and it is this which we called d in definition 153(c). Consequently, if we think of $\Delta[1]_t \Rightarrow A$ as a regular subset of $\Delta[1] \Rightarrow A$ (as in lemma 147) then the stratified map $\mathbf{i}$ from above factors as:

$$A \xrightarrow{d \Rightarrow A} \Delta[1]_t \Rightarrow A \xrightarrow{\subseteq_r} \Delta[1] \Rightarrow A$$

However, d is a primitive f-extension and so, by theorem 55(2), it follows that $d \Rightarrow A$ is a stratified isomorphism. In other words, the image of $\mathbf{i}$, which we know to be the set of identities in $\mathbb{P}(A)$, is precisely the regular subset $\Delta[1]_t \Rightarrow A \subseteq_r \Delta[1] \Rightarrow A$ and so an n -arrow $p \in \mathbb{P}(A)$ is an identity if and only if it extends to a stratified map:

$$\Delta[1]_t \otimes \Delta[n] \xrightarrow{p} A$$

- (4) A composable pair of n -arrows $p, q: \Delta[1] \otimes \Delta[n] \longrightarrow A$ corresponds to a unique stratified map $\Lambda_1[2] \otimes \Delta[n] \longrightarrow A$ that we may uniquely extend along the f-extension $\Lambda_1[2] \otimes \Delta[n] \xrightarrow{\subseteq_r} \Delta[2]_t \otimes \Delta[n]$ to a stratified map $w(p, q): \Delta[2]_t \otimes \Delta[n] \longrightarrow A$ and which is thus completely determined by the fact that $w(p, q) \circ (\Delta(\delta_2^2) \otimes \Delta[n]) = p$ and $w(p, q) \circ (\Delta(\delta_0^2) \otimes \Delta[n]) = q$. Now the composite of these n -arrows is given by $q * p \stackrel{\text{def}}{=} w(p, q) \circ (\Delta(\delta_1^2) \otimes \Delta[n])$ and we say that this composite is *witnessed* by the stratified map $w(p, q)$.

Observation 178 (complicially enriched categories). A *complicially enriched* category in $\underline{\mathbf{Cs}}$ is a complicial category $\mathbb{C}$ whose set of objects $\mathbb{C}_o$ is a 0-complicial set.

Applying lemma 173, this is equivalent to saying that $\mathbb{C}_o$ is 0-skeletal and thus isomorphic to the discrete stratified set on its set of 0-simplices. Consequently, it is easily seen that a complicially enriched category gives rise to an enriched category, in the sense of Kelly's book [19], whose set of objects is the set of 0-simplices of $\mathbb{C}_o$. Conversely, if $\mathcal{D}$ is a category (with a small set of objects) which is enriched in the cartesian category $(\underline{\mathbf{Cs}}, \times, \Delta[0])$ then the disjoint union $\bigsqcup_{d, d' \in \text{obj}(\mathcal{D})} \mathcal{D}(d, d')$ in $\underline{\mathbf{Cs}}$ inherits an internal category structure from $\mathcal{D}$ which makes it into the complicial set of arrows of a complicially enriched category in our sense. Indeed, these constructions demonstrate that the (2-)categories of complicially enriched categories in our sense, on the one hand, and categories enriched in the monoidal category $(\underline{\mathbf{Cs}}, \times, \Delta[0])$ in the sense of Kelly [19], on the other, are equivalent.

Notice that the inclusion of $\underline{\mathbf{Cs}}\text{-}\underline{\mathbf{Cat}}$ in $\underline{\mathbf{Cat}}(\underline{\mathbf{Cs}})$ has a right adjoint

$$\begin{array}{ccc} & \text{inclusion} & \\ \underline{\mathbf{Cs}}\text{-}\underline{\mathbf{Cat}} & \xrightarrow{\quad} & \underline{\mathbf{Cat}}(\underline{\mathbf{Cs}}) \\ & \text{\scriptsize $\perp$} & \\ & \xleftarrow{\quad \text{E} \quad} & \end{array}$$

where the complicially enriched category $\text{E}(\mathbb{C})$ can be constructed as the complicial full subcategory of $\mathbb{C}$ whose complicial set of objects is $|\mathbb{C}_o|_0 \subseteq_r \mathbb{C}_o$. In other words, $\text{E}(\mathbb{C})$ is the subcategory of $\mathbb{C}$ whose complicial set of arrows is the regular subset of $\mathbb{C}_a$ obtained by pulling the regular subset $|\mathbb{C}_o|_0 \times |\mathbb{C}_o|_0 \subseteq_r \mathbb{C}_o \times \mathbb{C}_o$ back along the stratified map $(\mathbf{s}, \mathbf{t}): \mathbb{C}_a \longrightarrow \mathbb{C}_o \times \mathbb{C}_o$. Furthermore, it is clear that if $f: \mathbb{C} \longrightarrow \mathbb{C}'$ is a complicial functor then $\text{E}(f)$ is constructed by restricting f to a complicial functor $f: \text{E}(\mathbb{C}) \longrightarrow \text{E}(\mathbb{C}')$.

Definition 179 (the category of paths in a complicial set). Define $\text{P}: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Cs}}\text{-}\underline{\mathbf{Cat}}$ to be the composite of the prism category functor $\mathbb{P}: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Cat}}(\underline{\mathbf{Cs}})$ and the coreflection functor

$E: \underline{\text{Cat}}(\underline{\text{Cs}}) \longrightarrow \underline{\text{Cs-Cat}}$. In other words, $P(A)$ is the complicial full subcategory of $\mathbb{P}(A)$ whose set of objects is the superstructure $|A|_0 \subseteq_r A$.

Observation 180 (an explicit description of the n -arrows of $P(A)$). By definition an n -arrow p of $\mathbb{P}(A)$ is in the complicial subcategory $P(A)$ if and only if $s(p)$ and $t(p)$ are both simplices in $|A|_0$. However we know, by observation 177(3), that for an explicitly presented n -arrow $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ we have $s(p) = p(\varepsilon_0^1 \circ \eta^n, \text{id}_{[n]})$ and $t(p) = p(\varepsilon_1^1 \circ \eta^n, \text{id}_{[n]})$, so it follows that p is in $P(A)$ iff all faces of these n -simplices with dimension > 0 are thin in A .

Observation 181 (discrete path categories). A trivial observation, which we shall use a couple of times in forthcoming arguments, is that if A is 0-complicial then the path category $\mathbb{P}(A)$ is complicially enriched (and consequently $\mathbb{P}(A) = P(A)$) and discrete as a category (cf. definition 74).

The first of these observations is an immediate consequence of the fact that the complicial set of objects of $\mathbb{P}(A)$ is A . The latter we prove by recalling that the primitive f-extension $d: \Delta[1]_t \longrightarrow \Delta[0]$ was constructed by extending the stratified map $\Delta(\eta^1): \Delta[1] \longrightarrow \Delta[0]$ through the inclusion $\Delta[1] \xrightarrow{\subseteq_c} \Delta[1]_t$. In other words, $\Delta(\eta^1)$ is the composite of an f-extension and an entire inclusion which is orthogonal to every 0-trivial stratified set, so it is itself orthogonal to every 0-complicial set. However, since A is 0-complicial it follows, by applying the fact that $\underline{\text{Cs}}_0$ is closed in $\underline{\text{Pcs}}$ under right closure (cf. definition 172), that the identities map $i = A \leftarrow \Delta(\eta^1)$ of $\mathbb{P}(A)$ is an isomorphism and so this category is discrete as required.

9.2. Path Categories and Superstructures.

Observation 182 (superstructures of complicially enriched categories). Observe that (for $m \in \mathbb{N}$) the complicial superstructure functor $|\cdot|_m$ of definition 112 preserves finite limits, since it is right adjoint to Th_m , so we know that it may be lifted to an endo-functor on $\underline{\text{Cat}}(\underline{\text{Cs}})$ as discussed in definition 60. Concretely, if $\mathbb{C}$ is a complicial category then its structural maps i , s and t restrict to maps between $|\mathbb{C}_o|_m \subseteq_r \mathbb{C}_o$ and $|\mathbb{C}_a|_m \subseteq_r \mathbb{C}_a$ additionally $|\mathbb{C}_a|_m$ is closed in $\mathbb{C}_a$ under the action of $*$. In other words, $|\mathbb{C}_a|_m$ is the underlying complicial set of arrows of a (regular) complicial subcategory of $\mathbb{C}$. Furthermore, the action of this lifted functor on complicial functors is simply one of restricting them to these regular sub-categories.

Of course, complicial sub-categories of complicially enriched categories are also complicially enriched and so our lifted superstructure functor restricts to an endo-functor on $\underline{\text{Cs-Cat}}$.

At the risk of confusing our notation a little, we actually call the subcategory obtained in this way the $(m+1)$ -dimensional superstructure of the complicially enriched category $\mathbb{C}$ and denote it using the notation $|\mathbb{C}|_{m+1}$ rather than following the conventions established for lifted functors in definition 60. While this “dimension shift” might cause a little consternation at first it is important since, as we shall see in a page or two, this subcategory really is the appropriate analogue of the $(m+1)$ -dimensional superstructure construction as generalised to the realm of complicially enriched categories.

Finally, it is convenient to round off this sequence of superstructure functors by defining another endo-functor $|\cdot|_0$ on $\underline{\text{Cs-Cat}}$, letting $|\mathbb{C}|_0$ be the discrete subcategory of all identities in $\mathbb{C}$. In other words, $|\mathbb{C}|_0$ is the smallest subcategory of $\mathbb{C}$ which contains all of its objects.

Taking successive superstructures of a complicially enriched category $\mathbb{C}$ it is immediate that we obtain a canonical filtered family $|\mathbb{C}|_0 \subseteq_r |\mathbb{C}|_1 \subseteq_r \dots \subseteq_r |\mathbb{C}|_n \subseteq_r \dots$ of regular subcategories the union of which is $\mathbb{C}$ itself.

Observation 183. If A is a complicial set then the path category $P(|A|_m)$ is also (trivially) a complicial subcategory of $P(A)$. Of course, an n -arrow $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ of $P(A)$ is in $P(|A|_m)$ iff it factors through the inclusion $|A|_m \subseteq_r A$ and, in turn, we may apply the adjunction $\text{Th}_m \dashv |\cdot|_m$ to show that this happens precisely when p extends to a stratified map:

$$\text{Th}_m(\Delta[1] \otimes \Delta[n]) \xrightarrow{p} A$$

It is instructive to compare this to the closely related complicial subcategory $|P(A)|_m$ for $m > 0$. By Yoneda's lemma, and the definition of superstructures of complicially enriched categories, an n -arrow in here corresponds to a stratified map

$$\Delta[n] \longrightarrow |P(A)_a|_{m-1} \subseteq_r |\Delta[1] \Rightarrow A|_{m-1}$$

which we may transpose under the adjunction $\text{Th}_m \dashv |\cdot|_m$ and then under the partial adjunction $\Delta[1] \otimes - \dashv \Delta[1] \Rightarrow *$ to show that it yields a stratified map:

$$\Delta[1] \otimes \text{Th}_{m-1}(\Delta[n]) \xrightarrow{p} A$$

In other words, our n -arrow $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ in $P(A)$ is an arrow in the subcategory $|P(A)|_m$ iff it extends to a stratified map with domain $\Delta[1] \otimes \text{Th}_{m-1}(\Delta[n])$.

Given these representations of the arrows of these two subcategories, the following lemma is now easily established:

Lemma 184. *If A is a complicial set then the complicial sub-categories $P(|A|_m)$ and $|P(A)|_m$ of $P(A)$ are in fact identical for each $m \geq 0$.*

Proof. First let us dispose with the special case $m = 0$, for which the definition of $|\cdot|_0$ given in observation 182 was selected. By observation 181 we know that $P(|A|_0)$ is a discrete subcategory of $P(A)$ whose complicial set of objects is $|A|_0 \subseteq_r A$. However, by definition we also know that $|P(A)|_0$ is the discrete subcategory of $P(A)$ with the same complicial set of objects and thus these two sub-categories are identical.

So from now on we assume that $m > 0$ and prove the stated equality as two inclusions:

$|P(A)|_m \subseteq_r P(|A|_m)$: We know that any r -simplices of $\Delta[1]$ with $r > 1$ are degenerate, and thus thin, so it follows that $\Delta[1] = \text{Th}_1(\Delta[1])$ and applying lemma 133 we see that:

$$\text{Th}_m(\Delta[1] \otimes \Delta[n]) \subseteq_e \text{Th}_1(\Delta[1]) \otimes \text{Th}_{m-1}(\Delta[n]) = \Delta[1] \otimes \text{Th}_{m-1}(\Delta[n])$$

Thus, from the explicit descriptions furnished by observation 183, we see that if an n -arrow p of $P(A)$ is an element of $|P(A)|_m$ then it extends to a stratified map with domain $\Delta[1] \otimes \text{Th}_{m-1}(\Delta[n])$ which may then be restricted to one with domain $\text{Th}_m(\Delta[1] \otimes \Delta[n])$ thus demonstrating that p is also in $P(|A|_m)$.

$P(|A|_m) \subseteq_r |P(A)|_m$: By definition we know that the complicial set of arrows of $|P(A)|_m$ is simply the complicial $(m-1)$ -superstructure $|P(A)_a|_{m-1}$ so it follows that the inclusion we wish to prove here holds if and only if the complicial set of arrows of $P(|A|_m)$ is $(m-1)$ -complicial. In other words, we need to demonstrate that for each $n > m-1$ any n -arrow p of $P(|A|_m)$ is thin. Consulting observation 177(2), we know that this will be the case if $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ maps the n -simplices $(\varepsilon_0^1 \circ \eta^n, \text{id}_{[n]})$ and $(\varepsilon_1^1 \circ \eta^n, \text{id}_{[n]};)$ and the $(n+1)$ -simplex $(\pi_1^{(1,n)}, \pi_2^{(1,n)})$ to thin simplices in A . However, since p is in $P(|A|_m)$ we know, from observation 183, that it extends to a stratified map $p: \text{Th}_m(\Delta[1] \otimes \Delta[n]) \longrightarrow A$ and since $n+1 > m$ it follows that p carries the $(n+1)$ -simplex above to a thin simplex in

A . Furthermore, since p is in $P(A)$ we may apply observation 180 to show that it maps both of the n -simplices above to thin simplices in A as required. $\square$

9.3. A Complicial Double Category with Connections. By iterating the category of prisms construction $\mathbb{P}$ we may construct an important complicial double category with connections.

Observation 185 (double prism categories). Of course, for each stratified set X the (partially) right adjoint functor $X \Rightarrow *$ preserves the (small) limits of $\underline{\mathbf{Cs}}$. It follows, by construction and the point-wise nature of limits in $\underline{\mathbf{Cat}}(\underline{\mathbf{Cs}})$, that the functor $\mathbb{P}: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Cat}}(\underline{\mathbf{Cs}})$ preserves all (small) limits. Consequently we may lift $\mathbb{P}$ to a LE-functor

$$\underline{\mathbf{Cat}}(\underline{\mathbf{Cs}}) \xrightarrow{\mathbb{P}} \underline{\mathbf{Cat}}(\underline{\mathbf{Cat}}(\underline{\mathbf{Cs}})) \stackrel{\text{def}}{=} \underline{\mathbf{Double}}(\underline{\mathbf{Cs}})$$

as discussed in definition 60. This constructs a vertically presented double category $\mathbb{P}(\mathbb{C})$ by applying $\mathbb{P}$ point-wise to the structural components of the complicial category $\mathbb{C}$. More explicitly, the double category $\mathbb{P}(\mathbb{C})$ has:

- complicial set of squares $\Delta[1] \Rightarrow \mathbb{C}_a$,
- complicial sets of vertical arrows $\mathbb{C}_a$ and horizontal arrows $\Delta[1] \Rightarrow \mathbb{C}_o$,
- complicial set of objects $\mathbb{C}_o$,
- vertical categories $\mathbb{C}$ of arrows and $\Delta[1] \Rightarrow \mathbb{C}$ of squares,
- horizontal categories $\mathbb{P}(\mathbb{C}_o)$ of arrows and $\mathbb{P}(\mathbb{C}_a)$ of squares,

Composing the functors $\mathbb{P}: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Cat}}(\underline{\mathbf{Cs}})$ and $\mathbb{P}: \underline{\mathbf{Cat}}(\underline{\mathbf{Cs}}) \longrightarrow \underline{\mathbf{Double}}(\underline{\mathbf{Cs}})$ we get a functor $\mathbb{P}^2: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Double}}(\underline{\mathbf{Cs}})$ and call $\mathbb{P}^2(A)$ the *double prism category* of the complicial set A .

Observation 186 (double prism categories under the microscope). From observation 65 we see that we may present $\mathbb{P}^2$ more symmetrically as $K_{\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}}}^l: \underline{\mathbf{Cs}} \longrightarrow \mathbb{T}_{\underline{\mathbf{Cat}}} \otimes \mathbb{T}_{\underline{\mathbf{Cat}}-\underline{\mathbf{Alg}}}(\underline{\mathbf{Cs}})$ (whose codomain is isomorphic to $\underline{\mathbf{Double}}(\underline{\mathbf{Cs}})$). Furthermore if we now apply observation 66, as we did in observation 177, we see that this in turn bears an explicit description as $K_{\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}} \otimes \Delta}: \underline{\mathbf{Cs}} \longrightarrow \mathbb{T}_{\underline{\mathbf{Cat}}} \otimes \mathbb{T}_{\underline{\mathbf{Cat}}} \otimes \mathbb{T}_{\underline{\mathbf{Cs}}-\underline{\mathbf{Alg}}}$ the external Kan functor. It follows that the complicial double category $\mathbb{P}^2(A)$ associated with a complicial set A bears the following explicit description:

- (1) The complicial set of squares of $\mathbb{P}^2(A)$ is $K_{\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}}}^l(A)([1], [1]) = (\Delta[1] \otimes \Delta[1]) \Rightarrow A$. Alternatively, using the explicit external presentation $K_{\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}} \otimes \Delta}(A)$ we see that the n -squares (that is the n -simplices in the complicial set of squares) of $\mathbb{P}^2(A)$ may be described as stratified maps of the form:

$$\Delta[1] \otimes \Delta[1] \otimes \Delta[n] \xrightarrow{\lambda} A \tag{49}$$

Furthermore, the action of a simplicial operator $\alpha: [m] \longrightarrow [n]$ on an n -square expressed in this form is given by $\lambda \cdot \alpha = \lambda \circ (\Delta[1] \otimes \Delta[1] \otimes \Delta(\alpha))$.

- (2) Consulting the internal presentation $K_{\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}}}^l(A)$ of $\mathbb{P}^2(A)$, it is clear that its horizontal and vertical categories of arrows of are both equal to $\mathbb{P}(A)$ and that its complicial set of objects is A itself. Furthermore, arguing just as in observation 177(4) we see that its sets of horizontal and vertical identity squares in are simply the regular subsets $(\Delta[1] \otimes \Delta[1]_t) \Rightarrow A$ and $(\Delta[1]_t \otimes \Delta[1]) \Rightarrow A$ of $(\Delta[1] \otimes \Delta[1]) \Rightarrow A$ respectively.

Applying the external presentation $K_{\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}} \otimes \Delta}(A)$ we see that the source and target maps for the horizontal and vertical categories of squares, when applied to the n -square

in display (49), may be given concretely by

$$\begin{aligned} s_h(\lambda) &= \lambda \circ (\Delta[1] \otimes \Delta(\varepsilon_0^1) \otimes \Delta[n]) & s_v(\lambda) &= \lambda \circ (\Delta(\varepsilon_0^1) \otimes \Delta[1] \otimes \Delta[n]) \\ t_h(\lambda) &= \lambda \circ (\Delta[1] \otimes \Delta(\varepsilon_1^1) \otimes \Delta[n]) & t_v(\lambda) &= \lambda \circ (\Delta(\varepsilon_1^1) \otimes \Delta[1] \otimes \Delta[n]) \end{aligned}$$

and if $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ is an n -arrow in $\mathbb{P}(A)$ then the horizontal and vertical identity squares on this are given by:

$$i_h(p) = p \circ (\Delta[1] \otimes \Delta(\eta^1) \otimes \Delta[n]) \quad i_v(p) = p \circ (\Delta(\eta^1) \otimes \Delta[1] \otimes \Delta[n])$$

Furthermore the horizontal and vertical identity squares of $\mathbb{P}^2(A)$ are those n -squares of the form shown in display (49) which extend to a stratified map with domain $\Delta[1] \otimes \Delta[1]_t \otimes \Delta[n]$ or $\Delta[1]_t \otimes \Delta[1] \otimes \Delta[n]$ respectively.

- (3) As in observation 177(4) if λ and λ' are a horizontally composable pair of n -squares then we may combine them to give a single stratified map $\Delta[1] \otimes \Delta[1]_t \otimes \Delta[n] \longrightarrow A$. and extend this along the f-extension $\Delta[1] \otimes \Delta[1]_t \otimes \Delta[n] \xrightarrow{\subseteq_r} \Delta[1] \otimes \Delta[2]_t \otimes \Delta[n]$ to form a *horizontal witness* $w_h(\lambda, \lambda'): \Delta[1] \otimes \Delta[2]_t \otimes \Delta[n] \longrightarrow A$. This witness is uniquely determined by the equations $w_h(\lambda, \lambda') \circ (\Delta[1] \otimes \Delta(\delta_2^2) \otimes \Delta[n]) = \lambda$ and $w_h(\lambda, \lambda') \circ (\Delta[1] \otimes \Delta(\delta_0^2) \otimes \Delta[n]) = \lambda'$ and we use it to define the horizontal composite by $\lambda' *_h \lambda = w_h(\lambda, \lambda') \circ (\Delta[1] \otimes \Delta(\delta_1^2) \otimes \Delta[n])$. Dually, if λ and λ' are vertically composable then we form a *vertical witness* $w_v(\lambda, \lambda'): \Delta[2]_t \otimes \Delta[1] \otimes \Delta[n] \longrightarrow A$ which is uniquely determined by the equations $w_v(\lambda, \lambda') \circ (\Delta(\delta_2^2) \otimes \Delta[1] \otimes \Delta[n]) = \lambda$ and $w_v(\lambda, \lambda') \circ (\Delta(\delta_0^2) \otimes \Delta[1] \otimes \Delta[n]) = \lambda'$ and we use it to define the vertical composite by $\lambda' *_v \lambda = w_v(\lambda, \lambda') \circ (\Delta(\delta_1^2) \otimes \Delta[1] \otimes \Delta[n])$.

The following theorem makes a pivotal contribution to understanding the structure of the double category $\mathbb{P}^2(A)$:

Theorem 187 ($\mathbb{P}^2(A)$ as a double category with connections). *If A is a complicial set then the double prism category $\mathbb{P}^2(A)$ possesses a canonical thinness structure $t\mathbb{P}^2(A)$ which extends $\mathbb{P}^2$ to a functor $\underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Double}}_T(\underline{\mathbf{Cs}})$. Furthermore, each $(\mathbb{P}^2(A), t\mathbb{P}^2(A))$ is a double category with connections in the sense of lemma 81.*

Proof. In order to construct the claimed thinness structure $t\mathbb{P}^2(A) \subseteq \mathbb{P}^2(A)$ (cf. definition 79), start by considering the f-almost $\mathbb{T}_{\mathbf{Cat}} \otimes \mathbb{T}_{\mathbf{Cat}}$ -coalgebra $\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}}$ which we used to define $\mathbb{P}^2$. Notice that the functor $\text{Th}_1: \underline{\mathbf{Strat}} \longrightarrow \underline{\mathbf{Strat}}$ preserves f-extensions, by lemma 171, and preserves colimits, since it is left adjoint to $|\cdot|_1$, from which it follows that it preserves all f-almost colimits. Consequently, the composite $\text{Th}_1 \circ (\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}}): \Delta \times \Delta \longrightarrow \underline{\mathbf{Strat}}$ is also an f-almost $\mathbb{T}_{\mathbf{Cat}} \otimes \mathbb{T}_{\mathbf{Cat}}$ -coalgebra, to which we may apply the construction of observation 63 in order to construct a second functor $t\mathbb{P}^2 \stackrel{\text{def}}{=} K_{\text{Th}_1 \circ (\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}})}^l$ from $\underline{\mathbf{Cs}}$ to $\underline{\mathbf{Double}}(\underline{\mathbf{Cs}})$.

Now, every stratified set X is an entire subset of the associated set $\text{Th}_1(X)$ so we may construct a canonical map $\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}} \longrightarrow \text{Th}_1 \circ (\Delta_{\mathbb{P}} \otimes \Delta_{\mathbb{P}})$ of our f-almost coalgebras (that is to say an arrow of $\mathbb{T}_{\mathbf{Cat}} \otimes \mathbb{T}_{\mathbf{Cat}}\text{-CoAlg}_{L_f}(\underline{\mathbf{Strat}})$) whose components are the entire inclusions $\Delta_{\mathbb{P}}([n]) \otimes \Delta_{\mathbb{P}}([m]) \xrightarrow{\subseteq_e} \text{Th}_1(\Delta_{\mathbb{P}}([n]) \otimes \Delta_{\mathbb{P}}([m]))$. As discussed in observation 63, this map of f-almost coalgebras gives rise to a derived natural transformation $K_{\subseteq_e}^l: t\mathbb{P}^2 \longrightarrow \mathbb{P}^2$ whose component at the complicial set A is obtained by applying $* \Rightarrow A$ to the entire inclusions of the last sentence and thus, by lemma 147, identifies $t\mathbb{P}^2(A)$ as a regular sub-double category of $\mathbb{P}^2(A)$.

Indeed we have $\text{Th}_1(\Delta_{\mathbb{P}}([1]) \otimes \Delta_{\mathbb{P}}([1])) = \text{Th}_1(\Delta[1] \otimes \Delta[1])$ so it follows that the set of squares of $t\mathbb{P}^2(A)$ is the regular subset $\text{Th}_1(\Delta[1] \otimes \Delta[1]) \Rightarrow A$ of the set of squares $(\Delta[1] \otimes$

$\Delta[1] \Rightarrow A$ of $\mathbb{P}(A)$. Also, by observation 133, we know that $\text{Th}_1(\Delta[1] \otimes \Delta[1])$ is an entire subset of both $\Delta[1]_t \otimes \Delta[1]$ and $\Delta[1] \otimes \Delta[1]_t$ so it follows, from observation 186, that the sets of horizontal and vertical identity squares in $\mathbb{P}^2(A)$ are actually regular subsets of the set of squares of $t\mathbb{P}^2(A)$. In other words, the pair $(\mathbb{P}^2(A), t\mathbb{P}^2(A))$ is indeed a double category with thinness as required.

Finally, by construction the regular inclusions $t\mathbb{P}^2(A) \xrightarrow{\subseteq_r} \mathbb{P}^2(A)$ are natural in A . In other words, if $f: A \rightarrow A'$ is a complicial map then the double functor $\mathbb{P}^2(f): \mathbb{P}^2(A) \rightarrow \mathbb{P}^2(A')$ lifts to a thinness preserving double functor $\mathbb{P}^2(f): (\mathbb{P}^2(A), t\mathbb{P}^2(A)) \rightarrow (\mathbb{P}^2(A'), t\mathbb{P}^2(A'))$, thus extending $\mathbb{P}^2$ to a functor from $\underline{\mathbf{Cs}}$ to the category of double categories with thinness $\underline{\mathbf{Double}}_T(\underline{\mathbf{Cs}})$ as stated.

To complete this proof, we must demonstrate that each double category with thinness $(\mathbb{P}^2(A), t\mathbb{P}^2(A))$ also satisfies the conditions given in lemma 81:

Conditions 81(i) and 81(ii). These two conditions have dual proofs, so we simply prove the latter regarding vertical arrows. Consider the following commutative square of inclusion maps

$$\begin{array}{ccc} \Delta[1] + \Delta[1] & \xrightarrow{\langle \Delta(\varepsilon_0^1) \otimes \Delta[1], \Delta(\varepsilon_1^1) \otimes \Delta[1] \rangle} & \text{Th}_1(\Delta[1] \otimes \Delta[1]) \\ \subseteq_e \downarrow & & \downarrow \subseteq_e \\ \Delta[1]_t + \Delta[1]_t & \xrightarrow{\langle \Delta(\varepsilon_0^1) \otimes \Delta[1], \Delta(\varepsilon_1^1) \otimes \Delta[1] \rangle} & \Delta[1] \otimes \Delta[1]_t \end{array}$$

in which the right hand vertical is an entire subset inclusion obtained by applying lemma 133 (with $n = 1$ and $m = 0$ we get $\text{Th}_1(\Delta[1] \otimes \Delta[1]) \subseteq_e \text{Th}_1(\Delta[1]) \otimes \text{Th}_0(\Delta[1]) = \Delta[1] \otimes \Delta[1]_t$) and the horizontal maps are those induced by the indicated stratified maps and the universal properties of the sums on the left hand side. Notice that the only simplices which are thin in $\Delta[1] \otimes \Delta[1]_t$ but not in $\text{Th}_1(\Delta[1] \otimes \Delta[1])$ are the 1-simplices $(\varepsilon_0^1 \circ \eta^1, \text{id}_{[1]})$ and $(\varepsilon_1^1 \circ \eta^1, \text{id}_{[1]})$, which are precisely the images of the 1-simplex $\text{id}_{[1]} \in \Delta[1]$ under the maps $\Delta(\varepsilon_0^1) \otimes \Delta[1]$ and $\Delta(\varepsilon_1^1) \otimes \Delta[1]$ respectively and it follows, by observation 158(a), that our square is a pasting square. Of course, if we apply the contravariant functor $* \Rightarrow A$ to this square it will carry colimits in the diagram to limits in $\underline{\mathbf{Cs}}$, giving us a pullback square

$$\begin{array}{ccc} (\Delta[1] \otimes \Delta[1]_t) \Rightarrow A & \xrightarrow{\subseteq_r} & \text{Th}_1(\Delta[1] \otimes \Delta[1]) \Rightarrow A \\ (\mathbf{s}_v, \mathbf{t}_v) \downarrow \lrcorner & & \downarrow (\mathbf{s}_v, \mathbf{t}_v) \\ (\Delta[1]_t \Rightarrow A) \times (\Delta[1]_t \Rightarrow A) & \xrightarrow{\subseteq_r} & (\Delta[1] \Rightarrow A) \times (\Delta[1] \Rightarrow A) \end{array}$$

In other words, the regular subset $(\Delta[1] \otimes \Delta[1]_t) \Rightarrow A$ of horizontal identity squares in $\mathbb{P}^2(A)$ is precisely the subset of $t\mathbb{P}^2(A)$ of those squares λ for which $\mathbf{s}_v(\lambda)$ and $\mathbf{t}_v(\lambda)$ are in the complicial subset of identities $\Delta[1]_t \Rightarrow A$ of the category $\mathbb{P}(A)$ of horizontal arrows, the desired result follows.

Conditions 81(iii) and 81(iv). These two conditions have dual proofs, so we simply prove the latter. It is easily demonstrated that we have a stratified map

$$\text{Th}_1(\Delta[1] \otimes \Delta[1]) \xrightarrow{\max} \Delta[1]$$

where $\max(\alpha, \beta)$ is the point-wise maximum of the simplicial operators $\alpha, \beta: [r] \longrightarrow [1]$, and that this satisfies the following identities:

$$\begin{aligned} \max \circ (\Delta[1] \otimes \Delta(\varepsilon_0^1)) &= \text{id}_{\Delta[1]} & \max \circ (\Delta(\varepsilon_0^1) \otimes \Delta[1]) &= \text{id}_{\Delta[1]} \\ \max \circ (\Delta[1] \otimes \Delta(\varepsilon_1^1)) &= \Delta(\varepsilon_1^1 \circ \eta^1) & \max \circ (\Delta(\varepsilon_1^1) \otimes \Delta[1]) &= \Delta(\varepsilon_1^1 \circ \eta^1) \end{aligned}$$

Applying the contravariant functor $* \Rightarrow A$ we get a stratified map $\max \Rightarrow A$ whose domain $\Delta[1] \Rightarrow A$ is the set of vertical arrows of $\mathbb{P}^2(A)$ and whose codomain is $t\mathbb{P}^2(A)$. Furthermore, applying $* \Rightarrow A$ to the identities above, and using the definitions of the various source and target maps of $\mathbb{P}(A)$ and $\mathbb{P}^2(A)$, we get the identities:

$$\begin{aligned} s_h \circ (\max \Rightarrow A) &= \text{id}_{\Delta[1] \Rightarrow A} & s_v \circ (\max \Rightarrow A) &= \text{id}_{\Delta[1] \Rightarrow A} \\ t_h \circ (\max \Rightarrow A) &= i \circ t & t_v \circ (\max \Rightarrow A) &= i \circ t \end{aligned}$$

In other words if q is a vertical arrow of $\mathbb{P}^2(A)$ then the thin square $\nu_q \stackrel{\text{def}}{=} (\max \Rightarrow A)(q)$ has horizontal and vertical source q and horizontal and vertical targets which are identities in $\mathbb{P}(A)$. This makes it a suitable candidate to be the connection square of condition 81(iv) as required. $\square$

10. DECALAGE FOR COMPLICIAL SETS AND COMPLICIALLY ENRICHED CATEGORIES

10.1. A Decalage Construction on Complicial Sets. We show that we may obtain the appropriate generalisation of the decalage construction to $\underline{\mathbf{Cs}}$ as a sub-functor of $\Delta[1] \Rightarrow *$.

Definition 188 (decalage on $\underline{\mathbf{Cs}}$). If A is a complicial set then define $\text{Dec}(A)$, its *decalage*, to be the complicial set obtained by taking the following pullback in $\underline{\mathbf{Cs}}$:

$$\begin{array}{ccc} \text{Dec}(A) & \xrightarrow{\subseteq_r} & \Delta[1] \Rightarrow A \\ \downarrow \lrcorner & & \downarrow \Delta(\varepsilon_0^1) \Rightarrow A \\ |A|_0 & \xrightarrow[\subseteq_r]{} & A \end{array} \quad (50)$$

As a functor, Dec is the point-wise pullback, in the functor category $[\underline{\mathbf{Cs}}, \underline{\mathbf{Cs}}]$, of the natural inclusion $|\cdot|_0 \xrightarrow{\subseteq_r} \text{id}_{\underline{\mathbf{Cs}}}$ along the natural transformation $\Delta(\varepsilon_0^1) \Rightarrow *: \Delta[1] \Rightarrow * \longrightarrow \text{id}_{\underline{\mathbf{Cs}}}$. In other words, for each stratified map $f: A \longrightarrow B$ the map $\Delta[1] \Rightarrow f: \Delta[1] \Rightarrow A \longrightarrow \Delta[1] \Rightarrow B$ restricts to a map $\text{Dec}(f): \text{Dec}(A) \longrightarrow \text{Dec}(B)$, thereby making our construction into a functor $\text{Dec}: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Cs}}$ for which the family of regular subset inclusions $\text{Dec}(A) \xrightarrow{\subseteq_r} \Delta[1] \Rightarrow A$ constitute a natural transformation from Dec to $\Delta[1] \Rightarrow *$.

Using the explicit description of the n -simplices of $\Delta[1] \Rightarrow A$ of observation 145, we may describe $\text{Dec}(A)$ as the regular subset of $\Delta[1] \Rightarrow A$ on those simplices $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ for which the simplex $(\Delta(\varepsilon_0^1) \Rightarrow A)(p) = p(\varepsilon_0^1 \circ \eta^n, \text{id}_{[n]})$ is in the regular subset $|A|_0 \subseteq_r A$.

Another way of re-stating this characterisation is to construct an entire superset $(\Delta[1] \otimes \Delta[n])'$ of $\Delta[1] \otimes \Delta[n]$ by making thin every simplex $(\alpha, \beta) \in \Delta[1] \otimes \Delta[n]$ which has dimension $r > 0$ and $\alpha(i) = 0$ for all $i \in [r]$. Then an n -simplex $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ in $\Delta[1] \Rightarrow A$ is in the regular subset $\text{Dec}(A)$ if and only if it extends to a stratified map $p: (\Delta[1] \otimes \Delta[n])' \longrightarrow A$.

We make Dec into a copointed endo-functor on $\underline{\mathbf{Cs}}$ by defining the counit $\partial: \text{Dec} \longrightarrow \text{id}_{\underline{\mathbf{Cs}}}$ to be the natural transformation obtained by composing the inclusion $\text{Dec} \xrightarrow{\subseteq_r} \Delta[1] \Rightarrow *$ with the natural transformation $\Delta(\varepsilon_1^1) \Rightarrow *: \Delta[1] \Rightarrow * \longrightarrow \text{id}_{\underline{\mathbf{Cs}}}$. In terms of the explicit presentation of simplices of $\text{Dec}(A)$, this counit maps an n -simplex $p: \Delta[1] \otimes \Delta[n] \longrightarrow A$ of $\text{Dec}(A)$ to the simplex $\partial_A(p) = (\Delta(\varepsilon_1^1) \Rightarrow A)(p) = p(\varepsilon_1^1 \circ \eta^n, \text{id}_{[n]})$ of A .

Notation 189. In the subsequent arguments we will have occasion to consider, and relate, three distinct decalage constructions, each one on a different category. In particular, we now have the classical decalage comonad on $\underline{\mathbf{Simp}}$, which we discussed in lemma 89, and the construction on $\underline{\mathbf{Cs}}$, introduced in the previous definition. We will be adding to these later in this subsection by describing an analogous copointed endo-functor on $\underline{\mathbf{Cs-Cat}}$.

In order to reduce notational clutter, we will overload the symbols Dec and ∂ in order to use them to denote the decalage construction on whichever of these categories is currently under consideration. Consequently, we will also be taking care to ensure that the reader may disambiguate this notation contextually.

Theorem 190. Suppose that $U: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Simp}}$ denotes the forgetful functor, which maps a complicial set to its underlying simplicial set, then there exists a natural isomorphism

$$\begin{array}{ccc} \underline{\mathbf{Cs}} & \xrightarrow{\text{Dec}} & \underline{\mathbf{Cs}} \\ U \downarrow & \cong \Downarrow \varphi & \downarrow U \\ \underline{\mathbf{Simp}} & \xrightarrow[\text{Dec}]{} & \underline{\mathbf{Simp}} \end{array} \quad (51)$$

in Cat which makes the pair (U, φ) into a strong transformation of copointed endo-functors from the decalage construction on Cs to that on Simp.

Most of the remainder of this subsection is devoted to proving this theorem, which we do via a sequence of observations and lemmas:

Observation 191 (strong transformations of copointed endo-functors). Transformations of copointed endo-functors are defined analogously to the comonad transformations discussed in observation 92. That is, the square of display (51) represents an endo-1-cell of the 2-category Cyl on the forgetful functor $U: \underline{\text{Cs}} \longrightarrow \underline{\text{Simp}}$ and we require that the pair consisting of the counits associated with the functor Dec on the categories Cs and Simp (respectively) is a cylinder (2-cell) from this endo-1-cell to the identity 1-cell on U .

Explicitly, this latter condition simply requires that the natural transformation φ must satisfy the equation $U\partial = (\partial U) \cdot \varphi$. Here it is worth observing that the use context of the symbol ∂ on either side of this equation implies that these instances represent different natural transformations. Since U is a functor from Cs to Simp we know that ∂ in the context $U\partial$ must be the counit for Dec on Cs and, conversely, that in the context ∂U it instead represents the counit of Dec on Simp.

The strong transformation postulated in theorem 190 simply demonstrates that the construction in definition 188 really deserves to be characterised as a generalisation of the decalage construction to the category of complicial sets.

Assuming for the remainder of this observation that theorem 190 holds and consulting observation 111, we know that the set of connected components of a complicial set is dependent only on the structure of its underlying simplicial set. In other words, we have a commuting triangle of functors

$$\begin{array}{ccc} \underline{\text{Cs}} & \xrightarrow{U} & \underline{\text{Simp}} \\ & \searrow \Pi_0 & \swarrow \Pi_0 \\ & \underline{\text{Set}} & \end{array}$$

and so we may apply the semi-simplicial version of lemma 93 to the strong transformation postulated in lemma 190 to give a natural isomorphism:

$$\begin{array}{ccc} \underline{\text{Cs}} & \xrightarrow{U} & \underline{\text{Simp}} \\ & \searrow N_{\text{Dec}, \Pi_0} & \swarrow N_{\text{Dec}, \Pi_0} \\ & \underline{\text{SSimp}} & \end{array}$$

Applying the semi-simplicial version of lemma 89 we may show that the nerve functor $N_{\text{Dec}, \Pi_0}: \underline{\text{Simp}} \longrightarrow \underline{\text{SSimp}}$ is isomorphic to the forgetful functor from Simp to SSimp. It follows from the triangle above that the nerve functor $N_{\text{Dec}, \Pi_0}: \underline{\text{Cs}} \longrightarrow \underline{\text{SSimp}}$ associated with the construction introduced in definition 188 is isomorphic to the forgetful functor from the category of complicial sets Cs to the category of semi-simplicial sets SSimp.

Observation 192 (constructing the natural transformation φ of theorem 190). We define the isomorphism φ in two steps, here we construct a (candidate) inverse $\varphi^*: \text{Dec} \circ U \longrightarrow U \circ \text{Dec}$ and in subsequent lemmas we demonstrate that this is indeed a natural isomorphism.

Consulting the construction of definition 88, and applying Yoneda's lemma for stratified sets, we see that if A is a complicial set then an n -simplex of $\text{Dec}(U(A))$ may be identified with

a stratified map $x: \Delta[n+1] \longrightarrow A$. Furthermore, under this identification the right action of simplicial operator ϕ on our n -simplex x in $\text{Dec}(\text{U}(A))$ is given by $x \cdot \phi = x \circ \Delta(\phi \oplus [0])$.

Of course we also know, from definition 188, that an n -simplex of $\text{U}(\text{Dec}(A))$ may be identified with a stratified map $p: (\Delta[1] \otimes \Delta[n])' \longrightarrow A$ and that under this identification $p \cdot \phi = p \circ (\Delta[1] \otimes \Delta(\phi))$ as discussed in observation 145.

Thus we are drawn to considering the relationship between $\Delta[n+1]$ and $(\Delta[1] \otimes \Delta[n])'$, so starting with an r -simplex (α, β) of $\Delta[1] \otimes \Delta[n]$ we define the r -simplex $e_n(\alpha, \beta)$ of $\Delta[n+1]$ by

$$(e_n(\alpha, \beta))(i) = \begin{cases} 0 & \text{if } \alpha(i) = 0, \\ \beta(i) + 1 & \text{if } \alpha(i) = 1. \end{cases}$$

The point-wise nature of this construction ensures that e_n respects the right actions of Δ on $\Delta[1] \otimes \Delta[n]$ and $\Delta[n+1]$ since these are given by pre-composition, in other words e_n is a simplicial map. Furthermore, e_n maps any simplex (α, β) which has $\alpha(i) = 0$ for some $i > 0$ to a degenerate simplex in $\Delta[n+1]$. It follows, directly from the analysis of lemma 130 and the definition of $(\Delta[1] \otimes \Delta[n])'$ in definition 88, that e_n carries thin simplices in $(\Delta[1] \otimes \Delta[n])'$ to degenerate simplices in $\Delta[n+1]$ and is thus a stratified map $e_n: (\Delta[1] \otimes \Delta[n])' \longrightarrow \Delta[n+1]$. We may now use this to define a function

$$\begin{array}{ccc} \text{Dec}(\text{U}(A))_n & \xrightarrow{(\varphi_A)_n} & \text{U}(\text{Dec}(A))_n \\ \parallel & \text{def} \parallel & \parallel \\ \text{Strat}(\Delta[n+1], A) & \xrightarrow{\text{Strat}(e_n, A)} & \text{Strat}((\Delta[n] \otimes \Delta[1])', A) \end{array}$$

which is natural in A by construction. Observe also that if $\phi: [m] \longrightarrow [n]$ is a simplicial operator then it is a matter of routine calculation to demonstrate that the following square of stratified maps commutes:

$$\begin{array}{ccc} (\Delta[1] \otimes \Delta[m])' & \xrightarrow{e_m} & \Delta[m+1] \\ \Delta[1] \otimes \Delta(\phi) \downarrow & & \downarrow \Delta(\phi \oplus [0]) \\ (\Delta[1] \otimes \Delta[n]) & \xrightarrow{e_n} & \Delta[n+1] \end{array} \quad (52)$$

Returning to the explicit descriptions furnished by the first couple of paragraphs of this observation, it is clear that we may use the commutativity of these squares to show that the collection of maps $(\varphi_A^*)_n$ ($n \in \mathbb{N}$) respect the actions of Δ on $\text{Dec}(\text{U}(A))$ and $\text{U}(\text{Dec}(A))$ and thus provide us with the components of a simplicial map $\varphi_A^*: \text{Dec}(\text{U}(A)) \longrightarrow \text{U}(\text{Dec}(A))$ as desired. To do so, simply consider an arbitrary simplicial operator ϕ and observe that we have $(\varphi_A^*)_n(x) \cdot \phi = (x \circ e_n) \cdot \phi = x \circ e_n \circ (\Delta(\phi) \otimes \Delta[1]) =^* x \circ \Delta([0] \oplus \phi) \circ e_m = (x \cdot \phi) \circ e_m = (\varphi_A^*)_m(x \cdot \phi)$ wherein the starred equality follows from (52).

Observation 193. By definition the component $(\varphi_A^*)_n = \text{Strat}(e_n, A)$ is an isomorphism iff e_n is orthogonal to A , so it follows that $\varphi_A^*: \text{Dec}(\text{U}(A)) \longrightarrow \text{U}(\text{Dec}(A))$ is an isomorphism for each complicial set A iff the stratified map $e_n: (\Delta[n] \otimes \Delta[1])' \longrightarrow \Delta[n+1]$ is an f-extension for each $n \in \mathbb{N}$. Consequently, all that remains in proving theorem 190 is to establish this latter condition.

To this end, define $(\Delta[1] \otimes \Delta[n])''$ to be the entire superset of $\Delta[1] \otimes \Delta[n]$ in which we make thin all those simplices (α, β) for which $\alpha(i) = 0$ for some $i > 0$. As discussed in the

last observation, the map e_n actually maps each of the thin simplices in $(\Delta[1] \otimes \Delta[n])''$ to a degenerate (and thus thin) simplex in $\Delta[n+1]$. However a little more is actually true, in fact e_n extends to a *regular* stratified map $(\Delta[1] \otimes \Delta[n])'' \longrightarrow \Delta[n+1]$.

We'll actually prove that $e_n: (\Delta[1] \otimes \Delta[n])' \longrightarrow \Delta[n+1]$ is an f-extension by decomposing our proof into two steps

- first show that the entire inclusion $(\Delta[1] \otimes \Delta[n])' \subseteq_e (\Delta[1] \otimes \Delta[n])''$ is a t-extension,
- then prove that the regular map $e_n: (\Delta[1] \otimes \Delta[n])'' \longrightarrow \Delta[n+1]$ is an f-extension,

and finally appeal to the fact that the composite of these two, $e_n: (\Delta[1] \otimes \Delta[n])' \longrightarrow \Delta[n+1]$ itself, is therefore an f-extension.

Lemma 194. *The entire inclusion $(\Delta[1] \otimes \Delta[n])' \xrightarrow{\subseteq_e} (\Delta[1] \otimes \Delta[n])''$ is a t-extension (and thus an f-extension).*

Proof. We aim to apply the explicit characterisation of t-extensions given in observation 126, so we define a sequence of subsets $\{tX_k\}_{k \in [n]}$ of $\Delta[1] \otimes \Delta[n]$ by:

$$tX_k \stackrel{\text{def}}{=} (t\Delta[1] \otimes t\Delta[n]) \cup \{(\alpha, \beta) \in \Delta[1] \otimes \Delta[n] \mid \dim(\alpha, \beta) \geq n - k + 1 \wedge \alpha(n - k + 1) = 0\}$$

It is a straightforward matter, using the fact that each α in the definition above is order preserving, to show that tX_0 is the set of thin simplices of $(\Delta[1] \otimes \Delta[n])'$, that tX_n is the set of thin simplices of $(\Delta[1] \otimes \Delta[n])''$ and that for all $0 \leq k < n$ we have $tX_k \subseteq tX_{k+1}$.

To verify the remaining condition of observation 126, suppose that $0 \leq k < n$ and that (α, β) is an r -simplex in $tX_{k+1} \setminus tX_k$ then from the definitions of these sets we know that $r \geq n - k > 0$, $\alpha(n - k) = 0$ and $\alpha(n - k + 1) = 1$. Furthermore, we also know that $\beta(n - k + 1) > \beta(n - k)$ since otherwise $(n - k)$ would witness that fact that (α, β) was a mediator simplex and thus an element of $t\Delta[1] \otimes t\Delta[n]$ (cf. observation 134).

So define a new $(r + 1)$ -simplex $(\alpha', \beta') \stackrel{\text{def}}{=} (\alpha \circ \sigma_{n-k}^r, \beta \circ \sigma_{n-k+1}^r)$ and observe that:

- The integer $(n - k + 1)$ witnesses the fact that (α', β') is a mediator simplex and is thus $(n - k + 1)$ -admissible.
- The $(n - k)^{\text{th}}$ face $(\alpha', \beta') \cdot \delta_{n-k}^{r+1}$ is equal to $(\alpha, \beta \circ \sigma_{n-k+1}^r \circ \delta_{n-k}^{r+1})$ which is a mediator simplex witnessed by $n - k$ and is thus an element of $t\Delta[1] \otimes t\Delta[n] \subseteq tX_k$.
- The $(n - k + 1)^{\text{th}}$ face $(\alpha', \beta') \cdot \delta_{n-k+1}^{r+1}$ is equal to our original simplex (α, β) .
- The $(n - k + 2)^{\text{th}}$ face $(\alpha', \beta') \cdot \delta_{n-k+2}^{r+1}$ is equal to $(\alpha \circ \sigma_{n-k}^r \circ \delta_{n-k+2}^{r+1}, \beta)$ which is an element of tX_k since $\alpha \circ \sigma_{n-k}^r \circ \delta_{n-k+2}^{r+1}(n - k + 1) = \alpha(n - k) = 0$.

In other words, (α', β') witnesses the extension of thinness to (α, β) as required by observation 126, which we may now apply to establish the proposition. $\square$

Lemma 195. *The stratified map $e_n: (\Delta[1] \otimes \Delta[n])'' \longrightarrow \Delta[n+1]$ is an f-extension.*

Proof. Start by considering the $(n+1)$ -simplex $(\rho_1^{n+1}, \sigma_0^n)$ of $(\Delta[1] \otimes \Delta[n])''$, where the simplicial operators $\rho_k^n: [r] \longrightarrow [1]$ are as defined in observation 165. From the definition of e_n , given in observation 192, we see that it carries $(\rho_1^{n+1}, \sigma_0^n)$ to the simplex $\text{id}_{[n+1]}$ in $\Delta[n+1]$ so, applying Yoneda's lemma, we see that the map $i_n = \ulcorner(\rho_1^{n+1}, \sigma_0^n)\urcorner: \Delta[n+1] \longrightarrow (\Delta[1] \otimes \Delta[n])''$ is a right inverse to $e_n: (\Delta[1] \otimes \Delta[n])'' \longrightarrow \Delta[n+1]$.

Consequently, we may argue as we did in the proof of corollary 164 and show that our desired result holds iff whenever A is a complicial set and $g, g': (\Delta[1] \otimes \Delta[n])'' \longrightarrow A$ are stratified maps with $g \circ i_n = g' \circ i_n$ which, by Yoneda's lemma, is the same as saying that $g(\rho_1^{n+1}, \sigma_0^n) = g'(\rho_1^{n+1}, \sigma_0^n)$ then $g = g'$.

But since the underlying simplicial set of $(\Delta[1] \otimes \Delta[n])''$ is $\Delta[1] \times \Delta[n]$ we know, by observation 45, that every one of its simplices is a face of some shuffle. It follows that the stratified maps g and g' coincide iff they act identically on each of the shuffles $(\rho_{k+1}^{n+1}, \sigma_k^n)$ of $\Delta[n] \times \Delta[1]$ ($k = 0, \dots, n$).

Now fix $0 < k \leq n$ and consider the simplex $(\rho_{k+1}^{n+1}, \sigma_k^n)$. Suppose that $\alpha: [r] \longrightarrow [n+1]$ is a face operator whose image contains $k-1$ and k , let $l \in [r]$ be the unique integer such that $\alpha(l) = k$ and consider the face $(\rho_{k+1}^{n+1}, \sigma_k^n) \cdot \alpha = (\rho_{k+1}^{n+1} \circ \alpha, \sigma_k^n \circ \alpha)$. It is clear that $l > 0$ and that $\rho_{k+1}^{n+1} \circ \alpha(l) = 0$ so it follows that this face is a thin simplex in $(\Delta[1] \otimes \Delta[n])''$. Quantifying over these face operators α it follows that our $(\rho_{k+1}^{n+1}, \sigma_k^n)$ of $(\Delta[1] \otimes \Delta[n])''$ is pre-degenerate at $k-1$ in there.

Returning to our stratified map $g: (\Delta[1] \otimes \Delta[n])'' \longrightarrow A$, we know that it preserves pre-degeneracy and so we may infer that the simplex $g(\rho_{k+1}^{n+1}, \sigma_k^n)$ in A is pre-degenerate at $k-1$. However since A is complicial, and thus well tempered by lemma 163, we may infer that this simplex is in fact degenerate at $k-1$.




FIGURE 6. An adjacent pair of shuffles in $\Delta[1] \otimes \Delta[n]$

Figure 6 depicts our shuffle $(\rho_{k+1}^{n+1}, \sigma_k^n)$ and the immediately preceding one $(\rho_k^{n+1}, \sigma_{k-1}^n)$. It clearly illustrates that they share the same k^{th} n -face and that this is the simplex $(\rho_k^n, \text{id}_{[n]})$, facts which may be verified by straightforward calculations with simplicial operators. From this, and the fact that g is a simplicial map, we immediately see that $g(\rho_k^n, \text{id}_{[n]})$ is the k^{th} n -face of $g(\rho_k^{n+1}, \sigma_{k-1}^n)$ and that the degeneracy result of the previous paragraph implies that the simplex $g(\rho_{k+1}^{n+1}, \sigma_k^n)$ is equal to its degenerated face $g(\rho_k^n, \text{id}_{[n]}) \cdot \sigma_{k-1}^n$. Combining these two, we get the equality $g(\rho_{k+1}^{n+1}, \sigma_k^n) = g(\rho_k^{n+1}, \sigma_{k-1}^n) \cdot (\delta_k^{n+1} \circ \sigma_{k-1}^n)$ which demonstrates that the action of g on the shuffle $(\rho_{k+1}^{n+1}, \sigma_k^n)$ is completely determined by its action on the adjacent shuffle $(\rho_k^{n+1}, \sigma_{k-1}^n)$.

Finally we can apply this result repeatedly for $k = n, n-1, \dots, 1$ to show that the action of g on each shuffle is determined by its action on the shuffle that immediately precedes it (under the linear ordering $\triangleleft$) and that, ultimately, the action of g on all shuffles, and thus on all simplices of $(\Delta[1] \otimes \Delta[n])''$, is completely determined by its action on the $\triangleleft$ -minimal shuffle $(\rho_1^{n+1}, \sigma_0^n)$. Of course, we can also apply exactly the same argument to g' and we know from

our initial assumption that these two stratified maps act identically on $(\rho_1^{n+1}, \sigma_0^n)$, it follows therefore that they must be identical maps as required. $\square$

Proof. (of theorem 190). Finally this follows on observing that the composite of the two f-extensions provided by these lemmas is the stratified map $e_n: (\Delta[n] \otimes \Delta[1])' \longrightarrow \Delta[n+1]$ which is therefore also an f-extension as required by observation 193. That the pair (U, φ) satisfies the cylinder condition required of a transformation of copointed endo-functors is a matter of trivial verification, directly from the definition of φ^* , which we leave to the reader. $\square$

10.2. A Path Construction on Complicially Enriched Categories. Now we turn to our primary motivation for introducing the double category with connection $(\mathbb{P}^2(A), t\mathbb{P}^2(A))$ in the previous section, that is to build an analogue of the functor $\Delta[1] \Rightarrow *$ on $\underline{\text{Cs-Cat}}$. In the next subsection we use this analogue to emulate the construction of the last subsection on $\underline{\text{Cs-Cat}}$.

Observation 196 (what do we mean by an “analogue” of $\Delta[1] \Rightarrow *$). First we should make precise the sense in which we shall be “building an analogue of $\Delta[1] \Rightarrow *: \underline{\text{Cs}} \longrightarrow \underline{\text{Cs}}$ on $\underline{\text{Cs-Cat}}$ ”. In fact, our intention is to construct a functor $I: \underline{\text{Cs-Cat}} \longrightarrow \underline{\text{Cs-Cat}}$ and a pair of natural transformations $\partial^0, \partial^1: I \longrightarrow \text{id}_{\underline{\text{Cs-Cat}}}$ and that this data should come equipped with a natural isomorphism

$$\begin{array}{ccc} \underline{\text{Cs}} & \xrightarrow{\Delta[1] \Rightarrow *} & \underline{\text{Cs}} \\ \text{P} \downarrow & \cong \Downarrow \theta & \downarrow \text{P} \\ \underline{\text{Cs-Cat}} & \xrightarrow{I} & \underline{\text{Cs-Cat}} \end{array} \quad (53)$$

making the triangles

$$\begin{array}{ccc} \text{P}(\Delta[1] \Rightarrow A) & \xrightarrow[\cong]{\theta_A} & I(\text{P}(A)) \\ & \searrow \text{P}(\Delta(\varepsilon_0^1) \Rightarrow A) \quad \swarrow \partial_{\text{P}(A)}^0 & \\ & \text{P}(A) & \end{array} \quad \begin{array}{ccc} \text{P}(\Delta[1] \Rightarrow A) & \xrightarrow[\cong]{\theta_A} & I(\text{P}(A)) \\ & \searrow \text{P}(\Delta(\varepsilon_1^1) \Rightarrow A) \quad \swarrow \partial_{\text{P}(A)}^1 & \\ & \text{P}(A) & \end{array} \quad (54)$$

commute for each compcial set A .

Our approach to building such a functor will be to start by identifying $\text{P}(\Delta[1] \Rightarrow A)$ as a substructure of $\mathbb{P}^2(A)$. We then show that when we may apply the functor $\mathbb{P}: \underline{\text{Cs}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ “point-wise” to a compcially enriched category, as described in definition 60, we obtain a double category the reflection dual of which is in fact obtain a 2-category and that in the particular case of $\mathbb{P}(\text{P}(A))$ this is also a substructure of $\mathbb{P}^2(A)$. Finally we show that we may apply the double category of squares construction to $\mathbb{P}(\text{P}(A))^r$ to re-construct a part of $\mathbb{P}^2(A)$ which contains $\text{P}(\Delta[1] \Rightarrow A)$.

When handling path categories it will be convenient to restrict our attention to a certain sub-double category of $\mathbb{P}^2(A)$:

Definition 197. Let $\text{P}^{(2)}(A)$ denote the sub-double category of $\mathbb{P}^2(A)$ consisting of those squares λ for which each one of the four “corner” objects $\mathbf{s}(s_h(\lambda)) = \mathbf{s}(s_v(\lambda))$, $\mathbf{s}(t_h(\lambda)) = \mathbf{t}(s_v(\lambda))$, $\mathbf{t}(s_h(\lambda)) = \mathbf{s}(t_v(\lambda))$ and $\mathbf{t}(t_h(\lambda)) = \mathbf{t}(t_v(\lambda))$ are simplices in the 0-superstructure

$|A|_0$ of A . Alternatively, a square λ is in $P^{(2)}(A)$ iff $s_h(\lambda)$ and $t_h(\lambda)$ are both in the complicial subcategory $P(A) \subseteq_r \mathbb{P}(A)$ or, equivalently, iff $s_v(\lambda)$ and $t_v(\lambda)$ are both in $P(A)$.

That this set of squares is indeed closed in $\mathbb{P}^2(A)$ under its horizontal and vertical category structures is a trivial consequence of the fact that the corner objects of composites, sources and targets of squares are all corners of one or other of the original squares being operated upon (cf. observation 73). So if the original squares all have corners in $|A|_0$ then so do their composites etcetera.

Furthermore, let $tP^{(2)}(A)$ denote the intersection $P^{(2)}(A) \cap t\mathbb{P}^2(A)$ then, by the same reasoning as the last paragraph applied to theorem 187, it follows that $(P^{(2)}(A), tP^{(2)}(A))$ is also a double category with connection.

Observation 198 ($P(\Delta[1] \Rightarrow A)$ as a substructure of $P^{(2)}(A)$). Suppose that $\mathbb{D}$ is a complicial double category, then we may define an associated double category $E_h(\mathbb{D})$ by applying the co-reflector $E: \underline{\text{Cat}}(\underline{\text{Cs}}) \longrightarrow \underline{\text{Cs}}\text{-}\underline{\text{Cat}}$ point-wise to the vertical presentation of $\mathbb{D}$ (cf. definition 60). In other words, $E_h(\mathbb{D})$ is the regular sub-double category of $\mathbb{D}$ of those squares and horizontal arrows whose horizontal source and target are both in the superstructure $|\text{arr}_v(\mathbb{D})|_0 \subseteq_r \text{arr}_v(\mathbb{D})$. Observe also that, by definition, its horizontal categories of arrows and squares are actually complicially enriched, since they are $E(\text{arr}_h(\mathbb{D}))$ and $E(\text{sq}_h(\mathbb{D}))$ respectively.

From observation 185, we know that the vertical presentation of $\mathbb{P}^2(A)$ is constructed by applying the left exact functor $\mathbb{P}: \underline{\text{Cs}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ point-wise to the prism category $\mathbb{P}(A) \in \underline{\text{Cat}}(\underline{\text{Cs}})$. It follows, therefore, that the vertical presentation of $E_h(\mathbb{P}^2(A))$ is obtained by applying $P = E \circ \mathbb{P}$ point-wise to $\mathbb{P}(A)$, or in other words it is simply the double category we would usually refer to as $P(\mathbb{P}(A))$.

Since the complicial sets of arrows and objects of $\mathbb{P}(A)$ are $\Delta[1] \Rightarrow A$ and A respectively it follows that the horizontal categories of squares and arrows of $P(\mathbb{P}(A))$ are $P(\Delta[1] \Rightarrow A)$ and $P(A)$ respectively. Additionally, the source and target maps of $\mathbb{P}(A)$ are $s = \Delta(\varepsilon_0^1) \Rightarrow A$ and $t = \Delta(\varepsilon_1^1) \Rightarrow A$ therefore the vertical presentation of $P(\mathbb{P}(A))$ has vertical source and target functors given by $s_v = P(\Delta(\varepsilon_0^1) \Rightarrow A)$ and $t_v = P(\Delta(\varepsilon_1^1) \Rightarrow A)$. These observations explain our interest in the sub-double category $P(\mathbb{P}(A))$, in short it provides us with a convenient encapsulation of the structures on the left hand side of the triangles in display (54) of observation 196 in terms of the vertical presentation of the double category $\mathbb{P}^2(A)$.

Finally, consulting definition 197 we see that the fact that $P(\mathbb{P}(A))$ has $P(A)$ as its horizontal category of arrows implies that it is actually contained within the sub-double category $P^{(2)}(A)$ of $\mathbb{P}^2(A)$ and it follows immediately that we actually have $P(\mathbb{P}(A)) = E_h(P^{(2)}(A))$.

Observation 199 (2-categories from complicially enriched categories). Suppose that $\mathbb{C}$ is a complicially enriched category and consider the complicial double category $\mathbb{P}(\mathbb{C})$ discussed in observation 185. We know that its vertical presentation is constructed by applying the functor $\mathbb{P}: \underline{\text{Cs}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ “point-wise” to the structural components of $\mathbb{C}$ and in particular that its complicial category of horizontal arrows $\text{arr}_h(\mathbb{P}(\mathbb{C}))$ is equal to $\mathbb{P}(\mathbb{C}_o)$. Of course since $\mathbb{C}$ is complicially enriched we know, by the definition in observation 178, that its set of objects $\mathbb{C}_o$ is 0-complicial and so we can infer, by observation 181, that $\mathbb{P}(\mathbb{C}_o)$ is the discrete category on $\mathbb{C}_o$ itself. It follows therefore that the reflection dual $\mathbb{P}(\mathbb{C})^r$ is a complicial 2-category.

Consider now the path category $P(A)$. We know that it is a regular subcategory of $\mathbb{P}(A)$, so it follows that the complicial double category $\mathbb{P}(P(A))$ is a regular sub-double category of $\mathbb{P}^2(A)$. More precisely, from observation 185 we know that the horizontal category of squares of $\mathbb{P}(P(A))$ is formed by applying the functor $\mathbb{P}: \underline{\text{Cs}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ to the complicial set $P(A)_a$

which was defined as a pullback in observation 178. However $\mathbb{P}$ preserves pullbacks and it follows that the complicial category $\text{sq}_h(\mathbb{P}(\mathbb{P}(A))) = \mathbb{P}(\mathbb{P}(A)_a)$ is the regular subcategory of $\text{sq}_h(\mathbb{P}^2(A)) = \mathbb{P}(\mathbb{P}(A)_a)$ obtained by pulling the discrete subcategory $\mathbb{P}(|A|_0) \times \mathbb{P}(|A|_0)$ of $\mathbb{P}(A) \times \mathbb{P}(A)$ back along the map $(\mathbb{P}(\mathbf{s}), \mathbb{P}(\mathbf{t})) = (\mathbf{s}_v, \mathbf{t}_v): \text{sq}_h(\mathbb{P}^2(A)) \longrightarrow \mathbb{P}(A) \times \mathbb{P}(A)$.

However, observation 181 tells us that $\mathbb{P}(|A|_0)$ is actually the discrete category on $|A|_0$, so it follows that $\mathbb{P}(\mathbb{P}(A))$ is simply the regular sub-double category of $\mathbb{P}^2(A)$ consisting of those squares λ whose vertical source $\mathbf{s}_v(\lambda)$ and target $\mathbf{t}_v(\lambda)$ are each horizontal identities in $\mathbb{P}(A)$ on some object in $|A|_0$. Taking reflection duals, it follows that we can summarise this description of $\mathbb{P}(\mathbb{P}(A))$ using the sub-double category $\mathbb{P}^{(2)}(A)$ of definition 197 and the 2-category of globs construction of observation 78 and simply observe that the 2-categories $\mathbb{P}(\mathbb{P}(A))^r$ and $\text{Glob}(\mathbb{P}^{(2)}(A))^r$ are identical (as substructures of $\mathbb{P}^2(A)^r$).

Lemma 200. *We may construct a functor $\mathbb{P}^*: \underline{\text{Cs-Cat}} \longrightarrow \underline{2\text{-Cat}}(\underline{\text{Cs}})$ and a family of isomorphisms $\mathbb{P}^*(\mathbb{P}(A)) \cong \text{Glob}(\mathbb{P}^{(2)}(A))$ that is natural in the complicial set A . Furthermore, applying the double category of pasting squares construction (cf. observation 76) we obtain the natural isomorphism depicted in the following diagram:*

$$\begin{array}{ccc}
 \underline{\text{Cs}} & \xrightarrow{\quad \mathbb{P} \quad} & \underline{\text{Cs-Cat}} \\
 \mathbb{P}^{(2)} \downarrow & \xRightarrow[\phi]{\cong} & \downarrow \mathbb{P}^* \\
 \underline{\text{Double}}(\underline{\text{Cs}}) & \xleftarrow[\text{Sq}]{} & \underline{2\text{-Cat}}(\underline{\text{Cs}})
 \end{array} \tag{55}$$

Proof. Taking our lead from the last observation, we define $\mathbb{P}^*: \underline{\text{Cs-Cat}} \longrightarrow \underline{2\text{-Cat}}(\underline{\text{Cs}})$ to be the functor which takes a complicial category $\mathbb{C}$, maps it to the double category $\mathbb{P}(\mathbb{C})$, applies the reflection dual to obtain a 2-category $\mathbb{P}(\mathbb{C})^r$ **and then** applies the 2-categorists 2-cellular dual to obtain $\mathbb{P}^*(\mathbb{C}) \stackrel{\text{def}}{=} (\mathbb{P}(\mathbb{C})^r)^{\text{co}}$. Our reason for applying this final duality will make itself clear momentarily, for now it is enough to observe that this construction, as the composite of three functorial constructions, is itself functorial.

Everything else follows from the fact that $(\mathbb{P}^{(2)}(A), t\mathbb{P}^{(2)}(A))$ is a complicial double category with connection for each complicial set A . In particular, observation 82 provides us with a natural isomorphism $\text{Glob}(\mathbb{P}^{(2)}(A))^r \cong \text{Glob}(\mathbb{P}^{(2)}(A))^{\text{co}}$ and observation 199 shows that its left hand side is actually identical to $\mathbb{P}(\mathbb{P}(A))^r$. So applying the 2-cellular dual $(-)^{\text{co}}$, and using the fact that it is involutive, we get the first of the natural isomorphisms asked for in the statement of this lemma $\mathbb{P}^*(\mathbb{P}(A)) \stackrel{\text{def}}{=} (\mathbb{P}(\mathbb{P}(A))^r)^{\text{co}} \cong \text{Glob}(\mathbb{P}^{(2)}(A))$, which explains why we defined $\mathbb{P}^*$ as we did.

Applying the pasting squares construction $\text{Sq}: \underline{2\text{-Cat}}(\underline{\text{Cs}}) \longrightarrow \underline{\text{Double}}(\underline{\text{Cs}})$ to this natural isomorphism we get $\text{Sq}(\mathbb{P}^*(\mathbb{P}(A))) \cong \text{Sq}(\text{Glob}(\mathbb{P}^{(2)}(A)))$ and then lemma 81 provides another isomorphism $\text{Sq}(\text{Glob}(\mathbb{P}^{(2)}(A))) \cong \mathbb{P}^{(2)}(A)$ which is also natural in A . Finally, composing the two we get the natural isomorphism ϕ asked for in display (55) of the statement. $\square$

Observation 201 (a more concrete description of $\phi: \mathbb{P}^{(2)} \longrightarrow \text{Sq} \circ \mathbb{P}^* \circ \mathbb{P}$). Tracing through the construction of lemma 81 and those of the last few observations, it is clear that the double categories $\mathbb{P}^{(2)}(A)$ and $\text{Sq}(\mathbb{P}^*(\mathbb{P}(A)))$ as we've constructed them share precisely the same categories of horizontal and vertical arrows, namely $\mathbb{P}(A)$, and that the double functor $\phi_A: \mathbb{P}^{(2)}(A) \longrightarrow \text{Sq}(\mathbb{P}^*(\mathbb{P}(A)))$ acts trivially on these. In other words, it carries each square

λ in $P^{(2)}(A)$ to a pasting square of the form:

$$\begin{array}{ccc}
 \bullet & \xrightarrow{s_v(\lambda)} & \bullet \\
 s_h(\lambda) \downarrow & \lambda^* \Downarrow & \downarrow t_h(\lambda) \\
 \bullet & \xrightarrow{t_v(\lambda)} & \bullet
 \end{array}$$

Returning again to lemma 81, we see that the 2-cell λ^* is constructed by composing λ above and below with compatible connection squares in $P^{(2)}(A)$. By observation 199 this vertical composite $\nu_{t_v(\lambda)} * \lambda * \mu_{s_v(\lambda)}$ is actually in the sub-double category $\mathbb{P}(P(A))$ and is easily seen to provide a 2-cell with the appropriate orientation in the dual 2-category $\mathbb{P}^*(P(A))$.

Observation 202. Given the result of lemma 200 and the discussion in observation 198, the result alluded to in observation 196 is now a matter of mere formality. To illuminate that comment, define the functor $\mathbb{I}: \underline{\text{Cs-Cat}} \longrightarrow \underline{\text{Double}}(\underline{\text{Cs}})$ to be the composite

$$\underline{\text{Cs-Cat}} \xrightarrow{\mathbb{P}^*} \underline{2\text{-Cat}}(\underline{\text{Cs}}) \xrightarrow{\text{Sq}} \underline{\text{Double}}(\underline{\text{Cs}}) \xrightarrow{E_h} \underline{\text{Double}}(\underline{\text{Cs}})$$

and observe that on applying the functor E_h to the natural isomorphism ϕ of lemma 200, and appealing to the equality $P(\mathbb{P}(A)) = E_h(P^{(2)}(A))$ of observation 198, we get a natural isomorphism:

$$\begin{array}{ccc}
 \underline{\text{Cs}} & \xrightarrow{\mathbb{P}} & \underline{\text{Cat}}(\underline{\text{Cs}}) \\
 P \downarrow & \cong \Downarrow \theta & \downarrow P \\
 \underline{\text{Cs-Cat}} & \xrightarrow{\mathbb{I}} & \underline{\text{Double}}(\underline{\text{Cs}})
 \end{array} \tag{56}$$

Examining the composite at the top right of this diagram we see, by observation 198, that if A is a complicial set then the double category $P(\mathbb{P}(A))$ has horizontal categories of arrows $P(A)$ and squares $P(\Delta[1] \Rightarrow A)$ and that its vertical source and target maps are $P(\Delta(\varepsilon_0^1) \Rightarrow A)$ and $P(\Delta(\varepsilon_1^1) \Rightarrow A)$ respectively. As also observed there, these are simply the various structures which appear on the right of the triangles in display (54) of observation 196.

Taking this as our lead, we turn to the lower left composite of display (56). Consulting observation 196, it is clear that if $\mathbb{C}$ is a complicially enriched category then the natural candidate for $I(\mathbb{C}) \in \underline{\text{Cs-Cat}}$ is the horizontal category of squares of $\mathbb{I}(\mathbb{C})$, which we know to be complicially enriched (cf. observation 198). Furthermore, by construction, the horizontal category of arrows of $\mathbb{I}(\mathbb{C})$ is actually $\mathbb{C}$ itself, and so the choice $I(\mathbb{C}) \stackrel{\text{def}}{=} \text{sq}_h(\mathbb{I}(\mathbb{C}))$ also presents us with obvious candidates for the complicial functors $\partial_{\mathbb{C}}^0, \partial_{\mathbb{C}}^1: I(\mathbb{C}) \longrightarrow \mathbb{C}$ which we can take to be the vertical source and target maps of $\mathbb{I}(\mathbb{C})$ respectively. Now suppose that $f: \mathbb{C} \longrightarrow \mathbb{C}'$ is a functor in $\underline{\text{Cs-Cat}}$ then the double functor $\mathbb{I}(f): \mathbb{I}(\mathbb{C}) \longrightarrow \mathbb{I}(\mathbb{C}')$ acts like f on horizontal categories of arrows and so if we define $I(f)$ to be its action on horizontal categories of squares then the naturality conditions for $\partial_{\mathbb{C}}^0$ and $\partial_{\mathbb{C}}^1$ simply reduce to the fact that the double functor $\mathbb{I}(f)$ preserves vertical sources and targets.

Finally, having made this choice for I and the associated natural transformations ∂^0 and ∂^1 , the isomorphism θ of display (56) provides us with a natural family of isomorphisms

$$P(\Delta[1] \Rightarrow A) = \text{sq}_h(P(\mathbb{P}(A))) \xrightarrow[\cong]{\theta_A} \text{sq}_h(\mathbb{I}(P(A))) = I(P(A))$$

as required by display (53) of observation 196. Furthermore, each θ_A is actually a double functor so, in particular, the fact that it preserves vertical sources and targets, when combined with our identification of the vertical sources and targets of $P(\mathbb{P}(A))$, the definitions of $\partial_{P(A)}^0$ and $\partial_{P(A)}^1$ and the observation that θ acts as the identity on horizontal categories of arrows (cf. observation 201), establishes the triangle identities of display (54). $\square$

Observation 203 (a more explicit description of $I(\mathbb{C})$). If $\mathbb{C}$ is a complicially enriched category then $I(\mathbb{C})$ has $|\mathbb{C}_a|_0$ as its complicial set of objects and n -arrows which are pasting squares of the form

$$\begin{array}{ccc} \bullet & \xrightarrow{q} & \bullet \\ p \downarrow & \Downarrow \varphi & \downarrow p' \\ \bullet & \xrightarrow{q'} & \bullet \end{array} \quad (57)$$

where p and p' are n -arrows in $|\mathbb{C}|_0 \subseteq_r \mathbb{C}$ (which are its source and target object in $I(\mathbb{C})_o = |\mathbb{C}_a|_0$), q and q' are n -arrows in $\mathbb{C}$ and φ is an n -dimensional 2-cell in $\mathbb{P}^*(\mathbb{C})$ (which has $\mathbb{C}$ as its underlying category of 1-cells). Such pasting squares are thin in $I(\mathbb{C})_a$ precisely when q , q' and φ are all thin in their respective complicial sets, simplicial operators act on them point-wise and they compose by horizontal pasting. Furthermore the functors $\partial_{\mathbb{C}}^0$ and $\partial_{\mathbb{C}}^1$ simply project a pasting square to its vertical source q and target q' respectively.

10.3. A Decalage Construction on Complicially Enriched Categories. Finally we may transfer the decalage construction of subsection 10.1 to $\underline{\text{Cs-Cat}}$ using the endo-functor I which we constructed in observation 202, as follows:

Definition 204 (decalage on $\underline{\text{Cs-Cat}}$). If $\mathbb{C}$ is a complicially enriched category then define $\text{Dec}(\mathbb{C})$, its decalage, to be the complicially enriched category obtained by taking the pullback

$$\begin{array}{ccc} \text{Dec}(\mathbb{C}) & \xrightarrow{\subseteq_r} & I(\mathbb{C}) \\ \downarrow & \lrcorner & \downarrow \partial_{\mathbb{C}}^0 \\ |\mathbb{C}|_0 & \xrightarrow[\subseteq_r]{} & \mathbb{C} \end{array} \quad (58)$$

in $\underline{\text{Cs-Cat}}$. As a functor, this construction is the point-wise pullback, in the functor category $[\underline{\text{Cs-Cat}}, \underline{\text{Cs-Cat}}]$, of the natural inclusion $|\cdot|_0 \xrightarrow{\subseteq_r} \text{id}_{\underline{\text{Cs-Cat}}}$ along the natural transformation $\partial^0: I \longrightarrow \text{id}_{\underline{\text{Cs-Cat}}}$. We make Dec into a copointed endo-functor on $\underline{\text{Cs-Cat}}$ by defining the counit $\partial: \text{Dec} \longrightarrow \text{id}_{\underline{\text{Cs-Cat}}}$ to be the natural transformation obtained by composing the natural inclusion $\text{Dec} \xrightarrow{\subseteq_r} I$ with the natural transformation $\partial^1: I \longrightarrow \text{id}_{\underline{\text{Cs-Cat}}}$.

Observation 205 (an explicit description of Dec on $\underline{\text{Cs-Cat}}$). Applying the explicit description of $I(\mathbb{C})$ given in observation 203 and the definition of $|\mathbb{C}|_0$, we see that $\text{Dec}(\mathbb{C})$ is the complicial subcategory of $I(\mathbb{C})$ consisting of those pasting squares of the form depicted in

display (57) for which the 1-cell q is an identity n -arrow in $\mathbb{C}$. In other words, contracting the 1-cell q in that diagram to a point, we may depict the n -arrows of $\text{Dec}(\mathbb{C})$ as triangles:

$$\begin{array}{ccc} & \bullet & \\ p \swarrow & \varphi & \searrow p' \\ \bullet & \xrightarrow{q'} & \bullet \end{array} \quad (59)$$

With respect to this description, the component $\partial_{\mathbb{C}} : \text{Dec}(\mathbb{C}) \longrightarrow \mathbb{C}$ of the counit simply maps the n -arrow depicted in (59) to the n -arrow q' in $\mathbb{C}$.

We may easily establish a close relationship between our decalage constructions on $\underline{\mathbb{C}s}$ and $\underline{\mathbb{C}s}\text{-}\underline{\mathbb{C}at}$:

Theorem 206. *The natural isomorphism θ of observation 196, as constructed in observation 202, restricts to give rise to a natural isomorphism*

$$\begin{array}{ccc} \underline{\mathbb{C}s} & \xrightarrow{\text{Dec}} & \underline{\mathbb{C}s} \\ P \downarrow & \cong \Downarrow \psi & \downarrow P \\ \underline{\mathbb{C}s}\text{-}\underline{\mathbb{C}at} & \xrightarrow{\text{Dec}} & \underline{\mathbb{C}s}\text{-}\underline{\mathbb{C}at} \end{array} \quad (60)$$

which makes the pair (P, ψ) into a strong transformation of copointed endo-functors from the decalage construction on $\underline{\mathbb{C}s}$ to that on $\underline{\mathbb{C}s}\text{-}\underline{\mathbb{C}at}$.

Proof. This is simple calculation with pullbacks, which is best summarised in the following cubical diagram:

$$\begin{array}{ccccc} P(\text{Dec}(A)) \hookrightarrow & \xrightarrow{P(\subseteq_r)} & P(\Delta[1] \Rightarrow A) & \xrightarrow{P(\Delta(\varepsilon_1^1) \Rightarrow A)} & P(A) \\ \downarrow \lrcorner \cong \nearrow \psi_A & & \downarrow \lrcorner \cong \nearrow \theta_A & & \downarrow \partial_{P(A)}^1 \\ \text{Dec}(P(A)) \hookrightarrow & \xrightarrow{\subseteq_r} & I(P(A)) & & \\ \downarrow \lrcorner & & \downarrow P(\Delta(\varepsilon_0^1) \Rightarrow A) & & \downarrow \partial_{P(A)}^0 \\ P(|A|_0) \hookrightarrow & \xrightarrow{\subseteq_r} & P(A) & & \\ \downarrow \cong & & \downarrow \cong & & \\ |P(A)|_0 \hookrightarrow & \xrightarrow{\subseteq_r} & P(A) & & \end{array}$$

Here the front face is simply the pullback used to define $\text{Dec}(P(A))$ in display (60) of definition 206, the right-hand face is the first of the commuting triangles in display (54) of observation 196 and the equality in the lower left corner is that supplied by lemma 184. The back face is also a pullback since it is obtained by applying the left exact functor $P : \underline{\mathbb{C}s} \longrightarrow \underline{\mathbb{C}s}\text{-}\underline{\mathbb{C}at}$ to the pullback used to define $\text{Dec}(A)$ in display (51) of definition 190.

It follows therefore, by the pullback property of the front and back faces, that we get the induced isomorphism $\psi_A : P(\text{Dec}(A)) \longrightarrow \text{Dec}(P(A))$ which is uniquely determined by the property that it makes the upper face commute. Furthermore, it is clear that the naturality of θ_A immediately implies that ψ_A is also natural in the complicial set A .

Finally, consulting the definitions of the copointings of these decalage constructions given in definitions 190 and 206, we see that we may combine the upper face of our cube with the

triangle to its right, which is the second of the triangles from display (54) of observation 196, to demonstrate that ψ_A satisfies the compatibility condition required of a transformation of copointed endo-functors. $\square$

10.4. Semi-Simplicial Reconstruction. Using the decalage constructions of the last few subsections we may now apply the semi-simplicial version of lemma 93 to prove an exceedingly important result by which we may re-construct the underlying semi-simplicial structure of a complicial set A from the complicially enriched category $P(A)$. To do this, we must first introduce a suitable connected components functor on the category of complicially enriched categories.

Definition 207 (connected components of complicially enriched categories). If $\mathbb{C}$ is a complicially enriched category then let $\Pi_0(\mathbb{C})$, its *set of connected components*, be the set obtained by forming the coequaliser

$$(\mathbb{C}_a)_0 \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} (\mathbb{C}_o)_0 \xrightarrow{q_{\mathbb{C}}} \Pi_0(\mathbb{C})$$

in Set. In other words, we start with the set of objects of $\mathbb{C}$ and form $\Pi_0(\mathbb{C})$ from it by identifying any pair of objects which occurs as the source and target of the same 1-arrow in $\mathbb{C}$.

If $f: \mathbb{C} \longrightarrow \mathbb{C}'$ is a complicial functor then we know, by definition 68, that it preserves sources and targets, in other words its underlying maps $f: \mathbb{C}_a \longrightarrow \mathbb{C}'_a$ and $f_o: \mathbb{C}_o \longrightarrow \mathbb{C}'_o$ satisfy the equalities $f_o \circ s = s \circ f$ and $f_o \circ t = t \circ f$ and therefore that they induce a unique function $\Pi_0(f): \Pi_0(\mathbb{C}) \longrightarrow \Pi_0(\mathbb{C}')$ satisfying the equation $\Pi_0(f)(q_{\mathbb{C}}(c)) = q_{\mathbb{C}'}(f_o(c))$ for each $c \in (\mathbb{C}_o)_0$. This action on complicial functors clearly makes Π_0 into a functor $\Pi_0: \underline{\mathbf{Cs-Cat}} \longrightarrow \underline{\mathbf{Set}}$.

Observation 208 (relating Π_0 on $\underline{\mathbf{Cs}}$ and $\underline{\mathbf{Cs-Cat}}$). We would like to use our connected components functors Π_0 on $\underline{\mathbf{Cs}}$ and $\underline{\mathbf{Cs-Cat}}$ in applying (the semi-simplicial version of) lemma 93 to the strong transformation of copointed decalage functors provided by theorem 206. However, in order to do so we need to be able to demonstrate that we have a family of isomorphisms $\Pi_0(P(A)) \cong \Pi_0(A)$ which is natural in the complicial set A . These may be constructed with the aid of the following diagram

$$\begin{array}{ccccc} (P(A)_a)_0 & \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} & (P(A)_o)_0 & \xrightarrow{q_{P(A)}} & \Pi_0(P(A)) \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ A_1 & \begin{array}{c} \xrightarrow{A_{\varepsilon_0^1}} \\ \xrightarrow{A_{\varepsilon_1^1}} \end{array} & A_0 & \xrightarrow{q_A} & \Pi_0(A) \end{array}$$

in which the upper and lower horizontal “forks” are the defining coequalisers for $\Pi_0(P(A))$ (definition 207) and $\Pi_0(A)$ (definition 111) respectively.

The vertical isomorphism in the centre is, in fact, an equality since we know that the complicial set of objects $P(A)_o$ is $|A|_0 \subseteq_r A$ and that $|A|_0$ and A have the same sets of 0-simplices so it follows that $(P(A)_o)_0 = A_0$. To construct the vertical isomorphism to the left, we start by observing that every 0-arrow of $P(A)$ is (trivially) in its subcategory $P(A)$ and so, by observation 177, we know that the elements of $(P(A)_a)_0$ bear a canonical representation as stratified maps $p: \Delta[1] \cong \Delta[1] \otimes \Delta[0] \longrightarrow A$. In other words, $(P(A)_a)_0$ may be identified

with the homset $\underline{\text{Strat}}(\Delta[1], A)$ and the vertical arrow on the left of our diagram is simply Yoneda's isomorphism. It is now a routine matter to check that the left-hand side of our diagram commutes serially (cf. [22]) and it follows that the coequaliser property of our forks ensures that we get the induced isomorphism indicated as a dotted vertical arrow to the right of the diagram. The naturality of this family of isomorphisms in A is now a trivial consequence of their definition.

Theorem 209 (semi-simplicial reconstruction). *Let $R: \underline{\text{Cs-Cat}} \longrightarrow \underline{\text{SSimp}}$ denote the semi-simplicial nerve functor N_{Dec, Π_0} associated with the decalage construction on $\underline{\text{Cs-Cat}}$ (of definition 204) under the connected components functor $\Pi_0: \underline{\text{Cs-Cat}} \longrightarrow \text{Set}$ (of definition 207). Then we have a natural isomorphism*

$$\begin{array}{ccc} \underline{\text{Cs}} & \xrightarrow{P} & \underline{\text{Cs-Cat}} \\ & \searrow U \quad \cong \quad \swarrow R & \\ & \underline{\text{SSimp}} & \end{array} \quad (61)$$

where U on the left-hand side is simply the forgetful functor from $\underline{\text{Cs}}$ to $\underline{\text{SSimp}}$. Consequently, we often refer to R as the **semi-simplicial reconstruction functor**.

Proof. Observation 208 and theorem 206 provide us with the data which allows us to apply the semi-simplicial version of lemma 93 to construct a natural isomorphism:

$$\begin{array}{ccc} \underline{\text{Cs}} & \xrightarrow{P} & \underline{\text{Cs-Cat}} \\ & \searrow N_{\text{Dec}, \Pi_0} \quad \cong \quad \swarrow N_{\text{Dec}, \Pi_0} & \\ & \underline{\text{SSimp}} & \end{array}$$

However, by definition the functor on the right-hand diagonal is simply our semi-simplicial reconstruction functor $R: \underline{\text{Cs-Cat}} \longrightarrow \underline{\text{SSimp}}$ and by observation 191 we know that to one on the left-hand diagonal is (isomorphic to) the forgetful functor $U: \underline{\text{Cs}} \longrightarrow \underline{\text{SSimp}}$. In other words, the natural isomorphism thus constructed may be composed with the isomorphism $U \cong N_{\text{Dec}, \Pi_0}: \underline{\text{Cs}} \longrightarrow \underline{\text{SSimp}}$ to provide the natural isomorphism of display (61) in the statement of the theorem as required. $\square$

We now proceed to establish some important properties of semi-simplicial reconstruction:

Lemma 210. *The decalage functor $\text{Dec}: \underline{\text{Cs-Cat}} \longrightarrow \underline{\text{Cs-Cat}}$ preserves all (small) limits. Furthermore, its composite with the connected components functor $\Pi_0 \circ \text{Dec}: \underline{\text{Cs-Cat}} \longrightarrow \text{Set}$ simply maps each complicially enriched category $\mathbb{C}$ to a set which is naturally isomorphic to its set of objects $(\mathbb{C}_o)_0$ and so this functor also preserves all (small) limits.*

It follows, therefore, that the semi-simplicial reconstruction functor $R: \underline{\text{Cs-Cat}} \longrightarrow \underline{\text{SSimp}}$ also preserves all limits and that, as a consequence, it preserves monomorphisms.

Proof. The limit of a diagram in $\underline{\text{Cs-Cat}}$ is formed by forgetting the category structures on its vertices, calculating the limit of the resulting diagram in $\underline{\text{Cs}}$ (which is a reflective full subcategory of $\underline{\text{Strat}}$ and thus closed in there under all limits) and then defining the category structure on the consequent complicial set “point-wise”. It is therefore a matter of routine verification, using the explicit description of $\text{Dec}(\mathbb{C})$ given in observations 203 and 205, to show that Dec preserves limits that are constructed in this way.

To prove the second of these results, we fix a complicially enriched category $\mathbb{C}$ and exploit the explicit description of $\text{Dec}(\mathbb{C})$ given in observation 205 to construct the following diagram of sets and functions:

$$\begin{array}{ccccc}
 & \xleftarrow{g} & & \xleftarrow{f} & \\
 (\text{Dec}(\mathbb{C})_a)_0 & \xrightarrow{s} & (\text{Dec}(\mathbb{C})_o)_0 & \xrightarrow{q} & (\mathbb{C}_o)_0 \\
 & \xrightarrow{t} & & &
 \end{array} \quad (62)$$

Here the maps s and t are the source and target maps of $\text{Dec}(\mathbb{C})$ which simply take an arrow depicted as a triangle in display (59) to the 0-arrows p and p' respectively. To construct q , f and g , notice first that, by construction, we have $\text{Dec}(\mathbb{C})_o = |\mathbb{C}_a|_0$ and we know that $|\mathbb{C}_a|_0$ and $\mathbb{C}_a$ have the same sets of 0-simplices so we have that $(\text{Dec}(\mathbb{C})_o)_0 = (\mathbb{C}_a)_0$. It follows that we may take q and f to be the source and identity functions of the category $\mathbb{C}$ respectively. Finally, the action of g is best described in diagrammatic terms:

$$\begin{array}{ccc}
 (\text{Dec}(\mathbb{C})_o)_0 & \xrightarrow{g} & (\text{Dec}(\mathbb{C})_a)_0 \\
 \bullet \downarrow p & \xrightarrow{\quad} & \begin{array}{c} \bullet \\ \text{id} \parallel p \\ \bullet \end{array} \\
 \bullet & & \bullet \xrightarrow{p} \bullet
 \end{array}$$

It is now a routine matter to check that in the following diagram

$$\begin{array}{ccccc}
 (\text{Dec}(\mathbb{C})_o)_0 & \xrightarrow{g} & (\text{Dec}(\mathbb{C})_a)_0 & \xrightarrow{t} & (\text{Dec}(\mathbb{C})_o)_0 \\
 q \downarrow & & s \downarrow & & q \downarrow \\
 (\mathbb{C}_o)_0 & \xrightarrow{f} & (\text{Dec}(\mathbb{C})_o)_0 & \xrightarrow{q} & (\mathbb{C}_o)_0
 \end{array}$$

both squares commute and that its upper and lower horizontal composites are identities. Consequently display (62) is a split coequaliser and, comparing it with the coequaliser used to define Π_0 in definition 207, we see that it demonstrates that $(\mathbb{C}_o)_0$ is isomorphic to $\Pi_0(\text{Dec}(\mathbb{C}))$ as postulated. The naturality of this isomorphism follows easily as does the stated result regarding preservation of limits (since limits of complicially enriched categories are constructed point-wise in $\underline{\mathbf{Cs}}$).

Finally, returning to the construction of $R: \underline{\mathbf{Cs-Cat}} \longrightarrow \underline{\mathbf{SSimp}}$, as exposed in observation 86 and definition 87, we see that for each $n \in \mathbb{N}$ the functor $R(-)_n$ which applies R and then extracts the set of n -simplices of the result is, by definition, equal to $(\Pi_0 \circ \text{Dec}) \circ \text{Dec}^n$. However, by the two preservation results we have already established it follows that this latter functor preserves all limits. Therefore, quantifying over $n \in \mathbb{N}$ and appealing to the fact that limits are constructed point-wise (dimension by dimension) in $\underline{\mathbf{SSimp}}$, we see that R also preserves all limits.

Finally the comment regarding preservation of monomorphisms by R is a standard categorical result which applies to all pullback preserving functors (see [22]). $\square$

Theorem 211. *The semi-simplicial reconstruction functor $R: \underline{\mathbf{Cs-Cat}} \longrightarrow \underline{\mathbf{SSimp}}$ is faithful.*

Proof. This result hinges on the observation that if $\mathbb{C}$ is a complicially enriched category then the path category $P(\mathbb{C}_a)$ constructed from its complicial set of arrows is actually (isomorphic

to) a regular subcategory of the complicially enriched decalage category $\text{Dec}(\mathbb{C})$. In particular it is easily seen, directly from the definition of $\mathbb{P}^*(\mathbb{C})$ in lemma 200 and the explicit description of $\text{Dec}(\mathbb{C})$ given in observations 203 and 205, that $\text{P}(\mathbb{C}_a)$ is naturally isomorphic to the subcategory of $\text{Dec}(\mathbb{C})$ of those triangles of the form depicted in display (59) for which q' is actually an identity arrow in $\mathbb{C}$.

So if $\mathbb{C}$ is a complicially enriched category then we may apply the semi-simplicial reconstruction functor R to the regular inclusion $\text{P}(\mathbb{C}_a) \hookrightarrow \text{Dec}(\mathbb{C})$ and appeal to the monomorphism preservation result of theorem 210 to demonstrate that we thereby obtain a monomorphism $R(\text{P}(\mathbb{C}_a)) \hookrightarrow R(\text{Dec}(\mathbb{C}))$ in SSimp. Now, by theorem 209 we have an isomorphism $U(\mathbb{C}_a) \cong R(\text{P}(\mathbb{C}_a))$ and furthermore, on consulting the construction of R as described in observation 86 and definition 87, we easily see that we also have $R(\text{Dec}(\mathbb{C})) = \text{Dec}(R(\mathbb{C}))$, where the functor Dec on the right-hand side is simply the usual decalage functor on SSimp. Of course all of these inclusions, isomorphisms and equalities are natural in the complicially enriched category $\mathbb{C}$ and on composing them we get a family of monomorphisms $m_{\mathbb{C}_a} : U(\mathbb{C}_a) \hookrightarrow \text{Dec}(R(\mathbb{C}))$ in SSimp which is also natural in $\mathbb{C} \in \text{Cs-Cat}$.

We now come to the crux of our argument, so suppose that $f, g : \mathbb{C} \rightarrow \mathbb{C}'$ are two functors in Cs-Cat and observe that the naturality of the family of monomorphisms constructed in the previous paragraph provides us with a serially commutative diagram:

$$\begin{array}{ccc} U(\mathbb{C}_a) & \xhookrightarrow{m_{\mathbb{C}}} & \text{Dec}(R(\mathbb{C})) \\ \begin{array}{c} U(f) \downarrow \\ U(g) \downarrow \end{array} & & \begin{array}{c} \text{Dec}(R(f)) \downarrow \\ \text{Dec}(R(g)) \downarrow \end{array} \\ U(\mathbb{C}'_a) & \xhookrightarrow{m_{\mathbb{C}'}} & \text{Dec}(R(\mathbb{C}')) \end{array}$$

So to prove that R is faithful suppose that $R(f) = R(g)$ and observe that then the vertical arrows on the right of this square are equal and so it follows, by serial commutativity, that the composites $m_{\mathbb{C}'} \circ U(f)$ and $m_{\mathbb{C}'} \circ U(g)$ in the lower left of the square must also be equal. However $m_{\mathbb{C}'}$ is a monomorphism so it follows that $U(f) = U(g)$ and we know that $U : \text{Cs} \rightarrow \text{SSimp}$ is faithful, since it only forgets about degeneracy actions and thinness without discarding any simplices, so it follows therefore that f and g coincide as maps on complicial sets of arrows and thus are identical functors as required. $\square$

Finally we may apply this last result to establish the theorem toward which we have been working throughout this entire section. Later on we shall see that this result leads directly to a proof that the ω -categorical nerve functor constructed by Street in [31] actually provides us with an equivalence between the categories of ω -categories and complicial sets. However, before stating and proving this pivotal result we pause to recall a simple categorical result which will be used in its proof and again in the next section:

Lemma 212. *Suppose that $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{E}$ is a composable pair of functors then*

- (i) *if the composite $G \circ F$ is fully faithful and G itself is faithful then F is also fully faithful, and*
- (ii) *if the composite $G \circ F$ is an equivalence and G itself is fully faithful then F is also an equivalence.*

Proof. This is an entirely trivial categorical result which we leave up to the reader to verify. $\square$

Theorem 213. *The path category functor $P: \underline{\mathbf{Cs}} \longrightarrow \underline{\mathbf{Cs-Cat}}$ is fully faithful.*

Proof. To prove this result we need to work a little harder to reconstruct the thinness and degeneracy information built into a complicial set A from its path category $P(A)$. To do so we'll exploit the fully faithful representation of well-tempered stratified sets as filtered semi-simplicial sets which we studied in lemma 119. Our argument can be summed up in the following diagram

$$\begin{array}{ccc}
 \underline{\mathbf{Cs}} & \xrightarrow{P} & \underline{\mathbf{Cs-Cat}} \\
 \downarrow |\cdot|_{\bullet} & \begin{array}{c} = \\ \downarrow [\mathbb{N}, P] \end{array} & \downarrow |\cdot|_{\bullet} \\
 [\mathbb{N}, \underline{\mathbf{Cs}}] & \xrightarrow{[\mathbb{N}, P]} & [\mathbb{N}, \underline{\mathbf{Cs-Cat}}] \\
 \searrow [\mathbb{N}, U] & \cong & \swarrow [\mathbb{N}, R] \\
 & [\mathbb{N}, \underline{\mathbf{SSimp}}] &
 \end{array}$$

in which the lower triangle is obtained by applying the 2-functor $[\mathbb{N}, -]$ to the triangle in display (61) of theorem 209. The left-hand vertical functor labelled $|\cdot|_{\bullet}$ maps a complicial set A to its filtered family of complicial superstructures $|A|_0 \subseteq_r |A|_1 \subseteq_r \dots \subseteq |A|_n \subseteq_r \dots$. It follows that the composite of the functors down the left hand side is simply the fully faithful representation of lemma 119 restricted to the category $\underline{\mathbf{Cs}}$ which is a subcategory of $\underline{\mathbf{Strat}}_W$ by lemma 163.

The right hand vertical, also called $|\cdot|_{\bullet}$, maps a complicially enriched category $\mathbb{C}$ to the filtered family of superstructures $|\mathbb{C}|_0 \subseteq_r |\mathbb{C}|_1 \subseteq_r \dots \subseteq_r |\mathbb{C}|_n \subseteq_r \dots$ in $\underline{\mathbf{Cs-Cat}}$ as defined in observation 182. Now we may immediately recast the result of lemma 184, which showed that $P(|A|_n) = |P(A)|_n$ for each $n \in \mathbb{N}$ and $A \in \underline{\mathbf{Cs}}$, to demonstrate the commutativity of the upper square in our diagram.

Furthermore, suppose that we are given two functors $f, g: \mathbb{C} \longrightarrow \mathbb{C}'$ in $\underline{\mathbf{Cs-Cat}}$ then, by definition, we have $|f|_{\bullet} = |g|_{\bullet}$ if and only if f and g agree when restricted to the superstructure $|\mathbb{C}|_n \subseteq_r \mathbb{C}$ for each $n \in \mathbb{N}$. However it is clear that any complicially enriched category $\mathbb{C}$ is equal to the union of its superstructures and so it follows that f and g must be equal as functors on $\mathbb{C}$. In other words, the functor $|\cdot|_{\bullet}$ on the right-hand side of our diagram is faithful as is the diagonal functor $[\mathbb{N}, R]$ on that same side, since it acts by applying the faithful semi-simplicial reconstruction functor R point-wise to filtered complicial sets and maps.

Now observe that our diagram provides us with an isomorphism between the fully faithful composite $[\mathbb{N}, U] \circ |\cdot|_{\bullet}$ on its left-hand side and the composite $[\mathbb{N}, R] \circ |\cdot|_{\bullet} \circ P$ on its right-hand side, which is thus also fully faithful. Furthermore we have shown that $[\mathbb{N}, R] \circ |\cdot|_{\bullet}$ is a composite of faithful functors and is thus itself faithful, so finally applying lemma 212(i) we may infer that P is fully faithful as postulated. $\square$

11. STREET'S ω -CATEGORICAL NERVE CONSTRUCTION

11.1. Parity Complexes. Street's nerve construction [31] proceeds by first providing an explicit construction of the free ω -category whose generators are the faces of a simplex and whose relations are suitably re-interpreted and oriented versions of the face relations. Later he introduced structures called *parity complexes* [33] in order to generalise this work to encompass cubes and a range of other polytopes which may be obtained as products or joins of simplices and globs (oriented globes).

We will need to understand and calculate with these free, geometrically derived ω -categories in the sequel and so we briefly review the important parts of the theory of parity complexes. For a more detailed analysis of the combinatorics of these structures we refer the reader to [33] and [34].

Definition 214 (pre-parity complexes). A *pre-parity complex* is a graded set $C = \bigcup_{n=0}^{\infty} C_n$ that comes equipped with a pair of operations which map each element $x \in C_n$ (for $n > 0$) to disjoint non-empty finite subsets $x^-, x^+ \subseteq C_{n-1}$. We say that the elements of C_n are n -dimensional. In order to simplify some calculations and definitions we will generally adopt the convention that if x is a 0-dimensional element then we take x^- and x^+ to be the empty set.

Following Street we will generally call the elements of x^- and x^+ the *negative faces* and *positive faces* of x respectively. We also reserve the symbols φ and ξ to vary over the set of parity symbols $\{+, -\}$ and use the notation $\neg\varphi$ to denote the opposite parity to φ . We also say that an integer i is of parity ξ if i is even and ξ is the parity $+$ or i is odd and ξ is the parity $-$.

We consider any subset S of C to be graded according to the grading of C and define its n -superstructure by $|S|_n \stackrel{\text{def}}{=} \bigcup_{m=0}^n S_m$. We will also tend to use the notation S^{-n} to denote the subset obtained by omitting the n -dimensional elements from S (that is $S^{-n} = S \setminus S_n$).

We say that a subset D is a sub-pre-parity complex of C (denoted $D \subseteq_p C$) if it is closed in C under face operations, that is if φ is an arbitrary parity symbol then for all $s \in D$ we have $s^\varphi \subseteq D$. The family of sub-pre-parity complexes of C is closed under unions and intersections and it follows that each subset S of C is contained in a unique smallest sub-pre-parity complex D , called the pre-parity complex *generated* by S .

When manipulating pre-parity complexes we will often have use for functions f which map elements one pre-parity complex to *sets* of the elements of another (or possibly the same) pre-parity complex. We will depict such functions using the crossed arrow notation $f: C \rightrightarrows D$ and if $S \subseteq C$ then we'll take the notation $f(S)$ to mean the union $\bigcup_{x \in S} f(x)$. In particular, if ξ is a parity symbol then under this notational convention we have $S^\xi = \bigcup_{x \in S} x^\xi$ the set of those elements which are ξ -parity faces of some element of S .

Street introduces the notations $S^\mp \stackrel{\text{def}}{=} S^- \setminus S^+$ and $S^\pm \stackrel{\text{def}}{=} S^+ \setminus S^-$ for the sets of negative (resp. positive) faces of elements of S which are not positive (resp. negative) faces of any element of S . He also introduces a binary perpendicularity relation $S \perp T$ which holds when $(S^+ \cap T^+) \cup (S^- \cap T^-) = \emptyset$, that is to say when none of the ξ -parity faces of some simplex in T is also an ξ -parity face of some simplex in S .

If $x, y \in C$ then we say that $x \perp y$ when the corresponding singleton sets are perpendicular. We say that a subset $S \subseteq C$ is *well-formed* if it has at most one 0-dimensional element and if whenever we have $x, y \in S$ with $x \neq y$ then $x \perp y$.

Finally, for $x, y \in C$ we write $x < y$ if $x^+ \cap y^- \neq \emptyset$ and $x \prec y$ if $x \in y^-$ or $y \in x^+$ then for $S \subseteq C$ we define $\triangleleft_S$ and $\blacktriangleleft_S$ to be the pre-orders obtained as the reflexive transitive closures of $<$ and $\prec$ on S (respectively). Notice that $\triangleleft_S$ is a sub-order of $\blacktriangleleft_S$ and that if $T \subseteq S$ then $\triangleleft_T$ and $\blacktriangleleft_T$ are sub-orders of the restrictions of $\triangleleft_S$ and $\blacktriangleleft_S$ to T (respectively). We also adopt the assumption that whenever order properties of a subset $S \subseteq C$ are referred to that it will be implicitly understood that the order $\triangleleft_S$ is intended.

Definition 215 (movement). Suppose that S, M and P are subsets of a pre-parity complex C then we say that S moves M to P when we have

$$P = (M \cup S^+) \setminus S^- \text{ and } M = (P \cup S^-) \setminus S^+$$

and we denote this relationship by $S: M \longrightarrow P$. The movement concept is fundamental to Street's use of parity complexes in studying ω -categories. The reader may find a study of the basic properties of this concept in propositions 2.1-2.4 of section 2 of [33].

Definition 216 (parity complexes). In general, we will not be that interested here in the detailed combinatorics involved in the definition and theory of parity complexes. However, for the record we will recall that a *parity complex* is a pre-parity complex C which satisfies the following axioms:

Axiom 1 $x^{--} \cup x^{++} = x^{+-} \cup x^{-+}$,

Axiom 2 x^- and x^+ are both well formed,

Axiom 3(a) $x \triangleleft_C y \triangleleft_C x$ implies $x = y$, and

Axiom 3(b) $x \triangleleft_C y$, $x \in z^\varphi$ and $y \in z^\xi$ imply that $\varphi = \xi$.

In general, much of the theory of parity complexes follows from these axioms, however at various stages Street found that he needed to introduce auxiliary assumptions to make certain arguments work. Notably, in order to correct the proof of his “excision of extremals” result in [34] he needed to assume that the ordering $\blacktriangleleft_C$ on C was anti-symmetric and that each element $x \in C$ satisfied a certain *globularity condition*. Furthermore, to extend his results to products of parity complexes (see later) he needed to assume that each factor complex *and its odd dual* satisfied these conditions.

However, we will not concern ourselves with the details of these various conditions here. Suffice it to say that Street demonstrated that every parity complex we shall meet in the sequel satisfies every one of these conditions. Consequently, from here on we shall take it as understood that when we say that such and such is a parity complex we tacitly assume that it satisfies all of the technical conditions required to make Street's excision of extremals argument work.

Indeed all of the examples discussed in [33] satisfy a stronger totality condition with regard to the $\blacktriangleleft_C$ relation which is of interest in its own right, but we shall not pursue that point here.

Finally, notice that if D is a sub-pre-parity complex of the parity complex C then, since the orderings $\triangleleft_D$ and $\blacktriangleleft_D$ are sub-orders of (the restrictions of) $\triangleleft_C$ and $\blacktriangleleft_C$ (respectively), it is clear that D is also a parity complex.

Observation 217 (ω -categories from parity complexes). If C is a any graded set we may define a simple ω -category $\mathcal{N}(C)$ with:

- cells (M, P) where M and P are finite subsets of C ,

- n -source and n -target operations given by

$$\begin{aligned} s_n(M, P) &= (|M|_n, M_n \cup |P|_{n-1}) \\ t_n(M, P) &= (|M|_{n-1} \cup P_n, |P|_n) \end{aligned}$$

for $n \geq 0$, and

- n -composition of an n -compatible pair given by:

$$(M, P) *_n (N, Q) = (M \cup N^{\neg n}, P^{\neg n} \cup Q)$$

The idea here is that M_n and P_n are the sets consisting of the n -dimensional elements of C which we consider as comprising the surface elements of the n -dimensional source and target hemispheres of the cell (M, P) . Given this interpretation, the meaning of the source and target maps should be clear. We form composites by taking unions of the sets of faces and eliding the common n -dimensional face along which the cells we are composing agree. Notice that a pair (M, P) is an n -cell in $\mathcal{N}(C)$ if and only if $M_n = P_n$ and $M_m = P_m = \emptyset$ for all $m > n$.

While this is, of itself, a somewhat routine and uninspiring construction, its interest comes from Street's observation that the free ω -category (see definition 218 below) generated by the elements of the parity complex C may be obtained as a sub- ω -category $\mathcal{O}(C)$ of $\mathcal{N}(C)$. To be precise, we define $\mathcal{O}(C)$ to be the subset of $\mathcal{N}(C)$ of those pairs (M, P) for which:

- both of M and P are both well-formed, non-empty subsets of C , and
- each of the subsets M and P moves M to P , that is symbolically we have $M: M \longrightarrow P$ and $P: M \longrightarrow P$.

It is easily seen that $\mathcal{O}(C)$ is closed in $\mathcal{N}(C)$ under sources and targets, however much of the detailed argument in [33] is devoted to demonstrating that it is also closed under compositions in $\mathcal{N}(C)$ and is thus a sub- ω -category of it.

Before stating Street's result, it is worth noting that each n -dimensional element $x \in C_n$ gives rise to an inductively defined pair of subsets given by

$$\begin{aligned} \mu(x)_m &= \pi(x)_m = \emptyset & \text{for } m > n \\ \mu(x)_n &= \pi(x)_n = \{x\} \\ \mu(x)_m &= \mu(x)_{m+1}^{\mp} \text{ and } \pi(x)_m = \pi(x)_{m+1}^{\pm} & \text{for } n > m \geq 0 \end{aligned}$$

which provide for us a cell $\langle x \rangle \stackrel{\text{def}}{=} (\mu(x), \pi(x))$ in $\mathcal{N}(C)$. In some places we adopt the convention that we may modify our use of the symbols μ and π by superscripting them with parity symbols. In particular, if χ represents a symbol in the set $\{\mu, \pi\}$ then χ^+ represents the same symbol and χ^- represents the opposite one.

We say that x is *relevant* if this pair is in fact a cell in $\mathcal{O}(C)$ and one of the basic postulates of Street's theory, which we are tacitly assuming, is that every element of a parity complex C should be relevant. Cells of the form $\langle x \rangle$ are called *atoms*.

Definition 218 (freely generated ω -categories (Street [31])). If $\mathbb{C}$ is an ω -category and $G \subseteq \mathbb{C}$ is a set of its cells then we let $|G|_n \stackrel{\text{def}}{=} G \cap |\mathbb{C}|_n$ and we grade G by letting $G_0 = |G|_0$ and $G_{n+1} \stackrel{\text{def}}{=} |G|_{n+1} \setminus |\mathbb{C}|_n$ for each $n \in \mathbb{N}$.

We say that $\mathbb{C}$ is *weakly generated* by G when for each $n \in \mathbb{N}$ the set $|\mathbb{C}|_n \cup G_{n+1}$ separates ω -functors with domain $|\mathbb{C}|_{n+1}$, in the sense that if $f, g: |\mathbb{C}|_{n+1} \longrightarrow \mathbb{D}$ is a pair of ω -functors then we may infer that $f = g$ whenever $f(c) = g(c)$ for all $c \in |\mathbb{C}|_n \cup G_{n+1}$.

We say that $\mathbb{C}$ is *generated* by G when for each $n \in \mathbb{N}$ the smallest sub- ω -category of $\mathbb{C}$ which contains the set $|\mathbb{C}|_n \cup G_{n+1}$ is the superstructure $|\mathbb{C}|_{n+1}$ itself.

We say that $\mathbb{C}$ is *freely generated* by G when, for all ω -categories $\mathbb{D}$, for all $n \in \mathbb{N}$, for all ω -functors $f: |\mathbb{C}|_n \longrightarrow \mathbb{D}$ and for all functions $g: G_{n+1} \longrightarrow \mathbb{D}$ such that $s_n(g(c)) = f(s_n(c))$ and $t_n(g(c)) = f(t_n(c))$ for each $c \in G_{n+1}$, there exists a unique $(n+1)$ -functor $h: |\mathbb{C}|_{n+1} \longrightarrow \mathbb{D}$ whose restriction to $|\mathbb{C}|_n$ is f and whose restriction to G_{n+1} is g . Street depicts this extension property as a serially commutative diagram

$$\begin{array}{ccc}
 G_{n+1} & \xrightarrow{\quad g \quad} & \mathbb{D} \\
 \downarrow \text{hook} & \nearrow h & \downarrow s_n, t_n \\
 |\mathbb{C}|_{n+1} & & \mathbb{D} \\
 \downarrow s_n, t_n & & \downarrow s_n, t_n \\
 |\mathbb{C}|_n & \xrightarrow{\quad f \quad} & \mathbb{D}
 \end{array}$$

in which f is a ω -functor and h is its unique ω -functorial “lift” extending the action of the function g on the $(n+1)$ -dimensional generators in G .

Notice that none of these generation properties place any restriction on the set G_0 . However, it will be convenient in what follows to assume that each one includes the postulate that $G_0 = |\mathbb{C}|_0$.

It is clear that the free generation and generation both imply weak generation. A little less straightforwardly, we may also prove that if $G \subseteq \mathbb{C}$ freely generates $\mathbb{C}$ then it generates it. To prove this assume that the subset G freely generates $\mathbb{C}$ and let $\mathbb{D}$ be the smallest sub- ω -category of $\mathbb{C}$ which contains $|\mathbb{C}|_n \cup G_{n+1}$. Applying the free generation property, we may extend the inclusion ω -functor $|\mathbb{C}|_n \hookrightarrow \mathbb{D}$ using the subset inclusion function $G_{n+1} \hookrightarrow \mathbb{D}$ to give an ω -functor $h: |\mathbb{C}|_{n+1} \longrightarrow \mathbb{D}$ which, by definition, maps elements of $|\mathbb{C}|_n \cup G_{n+1}$ to themselves. Composing h with the inclusion $\mathbb{D} \subseteq |\mathbb{C}|_{n+1}$ and applying the separation property of G as a weak generator we see that the resulting ω -functor is equal to the identity on $|\mathbb{C}|_{n+1}$ and it follows that the inclusion $\mathbb{D} \subseteq |\mathbb{C}|_{n+1}$ must in fact be an equality as required.

Theorem 219 (Street [33] and [34] and originally [31]). *The set $\mathcal{O}(C)$ is a sub- ω -category of $\mathcal{N}(C)$ and, furthermore, it is the freely generated ω -category on the set of atoms $\langle C \rangle \stackrel{\text{def}}{=} \{\langle x \rangle \mid x \in C\}$.*

Proof. See loc. cit. □

The following corollary is a triviality, but never-the-less it provides us with a useful decomposition result for free ω -categories constructed from parity complexes:

Lemma 220. *If C is a parity complex and $C^{(i)} \subseteq_p C$ ($i \in I$) is a family of sub-parity complexes with $C = \bigcup_{i \in I} C^{(i)}$ then the diagram of sub- ω -category inclusions*

$$\begin{array}{ccc}
 & \mathcal{O}(C^{(i)} \cap C^{(j)}) & \\
 \swarrow & \downarrow & \searrow \\
 \mathcal{O}(C^{(i)}) & & \mathcal{O}(C^{(j)}) \\
 \swarrow & \downarrow & \searrow \\
 & \mathcal{O}(C) &
 \end{array}$$

displays $\mathcal{O}(C)$ as the wide pushout of its sub- ω -categories $\mathcal{O}(C^{(i)})$ in $\omega\text{-}\underline{\text{Cat}}$.

Proof (sketch). Suppose that we are given ω -functors $f_i: \mathcal{O}(C^{(i)}) \longrightarrow \mathbb{C}$ forming a cocone under this diagram and that we've constructed a functor $f: |\mathcal{O}(C)|_n \longrightarrow \mathbb{C}$ which coincides with f_i on $|\mathcal{O}(C^{(i)})|_n$ for each $i \in I$. If x is an $(n+1)$ -dimensional element of C then x is an element of some $C^{(i)}$ and we have $f(s_n(\langle x \rangle)) = f_i(s_n(\langle x \rangle)) = s_n(f_i(\langle x \rangle))$ and $f(t_n(\langle x \rangle)) = f_i(t_n(\langle x \rangle)) = t_n(f_i(\langle x \rangle))$. Furthermore, if we have another j such that $x \in C^{(j)}$ then by the cocone property of the family f_i we know that $f_i(\langle x \rangle) = f_j(\langle x \rangle)$, and it follows that we may use the free generation property of $\mathcal{O}(C)$ to extend f uniquely to a functor on $|\mathcal{O}(C)|_{n+1}$ for which $f(\langle x \rangle) = f_i(\langle x \rangle)$ for all $i \in I$ and $x \in C_{n+1}^{(i)}$. But if f and f_i agree on these atoms they must agree in the whole of $|\mathcal{O}(C^{(i)})|_{n+1}$. Applying this extension result on successive skeleta, we may construct a unique ω -functor whose domain is their union $\mathcal{O}(C)$ and which restricts to f_i on each $\mathcal{O}(C^{(i)})$ as required. $\square$

The next two, very simple, technical results provide us with a convenient way to construct isomorphisms with freely generated ω -categories.

Lemma 221. *Let $e: \mathbb{C} \longrightarrow \mathbb{D}$ be an epimorphism of ω -categories and let $G \subseteq \mathbb{C}$ be a subset which weakly generates $\mathbb{C}$. Consider a subset $H \subseteq \mathbb{D}$, which we grade as in definition 218, and suppose that $e(G_0) \subseteq H_0$ and $e(G_{n+1}) \setminus |\mathbb{D}|_n \subseteq H_{n+1}$ (for each $n \in \mathbb{N}$) then H weakly generates $\mathbb{D}$.*

Proof. First note that e restricts to an epimorphism $e: |\mathbb{C}|_n \longrightarrow |\mathbb{D}|_n$ for each $n \in \mathbb{N}$ (by observation 84). Epimorphisms between 0-categories (sets) are all surjective and $G_0 = |\mathbb{C}|_0$ so from the condition on H_0 in the statement it is immediate that we must have $H_0 = |\mathbb{D}|_0$. To establish the separation property of weak generation, consider a pair of ω -functors $f, g: |\mathbb{D}|_{n+1} \longrightarrow \mathbb{E}$ with $f(d) = g(d)$ for all $d \in |\mathbb{D}|_n \cup H_{n+1}$. Of course, we know that $e(|\mathbb{C}|_n) \subseteq |\mathbb{D}|_n$ (since ω -functors carry n -cells to n -cells) and the condition on H_{n+1} in the statement is equivalent to $e(G_{n+1}) \subseteq |\mathbb{D}|_n \cup H_{n+1}$, therefore $e(|\mathbb{C}|_n \cup G_{n+1}) = e(|\mathbb{C}|_n) \cup e(G_{n+1}) \subseteq |\mathbb{D}|_n \cup H_{n+1}$. It follows, from the fact that f and g coincide on $|\mathbb{D}|_n \cup H_{n+1}$ by assumption, that $f(e(c)) = g(e(c))$ for each $c \in |\mathbb{C}|_{n+1}$. However G weakly generates $\mathbb{C}$ so we may infer that $f \circ e = g \circ e$ and use the epimorphism property of $e: |\mathbb{C}|_{n+1} \longrightarrow |\mathbb{D}|_{n+1}$ to show then that $f = g$ as required. $\square$

Observation 222. A partial converse to the last lemma is the simple observation that if $f: \mathbb{C} \longrightarrow \mathbb{D}$ is an ω -functor and H is a weak generator of $\mathbb{D}$ for which $H \subseteq f(\mathbb{C})$ then f is an epimorphism. In many cases we are presented with a jointly epimorphic family of ω -functors $\{e_i: \mathbb{C}_i \longrightarrow \mathbb{D}\}_{i \in I}$, rather than a single epimorphism, and a weak generator $G^{(i)}$ for each $\mathbb{C}_i$. In particular, this is the case when $\mathbb{D}$ is the colimit of some diagram and the ω -functors e_i are the components of the colimiting cocone. Of course, in such a case the coproduct ω -category $\coprod_{i \in I} \mathbb{C}_i$ is weakly generated by the subset $\coprod_{i \in I} G^{(i)}$ and we may apply the result of the last lemma to the induced epimorphism $\langle e_i \rangle_{i \in I}: \coprod_{i \in I} \mathbb{C}_i \longrightarrow \mathbb{D}$.

Lemma 223. *Suppose that $\mathbb{C}$ and $\mathbb{D}$ are ω -categories and that we have sets of cells $G \subseteq \mathbb{C}$ and $H \subseteq \mathbb{D}$ such that G **freely** generates $\mathbb{C}$ whereas H **weakly** generates $\mathbb{D}$. Assume further that we have an ω -functor $f: \mathbb{D} \longrightarrow \mathbb{C}$ which restricts to a bijection between the sets H_n and G_n for all $n \in \mathbb{N}$ then f is actually an **isomorphism** of ω -categories.*

Proof. Before proving this result, it's worth making a slight methodological point. Its important here that we've only made the assumption that H weakly generates $\mathbb{D}$. In all of the applications we have in mind $\mathbb{D}$ is obtained by taking some kind of quotient of a freely

generated ω -category. In such situations it's usually relatively easy to construct functors with domain $\mathbb{D}$ and even to prove weak generation by a set of cells, however it is rarely easy to directly establish free generation.

Our proof proceeds by induction on superstructures, demonstrating that f restricts to an isomorphism between each corresponding pair of superstructures. For the base case, definition 218 explicitly postulates that $G_0 = |\mathbb{C}|_0$ and $H_0 = |\mathbb{D}|_0$ and we assumed that f restricts to a bijection from H_0 to G_0 which thus provides a isomorphism between the discrete ω -categories $|\mathbb{C}|_0$ and $|\mathbb{D}|_0$.

To establish the induction step, we fix an $n \in \mathbb{N}$ and make the inductive hypothesis that the restriction $f: |\mathbb{D}|_n \longrightarrow |\mathbb{C}|_n$ is an isomorphism of ω -categories, suppose that it has inverse $g: |\mathbb{C}|_n \longrightarrow |\mathbb{D}|_n$ in $\omega\text{-Cat}$ and let $f^{-1}: G_{n+1} \longrightarrow H_{n+1}$ denote the postulated inverse to f on sets of $(n+1)$ -dimensional generators. Now consider an arbitrary $c \in G_{n+1}$ and observe that we have $\mathbf{s}_n(f^{-1}(c)) = g(f(\mathbf{s}_n(f^{-1}(c)))) = g(\mathbf{s}_n(f(f^{-1}(c)))) = g(\mathbf{s}_n(c))$ where the first equality holds since g and f are inverses on n -superstructures (by the inductive hypothesis), the second holds because f is an ω -functor (and thus commutes with n -sources) and the last follows from the fact that f and f^{-1} are inverses on sets of generators. We can argue similarly with n -targets and so it follows that we may apply the free generation property of G to extend g to a unique ω -functor $h: |\mathbb{C}|_{n+1} \longrightarrow |\mathbb{D}|_{n+1}$ with $h(c) = f^{-1}(c)$ for all $c \in G_{n+1}$. By definition, we easily see now that $f(h(c)) = c$ for all $c \in |\mathbb{C}|_n \cup G_{n+1}$ and $h(f(d)) = d$ for all $d \in |\mathbb{D}|_n \cup H_{n+1}$, so applying the weak generation properties of G and H we infer that $f \circ h = \text{id}_{|\mathbb{C}|_{n+1}}$ and $h \circ f = \text{id}_{|\mathbb{D}|_{n+1}}$. Thus the restriction $f: |\mathbb{D}|_{n+1} \longrightarrow |\mathbb{C}|_{n+1}$ is also an isomorphism of ω -categories, with inverse h .

Finally, our inductive argument has shown that f restricts to an isomorphism between each corresponding pair of superstructures and, using the fact that an ω -category is the union of its superstructures, it follows that it is itself an isomorphism of ω -categories. $\square$

In the sequel, we'll use a few properties of the ω -category $\mathcal{O}(C)$ generated by a parity complex C . These are an immediate consequence of Street's work in [33] and [34] but are not explicitly remarked upon there.

Observation 224. If S is a finite, non-empty subset of parity complex C then axiom 3(a) of definition 216 ensures that we may find a $\triangleleft_S$ minimal element s in S . By minimality it follows that $s^- \cap c^+ = \emptyset$ for all $c \in S$ and thus we know that $s^- \subseteq S^\mp$. Arguing dually, we also see that we have an element $t \in S$ with $t^+ \subseteq S^\pm$. In particular, these observations imply that both of the sets $S^\mp$ and $S^\pm$ are non-empty.

Applying this result to a cell (M, P) of $\mathcal{O}(C)$ we see that if M_n ($n > 0$) is non-empty then it contains elements s and t such that $s^- \subseteq M_n^\mp$ and $t^+ \subseteq M_n^\pm$. However, the movement property of our cell implies that $M_n^\mp \subseteq M_{n-1}$ and $M_n^\pm \subseteq P_{n-1}$, so it follows that $s^- \subseteq M_{n-1}$ and $t^+ \subseteq P_{n-1}$ and in particular that M_{n-1} and P_{n-1} are also non-empty. Of course, we may apply exactly the same argument to P_n to obtain precisely the same result under the assumption that it is non-empty.

Now, every cell (M, P) has a maximum n for which M_n or P_n is non-empty (since M and P are finite and non-empty) and by the movement property for $P_{n+1} = M_{n+1} = \emptyset$ we have $M_n = P_n$, it follows that (M, P) is a non-trivial n -cell. Applying the result of the last paragraph repeatedly, it follows also that M_r and P_r are non-empty for every $r < n$.

Lemma 225. Suppose that C is a parity complex and that there is some $x \in C_n$ ($n > 0$) which generates it, in the sense that C is the smallest sub-parity complex of itself which contains x ,

then if we are given an element $y \in C_m$ ($m > 0$) we may construct an $(m+1)$ -cell (M, P) in $\mathcal{O}(C)$ with $y \in M_m$ and $\mathbf{t}_m(M, P) = \mathbf{t}_m(\langle x \rangle)$.

Proof. The sub-parity complex generated by a subset $S \subseteq C$ is simply the smallest substructure of C which is closed under the face operations and which contains S . It follows, therefore, that $x \in C_n$ generates C if and only if $C_n = \{x\}$ and for all $y \in C_m$ with $m < n$ there exists a $z \in C_{m+1}$ such that $y \in z^- \cup z^+$. Notice also that this also implies that $C_m = \emptyset$ for $m > n$.

Armed with this observation, we prove our result by “downward” induction on m . The base case $m = n$ is trivial, since in that case $y = x$ and we may take the atom $\langle x \rangle$ itself as our $(m+1)$ -cell (M, P) .

So assume the inductive hypothesis, that our result is true at dimensions greater than m , and consider an element $y \in C_m$. We know, by our initial observation, that there is some $z \in C_{m+1}$ for which $y \in z^- \cup z^+$ and we may apply our inductive hypothesis to z to construct an $(m+2)$ -cell (N, Q) in $\mathcal{O}(C)$ which has $z \in N_{m+1}$ and $\mathbf{t}_{m+1}(N, Q) = \mathbf{t}_{m+1}(\langle x \rangle)$. However, suppose that $y \in z^+$ then, by the movement property of N_{m+1} , we know that $Q_m = (N_m \cup N_{m+1}^+) \setminus N_{m+1}^-$ so either $y \in Q_m$ or $y \in N_{m+1}^-$. In the first case there is nothing more to do since we can take $(M, P) = \mathbf{t}_m(N, Q)$, in the second case we can replace z by the element of N_{m+1}^- which counts y amongst its negative faces.

Thus from now on we may assume that $y \in z^-$, define $X = \{w \in N_{m+1} \mid z \triangleleft_N w\}$ and observe that since N is well formed, by the well formedness property of the cell (N, Q) , then so is the subset X . Furthermore we have:

$X^\pm \subseteq N_{m+1}^\pm \subseteq Q_m$, certainly we know that the second of these inclusions holds, by the movement property of the cell (N, Q) , so to prove the former suppose that $u \in X^+ \setminus N_{m+1}^\pm$ then we have

$$X^+ \setminus N_{m+1}^\pm = X^+ \setminus (N_{m+1}^+ \setminus N_{m+1}^-) = (X^+ \setminus N_{m+1}^+) \cup (X^+ \cap N_{m+1}^-) = X^+ \cap N_{m+1}^-$$

where the last of these equalities holds since $X \subseteq N_{m+1}$ which implies that $X^+ \subseteq N_{m+1}^+$, so it follows that we have a $v \in N_{m+1}$ and a $w \in X$ with $u \in w^+ \cap v^-$. However, the fact that this latter intersection is non-empty tells us that $w \triangleleft_N v$ and we know that $z \triangleleft_N w$, by the definition of X of which w is an element, so by transitivity we get $z \triangleleft_N v$ and thus we have $v \in X$, from which it follows that $u \in v^- \subseteq X^-$. In summary we have shown that $(X^+ \setminus N_{m+1}^\pm) \setminus X^- = X^\pm \setminus N_{m+1}^\pm$ is empty, or in other words that $X^\pm \subseteq N_{m+1}^\pm$ as required.

$z^- \subseteq (Q_m \cup X^-) \setminus X^+$, we know that $\triangleleft_N$ is reflexive so $z \in X$ and it follows that $z^- \subseteq Q_m \cup X^-$, therefore all we have to prove is that $z^- \cap X^+$ is empty. So suppose the converse, for a contradiction, and observe that it would imply that there is some $w \in X$ with $z^- \cap w^+ \neq \emptyset$, in which case we have $w \triangleleft_N z$, by definition, and $w \neq z$, since all elements have disjoint sets of negative and positive faces. However, by the definition of X , of which w is an element, we also know that $z \triangleleft_N w$ and when combined these three facts contradict axiom 3(a) of definition 216 as required.

Now, the first of these results implies that we may apply lemma 3.2 of [33] to the m -cell $\mathbf{t}_m(N, Q) = (|N|_{m-1} \cup Q_m, |Q|_m)$ and the subset X . Part (c) of that lemma allows us to construct an $(m+1)$ -cell $(M, P) \stackrel{\text{def}}{=} (|N|_{m-1} \cup Y \cup X, |Q|_m \cup X) \in \mathcal{O}(C)$, where Y is the subset $(Q_m \cup X^-) \setminus X^+$ discussed in the second of the results above. However, that analysis tells us that we have $z^- \subseteq Y$ and, since we originally selected z so that $y \in z^-$, it follows that $y \in Y = M_m$. Finally, by construction, it is clear that $\mathbf{t}_m(M, P) = \mathbf{t}_m(N, Q)$

and since we originally selected (N, Q) to have $\mathbf{t}_{m+1}(N, Q) = \mathbf{t}_{m+1}(\langle x \rangle)$ it follows that $\mathbf{t}_m(M, P) = \mathbf{t}_m(\langle x \rangle)$ as required. $\square$

Corollary 226. *Suppose that C is the parity complex of lemma 225 then the atom $\langle x \rangle$ is the **unique** non-trivial n -cell of the n -category $\mathcal{O}(C)$.*

Proof. Suppose that (M, P) is a non-trivial n -cell of $\mathcal{O}(C)$ then since x is the unique n -dimensional element of C we must have $M_n = P_n = \{x\}$. Now, for a contradiction, assume that $(M, P) \neq \langle x \rangle$, and observe that this assumption implies that there is some $m < n$ and some cell which is not an m -cell but which is m -composable with $\langle x \rangle$.

To establish this fact, observe that we may apply theorem 4.1 of [33] to construct an $m < n$ and two n -cells (N, Q) and (L, R) which are not m -cells and for which $(M, P) = (N, Q) *_m (L, R)$. Notice that one of these new cells must be a non-trivial n -cell, since (M, P) itself is, and indeed we may assume without loss of generality that this is actually equal to the atom $\langle x \rangle$. Were this not the case then we could simply iterate our argument and use the rank of cells, as in theorem 4.2 of [33], to show that this process must eventually terminate at the atom $\langle x \rangle$.

Without loss of generality we may assume that the composite we constructed is $\langle x \rangle *_m (L, R)$, were it the reverse composite we would simply argue dually. We know that (L, R) is not an m -cell and so, by observation 224, we may infer that there is some $y \in R_{m+1}$ such that $y^+ \subseteq R_m$. However $R_m = \mu(x)_m$, since $\langle x \rangle$ and (L, R) are m -composable, so we may infer that $y^+ \subseteq \mu(x)_m$.

Applying lemma 225 we obtain an $(m+2)$ -cell (M, P) which has $y \in M_{m+1}$ and which has $\mathbf{t}_{m+1}(M, P) = \mathbf{t}_{m+1}(\langle x \rangle)$. By the movement property for this cell we know that M_{m+1} moves $M_m = \mu(x)_m$ to $P_m = \pi(x)_m$ so in particular $\mu(x)_m = (\pi(x)_m \cup M_{m+1}^-) \setminus M_{m+1}^+$, however we have $y \in M_{m+1}$ thus $y^+ \subseteq M_{m+1}^+$ and it follows that $\mu(x)_m \cap y^+ = ((\pi(x)_m \cup M_{m+1}^-) \setminus M_{m+1}^+) \cap y^+ = \emptyset$. This, provides us with the desired contradiction, since in the last paragraph we constructed y so that $y^+ \subseteq \mu(x)_m$ and we know that y^+ itself is non-empty. $\square$

We may represent standard simplices, cubes and product polytopes as parity complexes and thereby construct appropriate ω -categorical models of such structures.

Observation 227 (ω -categorical simplex models). Following Street in [31], we may define the standard ω -simplex $\tilde{\Delta}$ to be the graded set for which $\tilde{\Delta}_r$ is the set of $(r+1)$ -element (indexed) subsets $\underline{v} = \{v_0 < v_1 < \dots < v_r\} \subseteq \mathbb{N}$ which we often simply write as $v_0 \dots v_r$. In essence each v_i is thought of as a distinct vertex of an infinite dimensional simplex and $v_0 \dots v_r$ is thought of as its r -dimensional face spanning the listed vertices.

Alternatively, we may represent an element $\underline{v} \in \tilde{\Delta}_r$ as the strictly order preserving map $\delta_{\underline{v}}$ from $[r]$ to $\mathbb{N}$ which maps i to v_i . In general, we will identify these representations, passing between them as the mood takes us and without comment.

Of course, if $\alpha: [p] \longrightarrow [r]$ is an arbitrary face operator then we may pre-compose it with $\delta_{\underline{v}} \in \tilde{\Delta}_r$ to give an element $\delta_{\underline{v}} \circ \alpha \in \tilde{\Delta}_p$. Using our vectorial notation we write $\underline{v} \cdot \alpha$ or $v_{\alpha(0)} v_{\alpha(1)} \dots v_{\alpha(p)}$ for the subset corresponding to the map $\delta_{\underline{v}} \circ \alpha$, and this action makes $\tilde{\Delta}$ into a semi-simplicial set. In terms of our vertex-wise notation we will sometimes use the traditional algebraic topologist's device $v_0 \dots \hat{v}_i \dots v_r$ where this denotes the face of $\underline{v}$ obtained by omitting the vertex v_i , in other words it is an alternative way of writing $\underline{v} \cdot \delta_i^r$.

Now we may use this semi-simplicial structure to define a pre-parity complex structure on $\tilde{\Delta}$ by:

$$\underline{v}^\xi = \{\underline{v} \cdot \delta_i^r \mid i \in [r] \text{ and } i \text{ is of parity } \xi\}$$

As demonstrated in [33] and [34], this actually makes $\tilde{\Delta}$ into a parity complex satisfying all of the conditions required for us to be able to apply theorem 219 and construct a free ω -category $\mathcal{O}(\tilde{\Delta})$.

We obtain an ω -categorical model of the n -simplex by restricting our attention to the sub-parity complex $\tilde{\Delta}[n]$ of $\tilde{\Delta}$ consisting of those elements $\underline{v}$ for which each v_i is actually an element of the ordered set $[n]$. In other words, this is the same as saying that the corresponding map $\delta_{\underline{v}}$ restricts to a face operator with codomain $[n]$. When manipulating the elements of $\tilde{\Delta}[n]$ as face operators well often apply a superscripted n , as in $\delta_{\underline{v}}^n$, to remind ourselves of this restricted codomain. The resulting sub- n -category $\mathcal{O}(\tilde{\Delta}[n]) \subseteq \mathcal{O}(\tilde{\Delta})$ was dubbed the n^{th} oriental in [31].

Notice that if $\underline{v}$ is an r -dimensional element of $\tilde{\Delta}[n]$ then $\underline{v}^- \cup \underline{v}^+$ is the set of all $(r-1)$ -dimensional subsets of $\underline{v}$. Applying this result recursively, it follows therefore that the sub-parity complex of $\tilde{\Delta}[n]$ generated by $\underline{v}$ consists of the set of all those elements $\underline{w} \in \tilde{\Delta}[n]$ which have $\underline{w} \subseteq \underline{v}$. In particular, it follows that the n -dimensional element $01\dots n$ generates $\tilde{\Delta}[n]$ as a parity complex. So we may apply corollary 226 to show that the atom $\langle 01\dots n \rangle$ is the unique non-trivial n -cell of the n -category $\mathcal{O}(\tilde{\Delta}[n])$.

Observation 228 (products of parity complexes and models of cubes). The product $C \times D$ of two pre-parity complexes C and D is the graded set defined by

$$(C \times D)_n = \bigcup_{p+q=n} C_p \times D_q$$

in which the ξ -parity faces of an element $(x, y) \in C_p \times D_q$ are given by

$$(x, y)^\xi = x^\xi \times \{y\} \cup \{x\} \times y^{\xi(p)}$$

where $\xi(p)$ is equal to ξ if p is even and is the opposite parity $\neg\xi$ if p is odd.

In [33] and [34] it is also shown that if C and D are parity complexes which satisfy the fairly ubiquitous conditions alluded to in definition 216 then $C \times D$ is also a parity complex which satisfies those conditions. This implies that we can apply theorem 219 to an arbitrary n -fold product $C_1 \times C_2 \times \dots \times C_n$ of such parity complexes to obtain a free ω -category $\mathcal{O}(C_1 \times C_2 \times \dots \times C_n)$.

A particular example of this is the ω -categorical model of the n -cube, which is defined to be the free ω -category on the n -fold product of the 1-simplex parity complex $\tilde{\Delta}[1]$ with itself.

Notice that if the element $x \in C_n$ generates C and $y \in D_m$ generates D (as in lemma 225) then we may show that (x, y) generates the product $C \times D$, as the reader may readily verify using the characterisation of generation given in the first paragraph of the proof of lemma 225. In particular we know, from observation 227, that the element $01\dots n$ generates $\tilde{\Delta}[n]$ so it follows that $(01\dots n, 01\dots m)$ generates the product $\tilde{\Delta}[n] \times \tilde{\Delta}[m]$ and thus that the atom $\langle 01\dots n, 01\dots m \rangle$ is the unique non-trivial $(n+m)$ -cell of the $(n+m)$ -category $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ (by corollary 226).

Observation 229 (defining functors between free ω -categories on parity complexes). In the sequel we will need to construct a number of ω -functors between ω -categories of the form $\mathcal{O}(C)$. To this end, consider a pair of graded sets C and D and a function $f: C \dashrightarrow D$ which maps elements of C to finite subsets of D and respects dimensions in the sense that for all $x \in C_n$ we have $f(x) \subseteq D_n$. We call such a function a *graded set morphism*.

As usual, if $S \subseteq C$ then we'll use $f(S)$ to denote the union $\bigcup_{x \in S} f(x)$ and observe that since f respects dimensions we have $f(S_n) = f(S)_n$, $f(|S|_n) = |f(S)|_n$ and $f(S^{-n}) = f(S)^{-n}$.

for each $n \in \mathbb{N}$. Furthermore it is clear that if S is finite then so is $f(S)$ and that if $T \subseteq C$ is another such subset then we have $f(S \cup T) = f(S) \cup f(T)$.

Notice that these are precisely the operations and conditions which are used to define the ω -category $\mathcal{N}(C)$. So since they are all preserved by f it follows easily that the function which maps a pair (M, P) in $\mathcal{N}(C)$ to the pair $(f(M), f(P))$ in $\mathcal{N}(D)$ is an ω -functor, for which we shall use the notation $\mathcal{N}(f)$.

Indeed we may construct a category GrSet with objects that are graded sets and arrows which are graded set morphisms. Composites and identities in this category are defined in the obvious way, that is $\text{id}_C(x) = \{x\}$ for a graded set C and $g \circ f(x) = g(f(x)) = \bigcup_{y \in f(x)} g(y)$ ($\forall x \in C$) for a composable pair $f: C \twoheadrightarrow D$ and $g: D \twoheadrightarrow E$. Now it is clear that the construction of $\mathcal{N}(f)$ given in the last paragraph makes $\mathcal{N}$ into a functor from this category GrSet to the category of ω -categories $\omega\text{-Cat}$.

In the case where C and D are parity complexes, we would now like to determine some easily verifiable conditions under which the induced ω -functor $\mathcal{N}(f): \mathcal{N}(C) \longrightarrow \mathcal{N}(D)$ restricts to give an ω -functor $\mathcal{O}(f): \mathcal{O}(C) \longrightarrow \mathcal{O}(D)$. A simple, but adequate, characterisation of those functions for which this holds is provided by lemma 231.

Observation 230. Before stating the next lemma, it is worth commenting on the fact that the conditions in its statement are motivated by a simple inductive characterisation of the $(n+1)$ -cells (M, P) of $\mathcal{O}(C)$. This states that the $(n+1)$ -cell $(M, P) \in \mathcal{N}(C)$ is in $\mathcal{O}(C)$ if and only if $s_n(M, P)$ and $t_n(M, P)$ are both in $\mathcal{O}(C)$ and M_{n+1} (which is equal to P_{n+1}) is well-formed and moves M_n to P_n .

To prove this characterisation first observe that, since we know that $\mathcal{O}(C)$ is closed under sources and targets in $\mathcal{N}(C)$, it is enough to fix an $(n+1)$ -cell $(M, P) \in \mathcal{N}(C)$ whose n -source and n -target are both in $\mathcal{O}(C)$ and show that the residual well-formedness and movement condition characterises when our $(n+1)$ -cell may actually be found to be in $\mathcal{O}(C)$.

Notice that since (M, P) is an $(n+1)$ -cell in $\mathcal{N}(C)$ we know that $M = |M|_n \cup M_{n+1}$ and $P = |P|_n \cup M_{n+1}$ and, furthermore, it is clear that well-formedness and movement are properties which respect dimension. It follows therefore that M and P are well-formed iff $|M|_n$, $|P|_n$ and M_{n+1} are all well-formed and that they each move M to P iff $|M|_n$ and $|P|_n$ each move $|M|_{n-1}$ to $|P|_{n-1}$ and M_{n+1} moves M_n to P_n .

However we have $s_n(M, P) = (|M|_n, M_n \cup |P|_{n-1}) \in \mathcal{O}(C)$ so it follows, by the definition of the cells that occur in $\mathcal{O}(C)$, that $|M|_n$ is non-empty and well-formed and that it moves $|M|_{n-1}$ to $|P|_{n-1}$. Similarly, considering $t_n(M, P) = (|M|_{n-1} \cup P_n, |P|_n) \in \mathcal{O}(C)$ we see, dually, that $|P|_n$ is also non-empty and well-formed and furthermore that it too moves $|M|_{n-1}$ to $|P|_{n-1}$. So finally, excluding these observations from the list of conditions in the last paragraph, we see that the remaining conditions required to show that (M, P) is a cell of $\mathcal{O}(C)$ are simply those postulated in the statement above. $\square$

Lemma 231. Suppose that C and D are parity complexes and $f: C \twoheadrightarrow D$ is a graded set morphism as discussed in observation 229. Then the ω -functor $\mathcal{N}(f): \mathcal{N}(C) \longrightarrow \mathcal{N}(D)$ restricts to give an ω -functor $\mathcal{O}(f): \mathcal{O}(C) \longrightarrow \mathcal{O}(D)$ if and only if f satisfies the following conditions:

- (a) for all $x \in C_0$ the subset $f(x) \subseteq D_0$ is a singleton, and
- (b) for all $n \in \mathbb{N}$ and $x \in C_{n+1}$ the subset $f(x) \subseteq D_{n+1}$ is well formed and moves $f(x^-)$ to $f(x^+)$ or symbolically $f(x): f(x^-) \longrightarrow f(x^+)$.

Consequently we have a subcategory Parity of GrSet, whose objects are parity complexes and whose arrows are those graded set morphisms which satisfy these two conditions, which we shall often refer to as **parity complex morphisms**.

Clearly then, our free ω -category construction lifts to a functor $\mathcal{O}: \underline{\text{Parity}} \longrightarrow \omega\text{-Cat}$ which acts on an arrow $f \in \underline{\text{Parity}}$ by restricting the action of $\mathcal{N}(f)$ as described.

Proof. The necessity of this equivalence follows immediately from the following observations:

- (i) If $x \in C_0$ then $\mathcal{N}(f)(\langle x \rangle) = (f(x), f(x))$ which is a 0-cell of $\mathcal{O}(D)$ if and only if $f(x)$ is a singleton subset of D_0 .
- (ii) If $x \in C_{n+1}$ then $\mathcal{N}(f)(\langle x \rangle) = (f(\mu(x)), f(\pi(x)))$ for which we have $f(\mu(x))_{n+1} = f(\pi(x))_{n+1} = f(x)$, $f(\mu(x))_n = f(x^-)$ and $f(\pi(x))_n = f(x^+)$. It follows, by observation 230, that $\mathcal{N}(f)(\langle x \rangle)$ is in $\mathcal{O}(D)$ if and only if $\mathbf{s}_n(\mathcal{N}(f)(\langle x \rangle)) = \mathcal{N}(f)(\mathbf{s}_n(\langle x \rangle))$ and $\mathbf{t}_n(\mathcal{N}(f)(\langle x \rangle)) = \mathcal{N}(f)(\mathbf{t}_n(\langle x \rangle))$ are in $\mathcal{O}(D)$ and $f(x)$ is a well formed set that moves $f(x^-)$ to $f(x^+)$.

We also prove sufficiency from these observations by inductively demonstrating that $\mathcal{N}(f)$ restricts to an ω -functor $|\mathcal{O}(C)|_n \longrightarrow \mathcal{O}(D)$ for each dimension n in turn and arguing that it therefore restricts in this way to the whole of $\mathcal{O}(C)$, since this is equal to the union of its superstructures. To that end we assume that f satisfies conditions (a) and (b) and proceed by induction on n .

The base case $n = 0$ of our induction is easy, since we know that the cells of $|\mathcal{O}(C)|_0$ are simply the atoms $\langle x \rangle$ for $x \in C_0$ so by applying observation (i) and condition (a) we see that $\mathcal{N}(f)$ restricts to an ω -functor $|\mathcal{O}(C)|_0 \longrightarrow \mathcal{O}(D)$.

So assume the inductive hypothesis, that $\mathcal{N}(f)$ restricts to an ω -functor from $|\mathcal{O}(C)|_n$ to $\mathcal{O}(D)$, and consider an arbitrary atom in $|\mathcal{O}(C)|_{n+1}$. Either this is in $|\mathcal{O}(C)|_n$ itself, in which case $\mathcal{N}(f)$ maps it into $\mathcal{O}(D)$ by hypothesis, or it is of the form $\langle x \rangle$ with $x \in C_{n+1}$. In the latter case the n -source and n -target of $\langle x \rangle$ are, of course, both cells of $|\mathcal{O}(C)|_n$ so by the inductive hypothesis it follows that $\mathcal{N}(f)(\mathbf{s}_n(\langle x \rangle))$ and $\mathcal{N}(f)(\mathbf{t}_n(\langle x \rangle))$ are both n -cells in $\mathcal{O}(D)$. Combining this information with condition (b) gives us enough to apply observation (ii) and infer that $\mathcal{N}(f)(\langle x \rangle)$ is in $\mathcal{O}(D)$. It follows therefore that $\mathcal{N}(f)$ maps every atom of dimension $\leq n + 1$ into $\mathcal{O}(D)$.

Now, by theorem 219 we know that the sub- ω -category $|\mathcal{O}(C)|_{n+1} \subseteq \mathcal{N}(C)$ is freely generated by the atoms of dimension $\leq n + 1$ and, in particular, this implies that every cell of $|\mathcal{O}(C)|_{n+1}$ may be obtained as some finite ω -categorical composite of such atoms. So since $\mathcal{N}(f)$ preserves these composites and maps all atoms of dimension $\leq n + 1$ into $\mathcal{O}(D)$ it follows that it restricts to an ω -functor $|\mathcal{O}(C)|_{n+1} \longrightarrow \mathcal{O}(D)$ as required. $\square$

Observation 232. As an aside, it is worth observing that the movement property of condition (b) of the last lemma may be summarised conveniently in the statement that the single equation

$$(f(x^\xi) \cup f(x)^{-\xi}) \setminus f(x)^\xi = f(x^{-\xi})$$

should hold for an arbitrary parity $\xi \in \{-, +\}$.

The following lemma and its corollary often allows us to determine the action of an ω -functor of the form $\mathcal{O}(f)$ on the atoms $\langle x \rangle$ in its domain without requiring us to perform detailed calculations directly on the combinatorially complex sets $\mu(x)$ and $\pi(x)$.

Lemma 233. Suppose that $f: C \longrightarrow D$ is a morphism of parity complexes and that $x \in C_n$ generates the sub-parity complex E of C . If we let $m \stackrel{\text{def}}{=} \max\{i \leq n \mid f(E_i) \neq \emptyset\}$ then the cell $\mathcal{O}(f)(\langle x \rangle)$ is a non-trivial m -cell in $\mathcal{O}(D)$.

Proof. Notice that our choice of m ensures that f maps every element of E of dimension $> m$ to the empty set, so it follows that $\mathcal{O}(f)$ maps every cell of $\mathcal{O}(E) \subseteq \mathcal{O}(C)$ to an m -cell of $\mathcal{O}(D)$. In particular since x generates E we know that the associated atom $\langle x \rangle$ is a cell in the sub- ω -category $\mathcal{O}(E) \subseteq \mathcal{O}(C)$ and so it follows that $\mathcal{O}(f)(\langle x \rangle)$ is indeed an m -cell in $\mathcal{O}(D)$. It remains to show that $\mathcal{O}(f)(\langle x \rangle)$ is actually a *non-trivial* m -cell.

To do this we select a $y \in E_m$ such that $f(y) \neq \emptyset$ and apply lemma 225 to obtain an $(m+1)$ -cell (M, P) in $\mathcal{O}(E)$ which has $y \in M_m$ and $t_m(M, P) = t_m(\langle x \rangle)$. Now, arguing as in the previous paragraph we know that $\mathcal{O}(f)(M, P)$ is an m -cell and it follows that $t_m(\mathcal{O}(f)(\langle x \rangle)) = \mathcal{O}(f)(t_m(\langle x \rangle)) = \mathcal{O}(f)(t_m(M, P)) = t_m(\mathcal{O}(f)(M, P)) = \mathcal{O}(f)(M, P)$. In particular this equation implies that $f(\pi(x)_m) = f(M_m)$ and, by construction, we know that $y \in M_m$ and it follows that $f(M_m) \neq \emptyset$ because y was originally chosen so that $f(y) \neq \emptyset$. So $f(\pi(x)_m) \neq \emptyset$ and it follows that $\mathcal{O}(f)(\langle x \rangle) = (f(\mu(x)), f(\pi(x)))$ has a non-empty set of m -dimensional elements and thus is a non-trivial m -cell in $\mathcal{O}(D)$ as postulated. $\square$

Corollary 234. *Under the conditions described in lemma 233, if F is the sub-parity complex of D generated by $f(E)$ and it may also be generated by a single element $y \in F$ then this element is m -dimensional, where m is defined as in the statement of lemma 233, and $\mathcal{O}(f)(\langle x \rangle) = \langle y \rangle$.*

Proof. We may construct the sub-parity complex F generated by a subset $S \subseteq D$ by closing S up in D under its face operations, which carry each element to subsets of elements of lower dimension. It follows therefore that the maxima of the dimensions of elements in S and in D must be equal. In particular, in lemma 233 we defined m to be the maximum dimension of elements in $f(E)$, so we may infer that the elements of F have maximum dimension F . Furthermore, since the single element y also generates F it follows the dimension of y must also be m .

Now, applying lemma 226 we see that the atom $\langle y \rangle$ is the unique non-trivial m -cell of the sub- ω -category $\mathcal{O}(F)$ of $\mathcal{O}(D)$. However, applying lemma 233 we know that $\mathcal{O}(f)(\langle x \rangle)$ is also a non-trivial m -cell in $\mathcal{O}(F)$ and so it follows that $\mathcal{O}(f)(\langle x \rangle) = \langle y \rangle$ as required. $\square$

Observation 235 (the product of parity complexes as a bifunctor). In fact, we may extend the product of parity complexes to a bifunctor from Parity $\times$ Parity to Parity by defining the product $f \times g: C \times D \longrightarrow E \times F$ of a pair of arrows $f: C \longrightarrow E$ and $g: D \longrightarrow F$ in Parity to be the function given by $(f \times g)(x, y) = f(x) \times g(y)$. Notice that condition (a) of lemma 231 holds trivially for this definition and, indeed, it is only slightly more difficult to prove that it also satisfies condition (b). To do this we consider an arbitrary element $(x, y) \in C_p \times D_q$ and show that:

$(f \times g)(x, y)$ is well formed: notice first that we know that $f(x) \subseteq E_p$ and $g(y) \subseteq F_q$ are well formed, since f and g are arrows of Parity, and suppose that (u, v) and (w, z) are arbitrary but distinct elements of the subset $(f \times g)(x, y) = f(x) \times g(y)$ of $E_p \times F_q$. We know that $(u, v)^\xi = u^\xi \times \{v\} \cup \{u\} \times v^{\xi(p)}$ and $(w, z)^\xi = w^\xi \times \{z\} \cup \{w\} \times z^{\xi(p)}$ and that the left hand factors in these unions are subsets of $E_{p-1} \times F_q$ whereas their right hand factors are subsets of the disjoint set $E_p \times F_{q-1}$. So it follows that if $(u, v)^\xi \cup (w, z)^\xi \neq \emptyset$ then either $v = z$, $u \neq w$ and $u^\xi \cap w^\xi \neq \emptyset$, which contradicts the well-formedness of $f(x)$, or $u = w$, $v \neq z$ and $v^{\xi(p)} \cap z^{\xi(p)} \neq \emptyset$, which contradicts the well-formedness of $g(y)$. It follows therefore that $(u, v)^\xi \cup (w, z)^\xi = \emptyset$ and thus that $f(x) \times g(y)$ is well formed as required.

$(f \times g)(x, y)$ moves $(f \times g)((x, y)^-)$ to $(f \times g)((x, y)^+)$: it is easily show that $((f \times g)(x, y))^\xi = f(x)^\xi \times g(y) \cup f(x) \times g(y)^\xi$ and that $(f \times g)((x, y)^\xi) = f(x^\xi) \times g(y) \cup$

$f(x) \times g(y^{\xi(p)})$ and it is again the case that the left factors of these unions are in the set $E_{p-1} \times F_q$ whereas their right factors are in the disjoint set $E_p \times F_{q-1}$. It follows that the movement property of observation 232 holds if and only if it can be verified independently on corresponding factors in these unions. However we have

$$\begin{aligned} (f(x^\xi) \times g(x) \cup f(x)^{-\xi} \times g(x)) \setminus (f(x)^\xi \times g(x)) &= \\ ((f(x^\xi) \cup f(x)^{-\xi}) \setminus f(x)^\xi) \times g(x) &= f(x)^{-\xi} \times g(x) \\ (f(x) \times g(x^{\xi(p)}) \cup f(x) \times g(x)^{-\xi(p)}) \setminus (f(x) \times g(x)^{\xi(p)}) &= \\ f(x) \times ((g(x^{\xi(p)}) \cup g(x)^{-\xi(p)}) \setminus g(x)^{\xi(p)}) &= f(x) \times g(x)^{-\xi(p)} \end{aligned}$$

where the final equalities in these calculations are simply obtained by applying the operations $- \times g(y)$ and $f(x) \times -$ to the movement properties of the parity complex morphisms f and g at the elements $x \in C_p$ and $y \in D_q$ respectively. $\square$

Observation 236 ($\tilde{\Delta}$ as a functor). Suppose that $\alpha: [m] \longrightarrow [n]$ is a simplicial operator and $\underline{v}$ is an element of $\tilde{\Delta}[m]$ then we define $\alpha(\underline{v})$ of $\tilde{\Delta}[n]$ to be the element determined by the equation $\delta_{\alpha(\underline{v})}^n = (\alpha \circ \delta_{\underline{v}}^m)^f$. In other words the face operator corresponding to $\alpha(\underline{v})$ is obtained by composing the operators α and $\delta_{\underline{v}}^m$ and then applying the face-degeneracy factorisation of observation 8. In terms of our vectorial notation, this operation corresponds to applying α to the vertices of $\underline{v}$ to construct a, possibly repetitious, list of vertices $\alpha(v_0), \alpha(v_1), \dots, \alpha(v_r)$ from which we discard any repeats before re-indexing to obtain $\alpha(\underline{v})$. Our intuition is that $\alpha(\underline{v})$ is the face of $\tilde{\Delta}[n]$ which spans the set of vertices obtained by applying α to the vertices of the face $\underline{v}$ of $\tilde{\Delta}[m]$.

Notice that while this operation is nicely functorial, in the sense that $(\beta \circ \alpha)(\underline{v}) = \beta(\alpha(\underline{v}))$, it is not, in general, the case that it respects dimensions since the dimension of $\alpha(\underline{v})$ will be strictly less than that of $\underline{v}$ whenever $\alpha(v_i) = \alpha(v_{i+1})$ for some i . However, using this operation we may still define an arrow $\tilde{\Delta}(\alpha): \tilde{\Delta}[m] \longrightarrow \tilde{\Delta}[n]$ of Parity by letting:

$$\tilde{\Delta}(\alpha)(\underline{v}) = \begin{cases} \{\alpha(\underline{v})\} & \text{if the dimensions of } \underline{v} \text{ and } \alpha(\underline{v}) \text{ are the same,} \\ \emptyset & \text{otherwise.} \end{cases}$$

The verification of conditions (a) and (b) of lemma 231 for this function is a simple matter, the only interesting part of which is the demonstration that $\tilde{\Delta}(\alpha)(\underline{v})$ moves $\tilde{\Delta}(\alpha)(\underline{v}^-)$ to $\tilde{\Delta}(\alpha)(\underline{v}^+)$ for each element $\underline{v} \in \tilde{\Delta}[m]$. Assuming that $\underline{v}$ has dimension r we argue by cases:

- **$\alpha(\underline{v})$ has dimension r** , in which case it is easily seen that $\tilde{\Delta}(\alpha)(\underline{v}^\xi) = \alpha(\underline{v})^\xi$ (for either parity ξ), but we know that every element moves its set of negative faces to its set of positive faces so it is immediate that $\tilde{\Delta}(\alpha)(\underline{v}) = \{\alpha(\underline{v})\}$ has the required movement property.
- **$\alpha(\underline{v})$ has dimension $r - 1$** , in which case we have $\alpha(v_i) = \alpha(v_{i+1})$ for exactly one $0 \leq i < r$ and it follows that $\alpha(\underline{v} \cdot \delta_j^r)$ has dimension $r - 2$ for all $j \neq i, i + 1$ and that $\alpha(\underline{v} \cdot \delta_i^r) = \alpha(\underline{v} \cdot \delta_{i+1}^r) = \alpha(\underline{v})$. However the faces $\underline{v} \cdot \delta_i^r$ and $\underline{v} \cdot \delta_{i+1}^r$ of $\underline{v}$ are of opposite parity and so it follows that $\tilde{\Delta}(\alpha)(\underline{v}^-) = \tilde{\Delta}(\alpha)(\underline{v}^+) = \{\alpha(\underline{v})\}$ and it is clear that the set $\tilde{\Delta}(\alpha)(\underline{v})$ which is empty moves any set to itself as required.
- **$\alpha(\underline{v})$ has dimension $\leq r - 2$** , in which case each element $\alpha(\underline{v} \cdot \delta_i^r)$ must also have dimension $\leq r - 2$, so $\tilde{\Delta}(\alpha)(\underline{v}) = \tilde{\Delta}(\alpha)(\underline{v}^-) = \tilde{\Delta}(\alpha)(\underline{v}^+) = \emptyset$ and it follows that the movement property is trivial in this case.

It is also a matter of trivial verification to show that this action on simplicial operators preserves composition, giving a functor $\hat{\Delta}: \Delta \longrightarrow \underline{\text{Parity}}$.

11.2. Collapsers and Stratified Parity Complexes.

Observation 237 (collapsing and collapsers). Suppose that $\mathbb{C}$ is an ω -category and that $S = \bigcup_{n>0} S_n$ is a graded subset of its cells. Then we say that an ω -functor $f: \mathbb{C} \longrightarrow \mathbb{D}$ *collapses* (the cells in) S iff for each $n \in \mathbb{N}$ it maps each cell c in the subset S_{n+1} of $\mathbb{C}$ to an n -cell $f(c)$ in $\mathbb{D}$. Since $\omega\text{-}\underline{\text{Cat}}$ possesses all colimits, it follows that we may form a quotient ω -category $\mathbb{C}/S$ of $\mathbb{C}$ which represents the ω -functors that collapse S

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow{q_S} & \mathbb{C}/S \\ & \searrow f & \swarrow \exists! \hat{f} \\ & \mathbb{D} & \end{array}$$

in the sense that we have a quotient ω -functor q_S , as depicted in the diagram above, which collapses S and through which any other ω -functor f which collapses S factors uniquely. An ω -functor which possesses this universal property is said to be a *collapser* of (the cells in) S .

At a couple of places in the sequel, we will have use for a standard and very simple factorisation result with respect to these collapsers. Suppose that S and T are two graded subsets of the cells of $\mathbb{C}$ with $T \subseteq S$ then in the diagram

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow{q_T} & \mathbb{C}/T \\ & \searrow q_S & \swarrow \exists! q \\ & \mathbb{C}/S & \end{array}$$

the ω -functor q_S collapses the cells in T , since it collapses those in S and $T \subseteq S$, so it factors through q_T to give a ω -functor which is itself a collapser. To be precise, q is a collapser of the graded subset $q_T(S \setminus T) \subseteq \mathbb{C}/T$ given by $q_T(S \setminus T)_r = \{q_T(x) \mid x \in S_r \text{ and } x \notin T_r\}$ for each $r > 0$.

Definition 238 (stratified parity complexes). By analogy with stratified sets, we define a *stratified parity complex* to be a pair (C, tC) in which C is a parity complex and tC is a subset of its elements, called *thin elements*, all of which must be of dimension > 0 . A *stratified morphism* $f: (C, tC) \longrightarrow (D, tD)$ between such structures is simply a parity complex morphism $f: C \longrightarrow D$ satisfying the thinness preservation axiom that $f(tC) \subseteq tD$. Equivalently this axiom states that for all $x \in tC$ we have $y \in tD$ for all $y \in f(x)$.

Suppose that $g: (D, tD) \longrightarrow (E, tE)$ is another stratified morphism then their composite $g \circ f: C \longrightarrow E$ in $\underline{\text{Parity}}$ also satisfies the thinness preservation axiom (with respect to tC and tE) and so we may lift the composition of $\underline{\text{Parity}}$ to construct a category $\underline{\text{SParity}}$ of stratified parity complexes and stratified morphisms between them. Notice that we may identify $\underline{\text{Parity}}$ with the full subcategory of those stratified parity complexes in $\underline{\text{SParity}}$ whose set of thin elements is empty, which we shall do routinely in the sequel.

In general, when discussing stratified parity complexes we will adopt the same notational conventions that we have been applying to stratified sets, including dropping explicit mention of their sets of thin elements (cf. definition 96). Having done this, we also adopt $U: \underline{\text{SParity}} \longrightarrow \underline{\text{Parity}}$ to denote the functor which forgets stratifications and use this when

we wish to notationally distinguish between a stratified parity complex C and its underlying parity complex $U(C)$.

It should also be clear that most of the basic concepts, constructions and standard structures introduced for stratified sets may be naturally transferred to stratified parity complexes. We leave the details of this process up to the reader, since we will be content to restrict ourselves to the relatively limited collection of results and generalisations discussed in the next few

Observation 239 (extending $\mathcal{O}$ to $\underline{\text{SParity}}$). If C is a stratified parity complex then we define a graded set $\langle tC \rangle$ of cells of the freely generated ω -category $\mathcal{O}(U(C))$ by $\langle tC \rangle_r = \{\langle x \rangle \mid x \in tC \cap C_r\}$ ($r = 1, 2, \dots$) and let $\mathcal{O}(C)$ to be the collapsed ω -category $\mathcal{O}(U(C))/\langle tC \rangle$. We'll generally denote the associated collapser by $q_t: \mathcal{O}(U(C)) \twoheadrightarrow \mathcal{O}(C)$ and extend our atom notation by letting $\langle x \rangle \in \mathcal{O}(C)$ denote the cell obtained by applying this collapser to the atom in $\mathcal{O}(U(C))$ which corresponds to the element $x \in C$. Of course this collapser is an epimorphism, so we may apply lemma 221 to show that the set

$$\langle C \rangle \stackrel{\text{def}}{=} \bigcup_{n \in \mathbb{N}} \{\langle x \rangle \mid x \in C_n \text{ and } \langle x \rangle \text{ is not an } (n-1)\text{-cell in } \mathcal{O}(C)\}$$

is a weak generator of $\mathcal{O}(C)$.

To extend this construction to a functor on $\underline{\text{SParity}}$, suppose that we are given a stratified morphism $f: C \twoheadrightarrow D$ and consider the ω -functor $\mathcal{O}(f): \mathcal{O}(U(C)) \longrightarrow \mathcal{O}(U(D))$ between the free ω -categories on underlying parity complexes. If x is an n -element in C then $\langle x \rangle$ is of course a non-trivial n -cell in $\mathcal{O}(U(C))$ and we know that $\mathcal{O}(f)$ maps it to an n -cell (N, P) in $\mathcal{O}(U(D))$ which has $N_n = P_n = f(x)$. So applying Street's Excision of Extremals algorithm, theorem 4.1 of [33], as discussed in the proof of theorem 4.2 of that paper, we see that the cell (N, P) may be expressed as a composite of atoms on elements in the sub-parity complex of D generated by $N \cup P$. In particular, all atoms in this decomposition are n -cells and the only ones that are non-trivial are those which are atoms on an element in $f(x)$. Now if x is thin then the thinness preservation property of f tells us that $f(x)$ is a set of thin n -elements in D , so we know that for each $y \in f(x)$ the quotient functor $q_t: \mathcal{O}(U(D)) \longrightarrow \mathcal{O}(D)$ collapses the atom $\langle y \rangle$ to an $(n-1)$ -cell and therefore that it collapses each cell in the atomic decomposition of (N, P) , and thus this cell itself, to an $(n-1)$ -cell in $\mathcal{O}(D)$.

In other words, we have shown that the composite ω -functor $q_t \circ \mathcal{O}(f): \mathcal{O}(U(C)) \longrightarrow \mathcal{O}(D)$ collapses the cells in the graded set $\langle tC \rangle$. From this we can infer, by the universal property of the quotient $\mathcal{O}(D)$, that this composite induces the unique (dashed) ω -functor which makes the following square commute:

$$\begin{array}{ccc} \mathcal{O}(U(C)) & \xrightarrow{\mathcal{O}(f)} & \mathcal{O}(U(D)) \\ q_t \downarrow & & \downarrow q_t \\ \mathcal{O}(C) & \overset{\exists! \mathcal{O}(f)}{\dashrightarrow} & \mathcal{O}(D) \end{array}$$

So we define $\mathcal{O}(f): \mathcal{O}(C) \longrightarrow \mathcal{O}(D)$ to be this induced ω -functor and it follows, from its uniqueness property, the functoriality of $\mathcal{O}$ on $\underline{\text{Parity}}$ and the fact that stratified morphisms compose as in the underlying category $\underline{\text{Parity}}$, that this action is functorial with respect to the category structure on $\underline{\text{SParity}}$ as required.

Suppose now that D is an entire sub-complex of the stratified parity complex C , in other words these share the same underlying parity complex $U(C)$ and have $tD \subseteq tC$. Then $\mathcal{O}(C)$ and $\mathcal{O}(D)$ are constructed by taking collapsers of $\mathcal{O}(U(C))$ with regard to the sets $\langle tC \rangle$ and $\langle tD \rangle$ of cells respectively. The latter of these is a subset of the former, so we may apply the factorisation result of observation 237 to show that the induced ω -functor $\mathcal{O}(\subseteq_e): \mathcal{O}(D) \longrightarrow \mathcal{O}(C)$ is a collapser of the graded set given by $\langle tC \setminus tD \rangle_r = \{\langle x \rangle \mid x \in tC \cap C_n \text{ and } x \notin tD\}$.

If we consider Parity to be the full subcategory of trivially stratified parity complexes in SParity, as mentioned in the last observation, then the functor $\mathcal{O}$ on SParity is indeed an extension of our original free ω -category functor on Parity.

Observation 240. Suppose that N is a stratified set in the full subcategory Simplex of definition 104, which according to convention has underlying simplicial set $\Delta[n]$. Then we may define a corresponding stratified parity complex $\tilde{N}$ which has underlying parity complex $\tilde{\Delta}[n]$ and in which an element $\underline{v}$ is thin iff the corresponding face $\delta_{\underline{v}}^n \in \Delta[n]$ is thin in N .

Furthermore, if $f: N \longrightarrow M$ is a stratified map in Simplex then we may apply the simplicial Yoneda lemma to show that there is a unique simplicial operator $\alpha: [n] \longrightarrow [m]$ for which $\Delta(\alpha): \Delta[n] \longrightarrow \Delta[m]$ is the simplicial map underlying f . Consider $\tilde{\Delta}(\alpha): \tilde{\Delta}[n] \longrightarrow \tilde{\Delta}[m]$ the corresponding parity complex morphism and recall that, from the definition given in observation 236, $\tilde{\Delta}(\alpha)(\underline{v}) = \{\underline{w}\}$ if and only if $\delta_{\underline{w}}^m = \alpha \circ \delta_{\underline{v}}^n = \Delta(\alpha)(\delta_{\underline{v}}^n)$ and that otherwise $\tilde{\Delta}(\alpha)(\underline{v}) = \emptyset$. In the former case, if $\underline{v}$ is thin in $\tilde{N}$ then we know, by definition, that $\delta_{\underline{v}}^n$ is thin in N and thus that $\Delta(\alpha)(\delta_{\underline{v}}^n) = \delta_{\underline{w}}^m$ is thin in M , since $\Delta(\alpha)$ underlies the stratified map f , so finally we see that $\underline{w}$ is thin in $\tilde{M}$. It follows that $\tilde{\Delta}(\alpha)$ provides us with a stratified parity complex morphism $\tilde{f}: \tilde{N} \longrightarrow \tilde{M}$ and that this construction enriches our operation on the objects of Simplex to a functor from that category into SParity.

Observation 241 (collapsed orientals). Of course, if we consider $t\Delta$ to be a full subcategory of Strat then it is, in fact, a subcategory of Simplex and it follows that we may restrict the functor of the last observation to a functor $\Delta: t\Delta \longrightarrow \text{SParity}$. This functor extends the functor $\tilde{\Delta}: \Delta \longrightarrow \text{Parity}$ of observation 236 by mapping $[n]_t$ to the stratified parity complex $\tilde{\Delta}[n]_t$ obtained from $\tilde{\Delta}[n]$ by making thin its unique n -dimensional element $01\dots n$.

Now, we've seen that the atom $\langle 01\dots n \rangle$ is the only non-trivial n -cell of the oriental $\mathcal{O}(\tilde{\Delta}[n])$ so it is clear that $\mathcal{O}(\tilde{\Delta}[n]_t)$ is an $(n-1)$ -category, since it is obtained by collapsing this unique n -cell. We call $\mathcal{O}(\tilde{\Delta}[n]_t)$ the n^{th} collapsed oriental. Our intuition here is that this is the appropriate ω -categorical model of the standard thin n -simplex.

Observation 242 (extending a few other constructions to SParity). If C and D are stratified parity complexes then their product is obtained by taking the underlying product of parity complexes, just as in observation 228, and defining its set of thin elements to be:

$$t(C \times D) \stackrel{\text{def}}{=} \{(x, y) \in C \times D \mid \text{either } x \in tC \text{ or } y \in tD\}$$

Furthermore, if $f: C \longrightarrow E$ and $g: D \longrightarrow F$ are stratified morphisms it is a matter of routine verification to check that their product as parity complex morphisms satisfies the thinness preservation axiom and thus that it provides a stratified morphism $f \times g: C \times D \longrightarrow E \times F$. It follows that we may canonically extend the product bifunctor on Parity to a bifunctor $\times: \text{SParity} \times \text{SParity} \longrightarrow \text{SParity}$ simply by lifting its action on underlying parity complex morphisms.

Another construction on stratified parity complexes that we shall have use for in the sequel is defined by analogy to the functors $\text{Th}_n: \underline{\text{Strat}} \longrightarrow \underline{\text{Strat}}$ introduced in definition 112. If C is a stratified parity complex and $n \in \mathbb{N}$ then $\text{Th}_n(C)$ is defined to be the stratified parity complex obtained from C by making thin all elements of dimension greater than n . Of course, it is clear that any stratified parity complex morphism $f: C \longrightarrow D$ lifts to a stratified morphism from $\text{Th}_n(C)$ to $\text{Th}_n(D)$, which we call $\text{Th}_n(f)$, making Th_n into an endo-functor on $\underline{\text{SParity}}$. The identity morphism on the underlying parity complex of C provides us with a canonical stratified morphism $i_C^n: C \longrightarrow \text{Th}_n(C)$ and the corresponding ω -functor $\mathcal{O}(i_C^n): \mathcal{O}(C) \longrightarrow \mathcal{O}(\text{Th}_n(C))$ is the collapser of the graded set of atoms of C on elements of dimension greater than n .

11.3. ω -Categorical Nerve Constructions.

Observation 243. Composing the functors $\tilde{\Delta}: \Delta \longrightarrow \underline{\text{Parity}}$ and $\mathcal{O}: \underline{\text{Parity}} \longrightarrow \omega\text{-Cat}$ we obtain a functor which represents the category of simplicial operators as a subcategory of $\omega\text{-Cat}$. Now, if $\alpha: [m] \longrightarrow [n]$ is a simplicial operator, then the corresponding ω -functor $\mathcal{O}(\tilde{\Delta}(\alpha)): \mathcal{O}(\tilde{\Delta}[m]) \longrightarrow \mathcal{O}(\tilde{\Delta}[n])$ is completely determined by its action on atoms, since they freely generate $\mathcal{O}(\tilde{\Delta}[m])$. In the sequel, it will be useful to have a more explicit description of this action on atoms.

Indeed, it is the case that $\mathcal{O}(\tilde{\Delta}(\alpha))$ is the unique ω -functor which maps each atom $\langle \underline{v} \rangle \in \mathcal{O}(\tilde{\Delta}[m])$ to the atom $\langle \alpha(\underline{v}) \rangle \in \mathcal{O}(\tilde{\Delta}[n])$. Unfortunately, this fact does not follow immediately from our argument so far, but it can be established by a somewhat tedious direct calculation. However we choose to take a different route and use an argument based on corollary 234. So let $E = \{ \underline{w} \in \tilde{\Delta}[m] \mid \underline{w} \subseteq \underline{v} \}$ be the sub-parity complex of $\tilde{\Delta}[m]$ generated by $\underline{v}$ and notice that if $\underline{w} \subseteq \underline{v}$ then $\alpha(\underline{w}) \subseteq \alpha(\underline{v})$ so it is clear that α maps elements of E into the sub-parity complex $F = \{ \underline{u} \in \tilde{\Delta}[n] \mid \underline{u} \subseteq \alpha(\underline{v}) \}$ generated by $\alpha(\underline{v})$ in $\tilde{\Delta}[n]$ and it follows that $\tilde{\Delta}(\alpha)$ restricts to a parity complex morphism from E to F . Now suppose that $\underline{u} = u_0 u_1 \dots u_r \in F_r$ (for some dimension r) then from the definition of F we know that the vertices of $\underline{u}$ are all vertices of $\alpha(\underline{v})$ and are thus of the form $\alpha(v_j)$ for some j . Consequently we may construct an element $\underline{w} \in E_r$ by letting $w_i = \min\{v_j \mid j \in [m] \text{ and } \alpha(v_j) = u_i\}$ for which we know, by construction, that $\alpha(\underline{w}) = \underline{u}$. It follows that we have $\tilde{\Delta}(\alpha)(\underline{w}) = \{\underline{u}\}$ and so $\underline{u} \in \tilde{\Delta}(\alpha)(E_r)$, or in other words we have succeeded in showing that the subset $\tilde{\Delta}(\alpha)(E)$ is actually equal to F itself. Now we may apply corollary 234 to conclude that $\tilde{\Delta}(\alpha)(\langle \underline{v} \rangle) = \langle \alpha(\underline{v}) \rangle$ as postulated.

We can also go further and extend this analysis to free ω -categories of the form $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$, which we will study in some detail in the next subsection. In this case if $\phi: [n] \longrightarrow [r]$ and $\psi: [m] \longrightarrow [s]$ are a pair of simplicial operators then we may easily adapt the argument of the last paragraph to show that if $(\underline{v}, \underline{w}) \in \tilde{\Delta}[n] \times \tilde{\Delta}[m]$ then $\mathcal{O}(\tilde{\Delta}(\phi) \times \tilde{\Delta}(\psi))(\langle \underline{v}, \underline{w} \rangle) = \langle \phi(\underline{v}), \psi(\underline{w}) \rangle$.

Observation 244 (Street's ω -categorical nerve). Our primary reason for extending $\tilde{\Delta}$ to a functor $\Delta \longrightarrow \underline{\text{Parity}}$ and forming the composite functor $\mathcal{O} \circ \tilde{\Delta}: \Delta \longrightarrow \omega\text{-Cat}$ is that we may now follow Street [31] and define an adjoint pair

$$\omega\text{-Cat} \begin{array}{c} \xleftarrow{\quad F_\omega \quad} \\ \perp \\ \xrightarrow{\quad N_\omega \quad} \end{array} \underline{\text{Simp}} \quad (63)$$

by Kan's construction (observation 63).

In other words, the right adjoint N_ω is constructed by “homming out” of the functor $\mathcal{O} \circ \tilde{\Delta}$, that is to say if $\mathbb{C}$ is an ω -category then the simplicial set $N_\omega(\mathbb{C})$ has $N_\omega(\mathbb{C})_n \stackrel{\text{def}}{=} \omega\text{-Cat}(\mathcal{O}(\tilde{\Delta}[n]), \mathbb{C})$ as its set of n -simplices and a right action given by the pre-composite $x \cdot \alpha \stackrel{\text{def}}{=} x \circ \mathcal{O}(\tilde{\Delta}(\alpha))$ where $x \in N_\omega(\mathbb{C})_n$ is an n -simplex and $\alpha: [m] \longrightarrow [n]$ is a compatible simplicial operator.

The left adjoint F_ω may be constructed as the left Kan extension of the composite functor $\mathcal{O} \circ \tilde{\Delta}: \Delta \longrightarrow \omega\text{-Cat}$ along the Yoneda embedding $\Delta: \Delta \longrightarrow \mathbf{Simp}$, that is we can write $F_\omega(X)$ as the weighted colimit $\text{colim}(X, \mathcal{O} \circ \tilde{\Delta})$ in $\omega\text{-Cat}$. A cocone under the diagram $\mathcal{O} \circ \tilde{\Delta}$ weighted by X is simply a family of ω -functors $f_x: \mathcal{O}(\tilde{\Delta}[n]) \longrightarrow \mathbb{C}$ for $x \in X$ (where $n = \dim(x)$) satisfying the naturality condition that for each simplicial operator $\alpha: [m] \longrightarrow [n]$ which is compatible with $x \in X$ we have $f_{x \cdot \alpha} = f_x \circ \mathcal{O}(\tilde{\Delta}(\alpha))$.

In future calculations we’ll use the notation $i_x: \mathcal{O}(\tilde{\Delta}[n]) \longrightarrow F_\omega(X)$ to denote the components of the colimiting cocone. Of course any other such cocone f_x induces a ω -functor $f: F_\omega(X) \longrightarrow \mathbb{C}$ which is uniquely determined by the equality $f \circ i_x = f_x$ for each $x \in X$. In particular, if $f: X \longrightarrow Y$ is a simplicial map then the ω -functor $F_\omega(f): F_\omega(X) \longrightarrow F_\omega(Y)$ is characterised by the fact that it is the unique ω -functor which makes the triangle

$$\begin{array}{ccc} & \mathcal{O}(\tilde{\Delta}[n]) & \\ i_x \swarrow & & \searrow i_{f(x)} \\ F_\omega(X) & \xrightarrow{F_\omega(f)} & F_\omega(Y) \end{array} \quad (64)$$

commute for each simplex $x \in X$ (where $n = \dim(x)$).

Observation 245 ($F_\omega(X)$ as a (freely) generated ω -category). Suppose that x is an n -simplex in X then we adopt the notation $\llbracket x \rrbracket$ to denote the cell $i_x(\langle 01\dots n \rangle) \in F_\omega(X)$. At the risk of confusing our nomenclature a little we’ll tend to refer to $\llbracket x \rrbracket$ as the *atom of $F_\omega(X)$ associated with the simplex x* .

Notice that if $\alpha: [m] \longrightarrow [n]$ is a simplicial operator then we have

$$i_x(\langle \alpha(01\dots m) \rangle) = (i_x \circ \mathcal{O}(\tilde{\Delta}(\alpha)))(\langle 01\dots m \rangle) = i_{x \cdot \alpha}(\langle 01\dots m \rangle) \stackrel{\text{def}}{=} \llbracket x \cdot \alpha \rrbracket \quad (65)$$

where the first equality follows by observation 243 and the second expresses the naturality of the cocone i_x . As a consequence if $\alpha: [m] \longrightarrow [n]$ is a face operator then

$$\llbracket x \cdot \alpha \rrbracket = i_x(\langle \alpha(01\dots m) \rangle) = i_x(\langle 01\dots n \rangle) \stackrel{\text{def}}{=} \llbracket x \rrbracket$$

where the second of these equalities holds because α is surjective and thus has $\alpha(01\dots m) = 01\dots n$.

Conversely, we may show that whenever x is a non-degenerate simplex of X then $\llbracket x \rrbracket$ is actually a non-trivial n -cell of $F_\omega(X)$. This can be done by employing the explicit description of $F_\omega(X)$ as a colimit to construct an ω -functor k_X from there into $\mathcal{N}(\tilde{X})$, the ω -category constructed from the graded set $\tilde{X}$ of non-degenerate simplices in X as in observation 217, which maps each atom $\llbracket x \rrbracket$ to a cell $(M, P) \in \mathcal{N}(\tilde{X})$ defined by:

$$M = \{x \cdot \delta_{\underline{v}}^n \mid \underline{v} \in \mu(01\dots n)\} \cap \tilde{X} \quad P = \{x \cdot \delta_{\underline{v}}^n \mid \underline{v} \in \pi(01\dots n)\} \cap \tilde{X}$$

Then, in the case where x is non-degenerate we see that $M_n = P_n = \{x\}$ so it follows that $k_X(\llbracket x \rrbracket)$ is a non-trivial n -cell in $\mathcal{N}(\tilde{X})$ and thus (cf. observation 84) that $\llbracket x \rrbracket$ itself is a non-trivial n -cell in $F_\omega(X)$ as required. We leave the detailed verification of this result as an exercise for the reader.

Indeed we may actually show that $F_\omega(X)$ is freely generated by its set of atoms:

$$\llbracket X \rrbracket \stackrel{\text{def}}{=} \{\llbracket x \rrbracket \mid x \in X\} = \{\llbracket x \rrbracket \mid x \text{ is a non degenerate simplex of } X\}$$

However this result will not be required here and so we leave its (relatively routine) proof as an exercise for the reader. Instead we will content ourselves with simply showing that $\llbracket X \rrbracket$ weakly generates $F_\omega(X)$. To do so we apply lemma 221, via the associated observation 222, to the jointly epimorphic family $i_x: \mathcal{O}(\tilde{\Delta}[n]) \longrightarrow F_\omega(X)$ of colimiting cocone components. Now if $\underline{v}$ is an m -dimensional element of $\tilde{\Delta}[n]$ we have $\underline{v} = \delta_{\underline{v}}^n(01\dots m)$ so it follows, from equation (65), that $i_x(\langle \underline{v} \rangle) = \llbracket x \cdot \delta_{\underline{v}}^n \rrbracket$. In other words, each ω -functor i_x maps the generators (atoms) of its domain $\mathcal{O}(\tilde{\Delta}[n])$ into $\llbracket X \rrbracket$, therefore the conclusion of lemma 221 is that this set weakly generates $F_\omega(X)$ as required.

In particular, it follows that if $f: X \longrightarrow Y$ is a simplicial map then the characterisation in display (64) can be re-expressed as stating that $F_\omega(f): F_\omega(X) \longrightarrow F_\omega(Y)$ is the unique ω -functor which is determined by the fact that for each non-degenerate simplex $x \in X$ it maps the cell $\llbracket x \rrbracket$ in $F_\omega(X)$ to the cell $\llbracket f(x) \rrbracket$ in $F_\omega(Y)$.

Observation 246. It turns out, however, that this nerve construction discards a little too much information to make it a useful representation of ω -categories as simplicial structures. To be precise, we do not preserve enough about the identities in the ω -categories we are representing and, consequently, there is no hope that we might be able to re-construct the compositional structures of an ω -category from its simplicial nerve. In [31] Street follows the lead set by Roberts in [26] and [27] and suggests that the appropriate approach to rectifying this deficiency would be to extend the nerve construction to the category of stratified sets

$$\omega\text{-}\underline{\text{Cat}} \begin{array}{c} \xrightarrow{\quad F_\omega \quad} \\ \perp \\ \xrightarrow{\quad N_\omega \quad} \end{array} \underline{\text{Strat}} \quad (66)$$

in which we can use appropriately selected sets of thin simplices to preserve enough information about identities between the composites that form their sources and targets.

To make this construction precise, we proceed as in observation 244 by forming the composite of the extended functors $\tilde{\Delta}: t\Delta \longrightarrow \underline{\text{SParity}}$ of observation 241 and $\mathcal{O}: \underline{\text{SParity}} \longrightarrow \omega\text{-}\underline{\text{Cat}}$ of observation 239. This composite extends $\mathcal{O} \circ \tilde{\Delta}: \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$ by mapping the thin simplex $[n]_t$ to the collapsed oriental $\mathcal{O}(\tilde{\Delta}[n]_t)$ and the operator $\varrho^n: [n] \longrightarrow [n]_t$ to the canonical collapser $q: \mathcal{O}(\tilde{\Delta}[n]) \longrightarrow \mathcal{O}(\tilde{\Delta}[n]_t)$. Of course, each of these quotient maps is an epimorphism and so it follows that the composite $\mathcal{O} \circ \tilde{\Delta}: t\Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$ is actually a $\mathbb{T}_{\underline{\text{Strat}}}$ -coalgebra. It follows that we may construct the adjunction shown in display (66) by applying Kan's construction, observation 63, to this coalgebra.

More explicitly, we may form the stratified nerve $N_\omega(\mathbb{C}) \in \underline{\text{Strat}}$ of an ω -category $\mathbb{C}$ by first forming its nerve in $\underline{\text{Simp}}$ and then making thin any n -simplex $x \in N_\omega(\mathbb{C})$ for which the corresponding ω -functor $x: \mathcal{O}(\tilde{\Delta}[n]) \longrightarrow \mathbb{C}$ carries the atom $\langle 01\dots n \rangle$ to an $(n-1)$ -cell in $\mathbb{C}$. Going in the other direction, to form $F_\omega(X)$ for a stratified set X we first apply the left adjoint of display (63) to the underlying simplicial set $U(X) \in \underline{\text{Simp}}$ to give an ω -category

$F_\omega(U(X))$ then we define a graded subset of the cells of $F_\omega(U(X))$ by

$$\llbracket tX \rrbracket_n \stackrel{\text{def}}{=} \{\llbracket x \rrbracket \mid x \in X \text{ is a non-degenerate, thin } n\text{-simplex}\}$$

and let $F_\omega(X)$ be the quotient $F_\omega(U(X))/\llbracket tX \rrbracket$ of observation 237. In other words, we may construct $F_\omega(U(X))$ as a colimit of orientals and then collapse those (images of) orientals in there which correspond to thin simplices in X . We'll generally denote the associated collapser by $q_t: F_\omega(U(X)) \longrightarrow F_\omega(X)$ and extend our atom notation by letting $\llbracket x \rrbracket \in F_\omega(X)$ denote the n -cell obtained by applying this collapser to the corresponding atom in $F_\omega(U(X))$ associated with the n -simplex $x \in X$.

Of course, the collapser $q_t: F_\omega(U(X)) \longrightarrow F_\omega(X)$ is an epimorphism and it carries the atom $\llbracket x \rrbracket \in F_\omega(U(X))$ associated with any thin n -simplex to an $(n-1)$ -cell in $F_\omega(X)$. Therefore we may apply lemma 221 to the weak generating set $\llbracket U(X) \rrbracket \subseteq F_\omega(U(X))$ to show that the set

$$\llbracket X \rrbracket \stackrel{\text{def}}{=} \bigcup_{n \in \mathbb{N}} \{\llbracket x \rrbracket \mid x \in X_n \text{ and } \llbracket x \rrbracket \text{ is not an } (n-1)\text{-cell in } F_\omega(X)\} \quad (67)$$

weakly generates the ω -category $F_\omega(X)$. Again, it follows that if $f: X \longrightarrow Y$ is a stratified map then the ω -functor $F_\omega(f): F_\omega(X) \longrightarrow F_\omega(Y)$ is completely characterised by the fact that for each simplex $x \in X$ it maps the cell $\llbracket x \rrbracket \in F_\omega(X)$ to the cell $\llbracket f(x) \rrbracket \in F_\omega(Y)$.

Suppose that Y is an entire subset of the stratified set X , in other words these stratified sets share the same underlying simplicial set and $tY \subseteq tX$. Then $F_\omega(X)$ and $F_\omega(Y)$ are constructed by taking collapsers of $F_\omega(U(X))$ with regard to the sets $\llbracket tX \rrbracket$ and $\llbracket tY \rrbracket$ of cells respectively. The latter of these is a subset of the former, so we may apply the factorisation result of observation 237 to show that the induced ω -functor $F_\omega(\subseteq_e): F_\omega(Y) \longrightarrow F_\omega(X)$ is a collapser of the graded set given by $\llbracket tX \setminus tY \rrbracket_r = \{\llbracket x \rrbracket \mid x \in tX \cap X_n \text{ and } x \notin tY\}$.

It might appear that we have introduced some ambiguity in our notation by using F_ω to denote the left adjoints from Simp and Strat into $\omega\text{-Cat}$ and at the same time identifying the category Simp with the full subcategory of Strat on those objects which are minimally stratified. However, we know that each thin simplex of a minimally stratified set X is degenerate, therefore $\llbracket tX \rrbracket = \emptyset$ and we see that, in this case, the quotient $F_\omega(X)$ is actually (isomorphic to) $F_\omega(U(X))$ itself.

Lemma 247. *For each $n \in \mathbb{N}$ the functors N_ω and F_ω restrict to give an adjunction*

$$\begin{array}{ccc} & F_n & \\ & \curvearrowright & \\ n\text{-Cat} & \xrightleftharpoons[N_n]{\perp} & \text{Strat}_n \end{array} \quad (68)$$

between the categories of n -categories and n -trivial stratified sets.

Proof. If $\mathbb{C}$ is an n -category and $r > n$ then an r -simplex of $N_\omega(\mathbb{C})$ is simply an ω -functor $x: \mathcal{O}(\tilde{\Delta}[r]) \longrightarrow \mathbb{C}$. However all cells in $\mathbb{C}$ are n -cells and so it follows that any such ω -functor must collapse the oriental that is its domain and thus, by definition, that our r -simplex is thin in $N_\omega(\mathbb{C})$. In other words, we see easily that $N_\omega(\mathbb{C})$ is indeed n -trivial as stated.

To prove that F_ω also restricts as described, consider an n -trivial stratified set X and recall that we know, from observation 108, that it admits a canonical representation as the colimit of a diagram $D_X: \mathbb{G}(X) \longrightarrow \text{Strat}$ of standard (thin) simplices. Furthermore the left adjoint Th_n to $|\cdot|_n$ preserves this colimit and we know that $\text{Th}_n(X) = X$, since in any n -trivial stratified set every simplex of dimension greater than n is already thin, so it follows that we may express X as a colimit of stratified sets of the form $\text{Th}_n(\Delta[r]_?)$. Applying the left adjoint

functor F_ω to this we see that $F_\omega(X)$ may be constructed as a colimit of ω -categories of the form $F_\omega(\text{Th}_n(\Delta[r]_?))$.

Now $F_\omega(\text{Th}_n(\Delta[r]_?))$ is constructed from $F_\omega(\Delta[r])$, the freely generated ω -category $\mathcal{O}(\tilde{\Delta}[r])$, by collapsing a graded set of cells which contains all atoms on elements of $\tilde{\Delta}[r]$ with dimension greater than n . However any r -cell in $\mathcal{O}(\tilde{\Delta}[r])$ may be expressed as a composite of atoms and all atoms of dimension r greater than n have been collapsed to $(r-1)$ -cells in $F_\omega(\text{Th}_n(\Delta[r]_?))$ so we may show, by a simple inductive proof on superstructures, that $F_\omega(\text{Th}_n(\Delta[r]_?))$ is in fact an n -category. Finally we know that $n\text{-}\underline{\text{Cat}}$ is closed in $\omega\text{-}\underline{\text{Cat}}$ under colimits so it follows that $F_\omega(X)$, as a colimit of n -categories, is actually an n -category itself as required. $\square$

Observation 248. We might also expect that a strong relationship should hold between $F_\omega(|X|_n)$, the ω -category associated with the n -superstructure of a stratified set X , and the n -superstructure $|F_\omega(X)|_n$. To construct a comparison between these two ω -categories, we apply F_ω to the inclusion $|X|_n \xrightarrow{\subseteq_r} X$ to obtain an ω -functor $F_\omega(\subseteq_r): F_\omega(|X|_n) \longrightarrow F_\omega(X)$ which is characterised by the fact that for each simplex $x \in |X|_n$ it maps the atom $\llbracket x \rrbracket \in F_\omega(|X|_n)$ to the corresponding atom $\llbracket x \rrbracket \in F_\omega(X)$. Now applying lemma 247 we find that $F_\omega(|X|_n)$ is an n -category and therefore that we may restrict our ω -functor to a canonical comparison $F_\omega(|X|_n) \longrightarrow |F_\omega(X)|_n$.

Indeed, this comparison ω -functor can be shown to be an isomorphism. However, a direct proof of this fact is a little involved and for our purposes here all we need is the observation that it is an epimorphism. To prove this, start with the fact that the set $\llbracket X \rrbracket$ of observation 246 weakly generates $F_\omega(X)$, from which we may infer immediately that its subset $|F_\omega(X)|_n \cap \llbracket X \rrbracket$ of n -cells weakly generates the superstructure $|F_\omega(X)|_n$. By the definition of $\llbracket X \rrbracket$ given in display (67) each cell in $|F_\omega(X)|_n \cap \llbracket X \rrbracket$ is an atom $\llbracket x \rrbracket \in |F_\omega(X)|_n$ on some non-thin r -simplex $x \in X$ with $r \leq n$, furthermore each such simplex is in the regular subset $|X|_n \subseteq_r X$ and thus provides an atom $\llbracket x \rrbracket \in F_\omega(|X|_n)$ which maps to $\llbracket x \rrbracket \in |F_\omega(X)|_n$ under our comparison ω -functor. In other words every one of the cells in our weak generator $|F_\omega(X)|_n \cap \llbracket X \rrbracket$ for $|F_\omega(X)|_n$ is mapped to by some cell in $F_\omega(|X|_n)$ under the comparison $F_\omega(|X|_n) \longrightarrow |F_\omega(X)|_n$, which is therefore an epimorphism by observation 222.

Theorem 249 (Street [32]). *The ω -categorical nerve functor $N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Strat}}$ takes each ω -category to a complicial set. It follows that the left adjoint $F_\omega: \underline{\text{Strat}} \longrightarrow \omega\text{-}\underline{\text{Cat}}$ carries f -extensions to isomorphisms and that we may restrict the adjunctions of displays (66) and (68) to:*

$$\begin{array}{ccc} \omega\text{-}\underline{\text{Cat}} & \xrightleftharpoons[\text{N}_\omega]{F_\omega} & \underline{\text{Cs}} \end{array} \quad \text{and} \quad \begin{array}{ccc} n\text{-}\underline{\text{Cat}} & \xrightleftharpoons[\text{N}_n]{F_n} & \underline{\text{Cs}}_n \end{array}$$

Proof. For a proof of this result see loc. cit., which essentially proceeds by establishing an optimal decomposition result for the cells of an oriental.

It is worth observing, however, that the result Street actually proves there looks, on the face of it, to be a little stronger than our compliciality condition. In particular his *admissible horn* notion is somewhat more liberal than the one we gave in notation 105, since it requires fewer faces to be thin for admissibility to hold. In turn, this means that his complicial set notion explicitly insists that a strictly greater number of horns should have unique fillers. It follows therefore that Street's result, which establishes compliciality in his apparently stronger sense, immediately implies compliciality in our sense which is actually the notion originally proposed by Roberts in [27].

It should be noted that there is no a priori reason to believe that Street's compliciality notion isn't strictly stronger than ours. However our slightly weaker notion is adequate for all of our work in earlier sections and its adoption frees us from having to verify Street's more complex admissibility condition in our calculations.

Indeed the equivalence of these notions is actually a simple corollary of the ultimate result in this work, which establishes Street's primary conjecture in [31] that the nerve functor actually provides us with a canonical equivalence between the categories $\omega\text{-}\underline{\text{Cat}}$ of ω -categories and $\underline{\text{Cs}}$ of complicial sets. $\square$

Observation 250 (N_ω as a finitely accessible functor). It is, of course, the case that $\omega\text{-}\underline{\text{Cat}}$ is an LFP-category. Limits and filtered colimits are formed in $\omega\text{-}\underline{\text{Cat}}$ in the familiar way for such categories, that is we take the limit or filtered colimit of the associated diagram of underlying sets in $\underline{\text{Set}}$ and then enrich that with an ω -category structure derived point-wise from those of the nodes of our original diagram. All finite ω -categories and quotients thereof are finitely presentable so, in particular, the orientals $\mathcal{O}(\tilde{\Delta}[n])$ and collapsed orientals $\mathcal{O}(\tilde{\Delta}[n])_t$ are all finitely presentable. It follows that the coalgebra $\mathcal{O} \circ \tilde{\Delta}: t\Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$, which we used in observation 246 to define $F_\omega \dashv N_\omega$ via Kan's construction, is finitely presented and thus that $N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Strat}}$ preserves all filtered colimits (cf. observation 63).

11.4. Relating Parity Complex Products and the Tensor Product of Stratified Sets. Earlier, we claimed in passing that the tensor product of complicial sets was the appropriate generalisation of Gray's lax tensor product [15] to the category of complicial sets. The main result of this subsection gives this bold claim validity by demonstrating a strong relationship between our tensor product and the free ω -categories generated by products of parity complexes.

Lemma 251. *We may define a canonical parity complex morphism $\nabla: \tilde{\Delta} \longrightarrow \tilde{\Delta} \times \tilde{\Delta}$ which maps each r -simplex $\underline{v}$ to the subset:*

$$\nabla(\underline{v}) \stackrel{\text{def}}{=} \left\{ (\underline{v} \cdot \mathbb{1}_1^{\vec{r}}, \underline{v} \cdot \mathbb{1}_2^{\vec{r}}) \mid \vec{r} \text{ is a partition of } r \right\}$$

Alternatively, if we express $\underline{v}$ using the vertex-wise notation $v_0 \dots v_r$ then we can write this as:

$$\nabla(v_0 \dots v_r) \stackrel{\text{def}}{=} \left\{ (v_0 \dots v_s, v_s \dots v_r) \in \tilde{\Delta}_s \times \tilde{\Delta}_{r-s} \mid s = 0, 1, \dots, r \right\}$$

Proof. Notice first that ∇ is indeed a morphism of graded sets, since for each $s = 0, 1, \dots, r$ we know that $(v_0 \dots v_s, v_s \dots v_r)$ is an element of $\tilde{\Delta}_s \times \tilde{\Delta}_{r-s}$ which is subset of $(\tilde{\Delta} \times \tilde{\Delta})_r \stackrel{\text{def}}{=} \bigcup_{i=0}^r \tilde{\Delta}_i \times \tilde{\Delta}_{r-i}$ and so it follows that $\nabla(v_0 \dots v_r) \subseteq (\tilde{\Delta} \times \tilde{\Delta})_r$. Furthermore, we have $\nabla(v_0) = \{(v_0, v_0)\}$ and so condition (a) of lemma 231 holds for our morphism and it remains to prove that condition (b) of that lemma also holds.

To that end, we start by working out what the sets of faces of an element $(v_0 \dots v_s, v_s \dots v_r)$ of $\nabla(v_0 \dots v_r)$ look like. Consulting the formulae for the sets of faces of simplices and products we see that its faces of parity ξ are $(v_0 \dots \hat{v}_i \dots v_s, v_s \dots v_r)$ if i is of parity ξ and $(v_0 \dots v_s, v_s \dots \hat{v}_i \dots v_r)$ if $(i - s)$ is of parity $\xi(s)$. However in the latter case it is clear that this condition also reduces to simply insisting that i is of parity ξ , therefore we have:

$$\begin{aligned} (v_0 \dots v_s, v_s \dots v_r)^\xi = & \{ (v_0 \dots \hat{v}_i \dots v_s, v_s \dots v_r) \mid 0 < s \text{ and } 0 \leq i \leq s \text{ and } i \text{ is of parity } \xi \} \cup \\ & \{ (v_0 \dots v_s, v_s \dots \hat{v}_i \dots v_r) \mid s < r \text{ and } s \leq i \leq r \text{ and } i \text{ is of parity } \xi \} \end{aligned} \quad (69)$$

Notice also that the first factor in this union is a subset of $\tilde{\Delta}_{s-1} \times \tilde{\Delta}_{r-s}$ and that its second factor is a subset of the disjoint set $\tilde{\Delta}_s \times \tilde{\Delta}_{r-s-1}$.

So suppose that we have a distinct pair of indices $s, t \in [r]$, assume w.l.o.g. that $0 \leq s < t \leq r$ and consider the intersection $(v_0 \dots v_s, v_s \dots v_r)^\xi \cap (v_0 \dots v_t, v_t \dots v_r)^\varphi$. We know that the first of the factors here is a subset of $(\tilde{\Delta}_{s-1} \times \tilde{\Delta}_{r-s}) \cup (\tilde{\Delta}_s \times \tilde{\Delta}_{r-s-1})$ and that its second is a subset of $(\tilde{\Delta}_{t-1} \times \tilde{\Delta}_{r-t}) \cup (\tilde{\Delta}_t \times \tilde{\Delta}_{r-t-1})$ therefore it follows, by the pairwise disjointness of the sets $\tilde{\Delta}_i \times \tilde{\Delta}_{r-i-1}$ for $i = 0, 1, \dots, r-1$, that our intersection is empty if $t > s+1$ and in the case where $t = s+1$ we have:

$$(v_0 \dots v_s, v_s \dots v_r)^\xi \cap (v_0 \dots v_{s+1}, v_{s+1} \dots v_r)^\varphi = \{(v_0 \dots v_s, v_s \dots \hat{v}_i \dots v_r) \mid i \text{ is of parity } \xi\} \cap \{(v_0 \dots \hat{v}_j \dots v_{s+1}, v_{s+1} \dots v_r) \mid j \text{ is of parity } \varphi\}$$

Notice, however, that those elements in the first factor which have $i > s$ cannot occur in the second factor, since all elements in there mention the vertex v_i in their second ordinate. Dually any element in the second factor with $j < s+1$ cannot also occur in the first factor. Thus it follows easily that the only possible element in this intersection must be obtained by dropping v_s in the first factor and v_{s+1} in the second factor and thus it is clear that we have:

$$(v_0 \dots v_s, v_s \dots v_r)^\xi \cap (v_0 \dots v_{s+1}, v_{s+1} \dots v_r)^\varphi = \begin{cases} \{(v_0 \dots v_s, v_{s+1} \dots v_r)\} & s \text{ is of parity } \xi \\ & \text{and } \varphi = \neg\xi, \\ \emptyset & \text{otherwise.} \end{cases}$$

From this calculation it immediately follows that $\nabla(v_0 \dots v_r)$ is a well formed subset of $\tilde{\Delta} \times \tilde{\Delta}$, since the only case which had $(v_0 \dots v_s, v_s \dots v_r)^\xi \cap (v_0 \dots v_t, v_t \dots v_r)^\varphi \neq \emptyset$ also had $\varphi = \neg\xi$, that is to say in that case φ and ξ were opposing, rather than equal, parity symbols.

To prove the movement property for $\nabla(v_0 \dots v_r)$ start by considering $\nabla(v_0 \dots v_r)^\xi$ which is obtained by taking the union of the sets in equation (69) for $s = 1, 2, \dots, r$ and thus may be written

$$\nabla(v_0 \dots v_r)^\xi = \{(v_0 \dots \hat{v}_i \dots v_s, v_s \dots v_r) \mid 0 \leq i \leq s \leq r \text{ and } i \text{ is of parity } \xi\} \cup \{(v_0 \dots v_s, v_s \dots \hat{v}_i \dots v_r) \mid 0 \leq s \leq i \leq r \text{ and } i \text{ is of parity } \xi\} \quad (70)$$

what's more our analysis of $(v_0 \dots v_s, v_s \dots v_r)^\xi \cap (v_0 \dots v_t, v_t \dots v_r)^\varphi$ in the last paragraph demonstrates that:

$$\nabla(v_0 \dots v_r)^- \cap \nabla(v_0 \dots v_r)^+ = \{(v_0 \dots v_s, v_{s+1} \dots v_r) \mid s = 0, 1, \dots, r-1\} \quad (71)$$

On the other hand, consider a face $v_0 \dots \hat{v}_i \dots v_r$ of $v_0 \dots v_r$ and observe that we have

$$\nabla(v_0 \dots \hat{v}_i \dots v_r) = \{(v_0 \dots \hat{v}_i \dots v_s, v_s \dots v_r) \mid i < s \leq r\} \cup \{(v_0 \dots v_s, v_s \dots \hat{v}_i \dots v_r) \mid 0 \leq s < i\}$$

so it follows that $\nabla(v_0 \dots v_r)^\xi$, which is the union of these sets taken over those $i = 0, 1, \dots, r$ with parity ξ , is given by

$$\nabla(v_0 \dots v_r)^\xi = \{(v_0 \dots \hat{v}_i \dots v_s, v_s \dots v_r) \mid 0 \leq i < s \leq r \text{ and } i \text{ is of parity } \xi\} \cup \{(v_0 \dots v_s, v_s \dots \hat{v}_i \dots v_r) \mid 0 \leq s < i \leq r \text{ and } i \text{ is of parity } \xi\} \quad (72)$$

which differs from the expression in (70) only to the extent that it insists on strict inequalities between i and s . Notice that each element of $\nabla(v_0 \dots v_r)^\xi$ has a vertex which is held in common by both of its ordinates, it follows that no element of the form $(v_0 \dots v_s, v_{s+1} \dots v_r)$ can be in

there so we may infer from equations (70), (71) and (72) that we can express $\nabla(v_0 \dots v_r)^\xi$ as the disjoint union:

$$\nabla(v_0 \dots v_r)^\xi = \nabla(v_0 \dots v_r^\xi) \sqcup \{(v_0 \dots v_s, v_{s+1} \dots v_r) \mid s = 0, 1, \dots, r-1\}$$

Finally, from this latter expression and equation (71) it follows immediately that

$$\nabla(v_0 \dots v_r^-) = \nabla(v_0 \dots v_r)^\mp \quad \text{and} \quad \nabla(v_0 \dots v_r^+) = \nabla(v_0 \dots v_r)^\pm$$

and it is trivially the case that any subset $\nabla(v_0 \dots v_r)$ moves $\nabla(v_0 \dots v_r)^\mp$ to $\nabla(v_0 \dots v_r)^\pm$. $\square$

Observation 252. It is clear that for any $n \in \mathbb{N}$ the parity complex morphism of the previous lemma restricts to a morphism $\nabla_n: \tilde{\Delta}[n] \longrightarrow \tilde{\Delta}[n] \times \tilde{\Delta}[n]$ in Parity. Furthermore, it is only slightly less routine to demonstrate that these form a natural family, in the sense that for each simplicial operator $\alpha: [m] \longrightarrow [n]$ we have a commuting square

$$\begin{array}{ccc} \tilde{\Delta}[m] & \xrightarrow{\nabla_m} & \tilde{\Delta}[m] \times \tilde{\Delta}[m] \\ \tilde{\Delta}(\alpha) \downarrow & & \downarrow \tilde{\Delta}(\alpha) \times \tilde{\Delta}(\alpha) \\ \tilde{\Delta}[n] & \xrightarrow{\nabla_n} & \tilde{\Delta}[n] \times \tilde{\Delta}[n] \end{array}$$

in Parity. To demonstrate this we pick an element $\underline{v} = v_0 \dots v_r$ of $\tilde{\Delta}[m]$ and observe that $\tilde{\Delta}(\alpha)(\underline{v}) = \emptyset$ iff there is some i such that $\alpha(v_i) = \alpha(v_{i+1})$ which in turn holds iff for each $s \in [r]$ either $\tilde{\Delta}(\alpha)(v_0 \dots v_s)$ or $\tilde{\Delta}(\alpha)(v_s \dots v_r)$ is the empty set. It follows then that one leg of this diagram carries $\underline{v}$ to the empty set if and only if the other one does. In the common non-empty case, it is now a routine matter to show that either way we traverse the diagram we get the set $\{(\alpha(v_0 \dots v_s), \alpha(v_s \dots v_r)) \mid s = 0, 1, \dots, r\}$.

Observation 253. The last two results provide us with the tools to construct a canonical cocone with vertex $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ under the diagram $D_{\Delta[n] \times \Delta[m]}: \mathbb{G}(\Delta[n] \times \Delta[m]) \longrightarrow \omega\text{-}\underline{\text{Cat}}$ which we used to construct $F_\omega(\Delta[n] \times \Delta[m]) \in \omega\text{-}\underline{\text{Cat}}$ in observation 244.

The component of this cocone at the r -simplex $(\alpha, \beta) \in \Delta[n] \times \Delta[m]$, qua object of $\mathbb{G}(\Delta[n] \times \Delta[m])$, is constructed by applying the functor $\mathcal{O}$ to the following composite of parity complex morphisms:

$$\tilde{\Delta}[r] \xrightarrow{\nabla_r} \tilde{\Delta}[r] \times \tilde{\Delta}[r] \xrightarrow{\tilde{\Delta}(\alpha) \times \tilde{\Delta}(\beta)} \tilde{\Delta}[m] \times \tilde{\Delta}[n]$$

Of course we need to check that these collectively satisfy the naturality condition for such a cocone. So suppose that the simplicial operator $\gamma: [s] \longrightarrow [r]$ represents an arrow in $\mathbb{G}(\Delta[n] \times \Delta[m])$ from the s -simplex (α', β') to the r -simplex (α, β) and observe that we have the commutative diagram

$$\begin{array}{ccccc} \tilde{\Delta}[s] & \xrightarrow{\nabla_s} & \tilde{\Delta}[s] \times \tilde{\Delta}[s] & & \\ \tilde{\Delta}(\gamma) \downarrow & & \downarrow \tilde{\Delta}(\gamma) \times \tilde{\Delta}(\gamma) & \searrow \tilde{\Delta}(\alpha') \times \tilde{\Delta}(\beta') & \\ & & & \tilde{\Delta}[n] \times \tilde{\Delta}[m] & \\ \tilde{\Delta}[r] & \xrightarrow{\nabla_r} & \tilde{\Delta}[r] \times \tilde{\Delta}[r] & \nearrow \tilde{\Delta}(\alpha) \times \tilde{\Delta}(\beta) & \end{array}$$

in Parity. Here the square on the left expresses the naturality property discussed in observation 252 and the triangle on the right commutes by the functoriality of $\tilde{\Delta}(-) \times \tilde{\Delta}(-)$ applied to the defining property of γ as an arrow of $\mathbb{G}(\Delta[n] \times \Delta[m])$, that being $(\alpha', \beta') = (\alpha, \beta) \cdot \gamma = (\alpha \circ \gamma, \beta \circ \gamma)$. So applying the functor $\mathcal{O}$ to this diagram we get the required co-cone naturality property and it follows that this information induces a comparison ω -functor $c^{n,m}: F_\omega(\Delta[n] \times \Delta[m]) \longrightarrow \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ which is uniquely determined by the property that the square

$$\begin{array}{ccc} \mathcal{O}(\tilde{\Delta}[r]) & \xrightarrow{\mathcal{O}(\nabla_r)} & \mathcal{O}(\tilde{\Delta}[r] \times \tilde{\Delta}[r]) \\ \downarrow i_{(\alpha, \beta)} & & \downarrow \mathcal{O}(\tilde{\Delta}(\alpha) \times \tilde{\Delta}(\beta)) \\ F_\omega(\Delta[n] \times \Delta[m]) & \xrightarrow{c^{n,m}} & \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m]) \end{array}$$

commutes for each simplex (α, β) in $\Delta[n] \times \Delta[m]$ (where $r = \dim(\alpha, \beta)$). Alternatively, using the notation of observation 245, we see that we may characterise $c^{n,m}$ as the unique ω -functor which has

$$c^{n,m}(\llbracket \alpha, \beta \rrbracket) = \mathcal{O}((\tilde{\Delta}(\alpha) \times \tilde{\Delta}(\beta)) \circ \nabla_r)(\langle 01\dots r \rangle) \quad (73)$$

for each simplex (α, β) in $\Delta[n] \times \Delta[m]$ (where $r = \dim(\alpha, \beta)$).

Finally it is easily seen that the family of ω -functors $c^{n,m}$ is natural, in the sense that the square

$$\begin{array}{ccc} F_\omega(\Delta[n] \times \Delta[m]) & \xrightarrow{c^{n,m}} & \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m]) \\ \downarrow F_\omega(\Delta(\phi) \times \Delta(\psi)) & & \downarrow \mathcal{O}(\tilde{\Delta}(\phi) \times \tilde{\Delta}(\psi)) \\ F_\omega(\Delta[r] \times \Delta[s]) & \xrightarrow{c^{r,s}} & \mathcal{O}(\tilde{\Delta}[r] \times \tilde{\Delta}[s]) \end{array} \quad (74)$$

commutes for each pair of simplicial operators $\phi: [n] \longrightarrow [r]$ and $\psi: [m] \longrightarrow [s]$. We leave it up to the reader to verify this claim, either by a simple diagram chase and an appeal to the uniqueness property of our defining colimit or by considering the action of these ω -functors on the atoms of $F_\omega(\Delta[n] \times \Delta[m])$ and using the explicit descriptions given in equation (73) and observations 243 and 245.

Observation 254. In section 7 we singled our two classes of simplices in $\Delta[n] \times \Delta[m]$, cylinders and mediators, which we found to be of great interest in our analysis of the stratified set $\Delta[n] \otimes \Delta[m]$. On order to extend our work in the last observation to the ω -category $F_\omega(\Delta[n] \otimes \Delta[m])$, we are lead to consider how the ω -functor $c^{n,m}: F_\omega(\Delta[n] \times \Delta[m]) \longrightarrow \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ acts on those atoms which correspond to simplices in these classes.

In the sequel we'll be interested in analysing cells $c^{n,m}(\llbracket \alpha, \beta \rrbracket)$ where (α, β) is an r -simplex of $\Delta[n] \times \Delta[m]$ and our primary tool in making such calculations will be the expression in equation (73). It is therefore worth observing that we may combine the definitions in observations 235, 236 and 251 to show that we have:

$$((\tilde{\Delta}(\alpha) \times \tilde{\Delta}(\beta)) \circ \nabla_r)(01\dots r) = \{(\alpha(01\dots s), \beta(s(s+1)\dots r)) \mid s = 0, 1, \dots, r\} \cap (\tilde{\Delta}[n] \times \tilde{\Delta}[m])_r \quad (75)$$

In other words, we form $\nabla_r(01\dots r)$ then apply α and β to respective components of successive pairs in this set and finally we intersect the result with $(\tilde{\Delta}[n] \times \tilde{\Delta}[m])_r$ to exclude any elements which ended up having dimension less than r .

Cylinders: Firstly let $r = n + m$ and consider the unique non-degenerate r -dimensional cylinder $(\pi_1^{n,m}, \pi_2^{n,m})$ in $\Delta[n] \times \Delta[m]$. Consulting the definition of these partition operators it is easy to see that

$$\pi_1^{n,m}(01\dots n) = 01\dots n \quad \text{and} \quad \pi_2^{n,m}(n(n+1)\dots r) = 01\dots m$$

so we see, by equation (75), that $(01\dots n, 01\dots m) \in ((\tilde{\Delta}(\pi_1^{n,m}) \times \tilde{\Delta}(\pi_2^{n,m})) \circ \nabla_r)(01\dots r)$. It follows therefore that $c^{n,m}(\llbracket \pi_1^{n,m}, \pi_2^{n,m} \rrbracket) = \mathcal{O}((\tilde{\Delta}(\pi_1^{n,m}) \times \tilde{\Delta}(\pi_2^{n,m})) \circ \nabla_r)(\langle 01\dots r \rangle)$ has non-empty sets of r -dimensional elements and is thus a non-trivial r -cell in $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$. However we know, by the comment at the end of observation 228, that the atom $\langle 01\dots n, 01\dots m \rangle$ is the unique non-trivial r -cell of $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ and we must have $c^{n,m}(\llbracket \pi_1^{n,m}, \pi_2^{n,m} \rrbracket) = \langle 01\dots n, 01\dots m \rangle$.

Now, referring back to observation 134, by definition we know that the partition (p, q) of r witnesses the r -simplex $(\alpha, \beta) \in \Delta[n] \times \Delta[m]$ to be a cylinder iff there are (necessarily unique) simplicial operators $\alpha': [p] \longrightarrow [n]$ and $\beta': [q] \longrightarrow [m]$ with $\alpha = \alpha' \circ \pi_1^{p,q}$ and $\beta = \beta' \circ \pi_2^{p,q}$. Furthermore, consulting the definition our partition operators $\pi_i^{p,q}$, it is clear that (α, β) is non-degenerate iff α' and β' are both face operators. In that case, we may define $\underline{v} = \alpha(01\dots r)$ and $\underline{w} = \beta(01\dots r)$, observe that these are elements of $\tilde{\Delta}[n]_p$ and $\tilde{\Delta}[m]_q$ respectively and easily show that $\alpha' = \delta_{\underline{v}}^n$ and $\beta' = \delta_{\underline{w}}^m$. Finally, the naturality property depicted in display (74), when applied to the operators $\delta_{\underline{v}}^n$ and $\delta_{\underline{w}}^m$ and evaluated at the atom $\llbracket \pi_1^{p,q}, \pi_2^{p,q} \rrbracket$, demonstrates that

$$c^{n,m}(\llbracket \alpha, \beta \rrbracket) = c^{n,m}(\llbracket \delta_{\underline{v}}^n \circ \pi_1^{p,q}, \delta_{\underline{w}}^m \circ \pi_2^{p,q} \rrbracket) = \mathcal{O}(\tilde{\Delta}(\delta_{\underline{v}}^n) \times \tilde{\Delta}(\delta_{\underline{w}}^m))(c^{p,q}(\llbracket \pi_1^{p,q}, \pi_2^{p,q} \rrbracket))$$

but we've just shown that

$$c^{p,q}(\llbracket \pi_1^{p,q}, \pi_2^{p,q} \rrbracket) = \langle 01\dots p, 01\dots q \rangle$$

so it follows that

$$\begin{aligned} c^{n,m}(\llbracket \alpha, \beta \rrbracket) &= \mathcal{O}(\tilde{\Delta}(\delta_{\underline{v}}^n) \times \tilde{\Delta}(\delta_{\underline{w}}^m))(\langle 01\dots p, 01\dots q \rangle) \\ &= \langle \delta_{\underline{v}}^n(01\dots p), \delta_{\underline{w}}^m(01\dots q) \rangle \\ &= \langle \underline{v}, \underline{w} \rangle \end{aligned}$$

where the second equality follows from observation 243 and the last one follows directly from the relationship between $\underline{v}$, $\underline{w}$ and the corresponding face operators $\delta_{\underline{v}}^n$, $\delta_{\underline{w}}^m$.

In summary, we have demonstrated that $c^{n,m}$ restricts to a bijection between the set of r -dimensional atoms in $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ and the set of atoms $\llbracket \alpha, \beta \rrbracket$ in $F_\omega(\Delta[n] \times \Delta[m])$ for which (α, β) is a non-degenerate cylinder in $\Delta[n] \times \Delta[m]$.

Mediators: Consider an arbitrary r -simplex $(\alpha, \beta) \in \Delta[n] \times \Delta[m]$ and observe that equations (73) and (75) imply that $c^{n,m}(\llbracket \alpha, \beta \rrbracket)$ is an $(r-1)$ -cell iff $(\alpha(01\dots s), \beta(s(s+1)\dots r)) \notin (\tilde{\Delta}[n] \times \tilde{\Delta}[m])_r$ (for all $s = 0, 1, \dots, r$). Of course, this condition holds if and only if either $\alpha(01\dots s)$ has dimension $< s$ (that is $\exists 0 \leq i < s$ s.t. $\alpha(i) = \alpha(i+1)$) or $\beta(s(s+1)\dots r)$ has dimension $< (r-s)$ (in other words $\exists s \leq j < r$ s.t. $\beta(j) = \beta(j+1)$).

In fact, if we let $s \leq r$ be the largest integer such that $\alpha(0) < \alpha(1) < \dots < \alpha(s)$ then this latter condition implies that there is some $s \leq j < r$ such that $\beta(j) = \beta(j+1)$ and then, by the maximality of s , we also have $\alpha(s) = \alpha(s+1)$. Conversely, if we had some $i \leq j$ such that $\alpha(i) = \alpha(i+1)$ and $\beta(j) = \beta(j+1)$ then it is clear that the triviality condition of the last paragraph immediately holds.

Notice that we've simply demonstrated that $c^{n,m}(\llbracket \alpha, \beta \rrbracket)$ is an $(r-1)$ -cell if and only if the condition of lemma 129(c) holds and we know, by corollary 130, that this characterises the thin simplices in the tensor product $\Delta[n] \otimes \Delta[m]$.

Of course, given that this result builds a strong and compelling link to our work on tensor products of stratified sets in sections 7 and 8, it would be tempting to claim that it was this observation which motivated our original definition of that tensor product. However this is by no means the case, since we arrived at our original definition of $\Delta[n] \otimes \Delta[m]$ through purely simplicial means and only later derived its close relationship to the free ω -category $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$.

These observations lead directly to, and are summarised by, the primary result of this subsection:

Theorem 255. *The ω -functor $c^{n,m}: F_\omega(\Delta[n] \times \Delta[m]) \longrightarrow \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ factors through the canonical collapse $q_t: F_\omega(\Delta[n] \times \Delta[m]) \longrightarrow F_\omega(\Delta[n] \otimes \Delta[m])$ and provides a comparison ω -functor*

$$F_\omega(\Delta[n] \otimes \Delta[m]) \xrightarrow{\bar{c}^{n,m}} \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$$

which inherits the naturality properties of $c^{n,m}$. If (p, q) is the partition which witnesses the non-degenerate r -simplex (α, β) as a cylinder then we have elements $\underline{v} = \alpha(01\dots r)$ in $\tilde{\Delta}[n]_p$ and $\underline{w} = \beta(01\dots r)$ in $\tilde{\Delta}[m]_q$ (for which $\alpha = \delta_{\underline{v}}^n \circ \pi_1^{p,q}$ and $\beta = \delta_{\underline{w}}^m \circ \pi_2^{p,q}$) and $\bar{c}^{n,m}$ maps the cell $\llbracket \alpha, \beta \rrbracket$ in $F_\omega(\Delta[n] \otimes \Delta[m])$ to the atom $\langle \underline{w}, \underline{v} \rangle$ in $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$. It follows that $\bar{c}^{n,m}$ restricts to a bijection between each set

$$C_r^{n,m} \stackrel{\text{def}}{=} \{ \llbracket \alpha, \beta \rrbracket \mid (\alpha, \beta) \text{ is a non-degenerate } r\text{-cylinder in } \Delta[n] \otimes \Delta[m] \}$$

and the set of r -dimensional atoms of $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$. Finally, $F_\omega(\Delta[n] \otimes \Delta[m])$ is weakly generated by the cells in the set $C^{n,m}$ and consequently we may apply lemma 223 to show that $\bar{c}^{n,m}$ is an **isomorphism** of ω -categories.

Proof. The latter part of the last observation demonstrated that the ω -functor $c^{n,m}$ maps atoms $\llbracket \alpha, \beta \rrbracket$ associated with those r -simplices $(\alpha, \beta) \in \Delta[n] \times \Delta[m]$ which are thin in $\Delta[n] \otimes \Delta[m]$ to $(r-1)$ -cells in $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$. In other words, $c^{n,m}$ collapses the cells in the graded set $\llbracket t(\Delta[n] \otimes \Delta[m]) \rrbracket \subseteq F_\omega(\Delta[n] \times \Delta[m])$ and it therefore factors through the quotient $F_\omega(\Delta[n] \otimes \Delta[m])$ as postulated. Checking the naturality properties of the resulting ω -functors $\bar{c}^{n,m}$ is a routine matter which we leave up to the reader. Notice also that the first half of the last observation demonstrated that $c^{n,m}$, and thus $\bar{c}^{n,m}$, acts on cells $\llbracket \alpha, \beta \rrbracket$ associated with non-degenerate cylinders as stated and thus that it restricts to a bijection between each $C_r^{n,m}$ and the set of r -dimensional atoms of $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$.

So all that remains is to apply induction on $(n+m)$ to prove that $C^{n,m}$ weakly generates the ω -category $F_\omega(\Delta[n] \otimes \Delta[m])$. The base case is trivial, so we adopt the inductive hypothesis, that $C^{p,q}$ weakly generates $F_\omega(\Delta[p] \otimes \Delta[q])$ whenever $p+q < n+m$, and consider family of stratified maps

$$\begin{aligned} \Delta(\delta_i^n) \otimes \Delta[m]: \Delta[n-1] \otimes \Delta[m] &\longrightarrow \Delta[n] \otimes \Delta[m] \quad \text{for } i = 0, 1, \dots, n \text{ and} \\ \Delta[n] \otimes \Delta(\delta_j^m): \Delta[n] \otimes \Delta[m-1] &\longrightarrow \Delta[n] \otimes \Delta[m] \quad \text{for } j = 0, 1, \dots, m. \end{aligned} \tag{76}$$

each of which factors through the regular boundary subset $\partial(\Delta[n] \otimes \Delta[m]) \subseteq_r \Delta[n] \otimes \Delta[m]$ of notation 166(ii) to provide a jointly epimorphic family of stratified maps with codomain $\partial(\Delta[n] \otimes \Delta[m])$. The functor F_ω preserves coproducts and epimorphisms, since it is left adjoint,

so applying it to this family we obtain a jointly epimorphic family of ω -functors with codomain $F_\omega(\partial(\Delta[n] \otimes \Delta[m]))$.

Clearly $\partial(\Delta[n] \otimes \Delta[m])$ is actually a regular subset of the $(n + m - 1)$ -superstructure $|\Delta[n] \otimes \Delta[m]|_{n+m-1}$ and indeed it is easily seen that if $H(\Delta[n], \Delta[m]) \subseteq_r \Delta[n] \otimes \Delta[m]$ is the regular subset defined in notation 166(iii) then $|\Delta[n] \otimes \Delta[m]|_{n+m-1} = \partial(\Delta[n] \otimes \Delta[m]) \cup H(\Delta[n], \Delta[m])$ so we have a pasting square:

$$\begin{array}{ccc} \partial H(\Delta[n], \Delta[m]) & \xrightarrow{\subseteq_r} & H(\Delta[n], \Delta[m]) \\ \downarrow \subseteq_r & & \downarrow \subseteq_r \\ \partial(\Delta[n] \otimes \Delta[m]) & \xrightarrow[\subseteq_r]{} & |\Delta[n] \otimes \Delta[m]|_{n+m-1} \end{array}$$

By lemma 168, we know that the upper horizontal in this square is an f-extension and so the lower horizontal inclusion is also an f-extension (by observation 158(b)). It follows, from theorem 249, that this inclusion is perpendicular to the nerve $N_\omega(\mathbb{C})$ of each ω -category or equivalently that the ω -functor $F_\omega(\subseteq_r): F_\omega(\partial(\Delta[n] \otimes \Delta[m])) \longrightarrow F_\omega(|\Delta[n] \otimes \Delta[m]|_{n+m-1})$ is an isomorphism.

We also know, from observation 248, that the regular inclusion $|\Delta[n] \otimes \Delta[m]|_{n+m-1} \subseteq_r \Delta[n] \otimes \Delta[m]$ provides an ω -functor $F_\omega(\subseteq_r): F_\omega(|\Delta[n] \otimes \Delta[m]|_{n+m-1}) \longrightarrow F_\omega(\Delta[n] \otimes \Delta[m])$ which restricts to an epimorphism with codomain $|F_\omega(\Delta[n] \otimes \Delta[m])|_{n+m-1}$. Observe now that the functoriality of F_ω implies that the family obtained by composing this latter epimorphic ω -functor with the composite of the isomorphism of the previous paragraph and the jointly epimorphic family of the paragraph before that gives a family of ω -functors which could otherwise be constructed by applying F_ω to the stratified maps in display 76 and factoring each one through $|F_\omega(\Delta[n] \otimes \Delta[m])|_{n+m-1} \subseteq F_\omega(\Delta[n] \otimes \Delta[m])$. As the composite of an epimorphism, an isomorphism and a jointly epimorphic family this too is a jointly epimorphic family.

If $(\alpha, \beta) \in \Delta[n-1] \otimes \Delta[m]$ is a non-degenerate cylinder then $F_\omega(\Delta(\delta_i^n) \otimes \Delta[m])(\llbracket \alpha, \beta \rrbracket) = \llbracket \delta_i^n \circ \alpha, \beta \rrbracket$ and $(\delta_i^n \circ \alpha, \beta)$ is a non-degenerate cylinder in $\Delta[n] \otimes \Delta[m]$, in other words the ω -functor $F_\omega(\Delta(\delta_i^n) \otimes \Delta[m])$ maps cells in $C^{n-1, m}$ into $|C^{n, m}|_{n+m-1}$. Similarly, each ω -functor $F_\omega(\Delta[n] \otimes \Delta(\delta_j^m))$ maps cells in $C^{m, n-1}$ into $|C^{n, m}|_{n+m-1}$. Consequently, we may apply lemma 221, and the associated observation 222, to the jointly epimorphic family of the last paragraph and the weakly generating sets $C^{n-1, m} \subseteq F_\omega(\Delta[n-1] \otimes \Delta[m])$ and $C^{m, n-1} \subseteq F_\omega(\Delta[n] \otimes \Delta[m-1])$ of the inductive hypothesis to show that $|C^{n, m}|_{n+m-1}$ weakly generates $|F_\omega(\Delta[n] \otimes \Delta[m])|_{n+m-1}$.

Finally, the only cell in the standard weak generator $\llbracket \Delta[n] \otimes \Delta[m] \rrbracket$ of $F_\omega(\Delta[n] \otimes \Delta[m])$ (cf. observation 246) which is not in $|F_\omega(\Delta[n] \otimes \Delta[m])|_{n+m-1}$ is the atom $\llbracket \pi_1^{n, m}, \pi_2^{n, m} \rrbracket$ because all simplices of $\Delta[n] \otimes \Delta[m]$ above dimension $(n + m)$ are degenerate and its only non-thin $(n + m)$ -simplex is the minimal shuffle $(\pi_1^{n, m}, \pi_2^{n, m})$. Therefore since $|C^{n, m}|_{n+m-1}$ weakly generates $|F_\omega(\Delta[n] \otimes \Delta[m])|_{n+m-1}$ we may infer that $C^{n, m} = |C^{n, m}|_{n+m-1} \cup \{\llbracket \pi_1^{n, m}, \pi_2^{n, m} \rrbracket\}$ weakly generates $F_\omega(\Delta[n] \otimes \Delta[m])$ as required. $\square$

Corollary 256. *For each pair of stratified sets N and M in Simplex the isomorphism of theorem 255 induces an isomorphism $\bar{c}^{N, M}: F_\omega(N \otimes M) \longrightarrow \mathcal{O}(\bar{N} \times \bar{M})$ which is uniquely*

determined by the fact that it makes the square

$$\begin{array}{ccc}
 F_\omega(\Delta[n] \otimes \Delta[m]) & \xrightarrow[\cong]{\bar{c}^{n,m}} & \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m]) \\
 F_\omega(\subseteq_e) \downarrow & & \downarrow \mathcal{O}(\subseteq_e) \\
 F_\omega(N \otimes M) & \xrightarrow[\exists! \bar{c}^{N,M}]{\cong} & \mathcal{O}(\tilde{N} \times \tilde{M})
 \end{array} \tag{77}$$

commute. In other words, $\bar{c}^{N,M}$ is characterised by the fact that it maps the cell $[\alpha, \beta] \in F_\omega(N \otimes M)$ associated with a non-thin r -cylinder $(\alpha, \beta) \in N \otimes M$ to the cell $\langle \underline{v}, \underline{w} \rangle \in \mathcal{O}(\tilde{N} \times \tilde{M})$ with $\underline{v} = \alpha(01\dots r)$ and $\underline{w} = \beta(01\dots r)$. It follows, therefore, that this family of isomorphisms is natural in $N, M \in \underline{\text{Simplex}}$.

Proof. First observe that both of the vertical ω -functors in display (77) are collapsers. The right hand one is simply the collapser we used to construct $\mathcal{O}(\tilde{N} \times \tilde{M})$ from the free ω -category $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ in observation 239, in other words this is the collapser of the graded subset of this latter ω -category of those atoms associated with elements which are thin in $\tilde{N} \times \tilde{M}$. The left hand one features in the commutative square

$$\begin{array}{ccc}
 F_\omega(\Delta[n] \boxtimes \Delta[m]) & \xrightarrow[\cong]{F_\omega(\subseteq_e)} & F_\omega(\Delta[n] \otimes \Delta[m]) \\
 F_\omega(\subseteq_e) \downarrow & & \downarrow F_\omega(\subseteq_e) \\
 F_\omega(N \boxtimes M) & \xrightarrow[\cong]{F_\omega(\subseteq_e)} & F_\omega(N \otimes M)
 \end{array}$$

where the horizontal ω -functors are obtained by applying F_ω to the t-extensions of lemma 139, and are therefore isomorphisms by Street's theorem 249. As discussed in observation 246 the left hand vertical in this square, and thus its isomorphic right hand vertical, is a collapser of the set of cells associated with those simplices which are thin in $N \boxtimes M$ but not thin in $\Delta[n] \boxtimes \Delta[m]$. Returning to definition 135, we see that the simplices which satisfy this condition are precisely the non-degenerate cylinders that are thin in $N \otimes M$.

Now we know, from theorem 255, that the isomorphism $\bar{c}^{n,m}$ at the top of display (77) maps the cell $[\alpha, \beta] \in F_\omega(\Delta[n] \otimes \Delta[m])$ associated with a non-degenerate cylinder (α, β) to a corresponding atom $\langle \underline{v}, \underline{w} \rangle \in \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[m])$ and that these are related by $\alpha = \delta_{\underline{w}}^n \circ \pi_1^{p,q}$ and $\beta = \delta_{\underline{v}}^m \circ \pi_2^{p,q}$. Consulting observation 134, we see that our cylinder is thin in $N \otimes M$ if either $\delta_{\underline{v}}^n$ is thin in N or $\delta_{\underline{w}}^m$ is thin in M and consequently we may infer, from the definitions of the corresponding stratified parity complexes $\tilde{N}$, $\tilde{M}$ and $\tilde{N} \times \tilde{M}$, that our non-degenerate cylinder (α, β) is thin in $N \otimes M$ iff the corresponding element $(\underline{v}, \underline{w})$ is thin in $\tilde{N} \times \tilde{M}$. It follows that $\bar{c}^{n,m}$ sets up a graded bijection between the sets of cells with respect to which the vertical ω -functors of display (77) are collapsers and thus that it induces the isomorphism at the bottom of that square as postulated. $\square$

11.5. An Inductive Proof of the Street-Roberts Conjecture.

Observation 257 (an outline of our proof of the Street-Roberts conjecture). We now have almost all of the technical tools required to complete our proof of the Street-Roberts conjecture, which is that the nerve functor $N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}$ is an equivalence of categories. Our approach will be to inductively prove that it restricts to an equivalence $N_n: n\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}_n$

for each $n \in \mathbb{N}$ and then use a simple colimiting argument to extend those equivalences to $N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}$

To complete this proof the only real work left is to show that we may construct an essentially commutative square

$$\begin{array}{ccc}
 \omega\text{-}\underline{\text{Cat}} & \xrightarrow{P} & (\omega\text{-}\underline{\text{Cat}})\text{-}\underline{\text{Cat}} \\
 N_\omega \downarrow & \cong & \downarrow N_{\omega\text{-}\underline{\text{Cat}}} \\
 \underline{\text{Cs}} & \xrightarrow{P^r} & \underline{\text{Cs}}\text{-}\underline{\text{Cat}}
 \end{array} \tag{78}$$

in which the horizontal functor along the top is the equivalence discussed in observation 85 and the functor along the bottom is the right handed version of the path category construction discussed in sections 9 and 10. Furthermore, the right vertical here is the functor obtained by applying the (product preserving) nerve functor N_ω point-wise to the hom- ω -categories of each $\omega\text{-}\underline{\text{Cat}}$ -enriched category.

Once we have constructed this square, the remainder of the proof is straightforward. First we combine lemma 247, lemma 184 and the discussion in observation 85 to show that for each $n \in \mathbb{N}$ the square in display (78) restricts to:

$$\begin{array}{ccc}
 (n+1)\text{-}\underline{\text{Cat}} & \xrightarrow{P} & (n\text{-}\underline{\text{Cat}})\text{-}\underline{\text{Cat}} \\
 N_{n+1} \downarrow & \cong & \downarrow N_{n\text{-}\underline{\text{Cat}}} \\
 \underline{\text{Cs}}_{n+1} & \xrightarrow{P^r} & \underline{\text{Cs}}_n\text{-}\underline{\text{Cat}}
 \end{array} \tag{79}$$

Applying induction we use this to show that N_ω restricts to an equivalence $N_n: n\text{-}\underline{\text{Cat}} \xrightarrow{\sim} \underline{\text{Cs}}_n$ for each $n \in \mathbb{N}$. The base case $n = 0$ follows immediately from lemma 173. For the inductive step we adopt the hypothesis that $N_n: n\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}_n$ is an equivalence, from which it is immediate that right hand vertical functor $N_{n\text{-}\underline{\text{Cat}}}$ of display (79) is an equivalence. Furthermore, as remarked in observation 85, the upper horizontal arrow P in this square is also an equivalence, so it follows that its upper right composite, and thus its isomorphic lower left composite, is an equivalence. However we know, by theorem 213, that the lower horizontal functor P^r is fully faithful, which allows us to apply lemma 212(ii) and demonstrate that the left hand vertical N_{n+1} is an equivalence as required.

Of course, this result immediately implies that the adjunction $F_n \dashv N_n: n\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}_n$ is an adjoint equivalence (cf. [22]) for each $n \in \mathbb{N}$. The unit and counit of each such adjunction is obtained by restricting the unit $\eta: \text{id}_{\underline{\text{Cs}}} \longrightarrow N_\omega \circ F_\omega$ and counit $\epsilon: F_\omega \circ N_\omega \longrightarrow \text{id}_{\omega\text{-}\underline{\text{Cat}}}$ of the adjunction $F_\omega \dashv N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}$. So it follows that each $F_n \dashv N_n$ is an adjoint equivalence iff the unit component η_A is an isomorphism whenever A is n -complicial and the counit component ϵ_C is an isomorphism whenever C is an n -category.

To extend this result to all components of η and ϵ , we exploit the fact that each ω -category or complicial set is equal to the union of its ascending sequence of superstructures. To elaborate, for each ω -category $\mathbb{C}$ the cocone of inclusions $i_n: |\mathbb{C}|_n \hookrightarrow \mathbb{C}$ displays $\mathbb{C}$ as the colimit $\text{colim}_{n \in \mathbb{N}} |\mathbb{C}|_n$ of the ascending chain $|\mathbb{C}|_0 \hookrightarrow |\mathbb{C}|_1 \hookrightarrow \dots \hookrightarrow |\mathbb{C}|_n \hookrightarrow \dots$ of superstructures in $\omega\text{-}\underline{\text{Cat}}$. Of course all colimits are preserved by the left adjoint F_ω , furthermore ascending chains are filtered diagrams and thus, by observation 250, the nerve functor N_ω also preserves the colimits of such chains. It follows therefore that the cocone

$F_\omega(N_\omega(i_n)): F_\omega(N_\omega(|\mathbb{C}|_n)) \longrightarrow F_\omega(N_\omega(\mathbb{C}))$ displays $F_\omega(N_\omega(\mathbb{C}))$ as the colimit of the chain: $F_\omega(N_\omega(|\mathbb{C}|_0)) \longrightarrow F_\omega(N_\omega(|\mathbb{C}|_1)) \longrightarrow \dots \longrightarrow F_\omega(N_\omega(|\mathbb{C}|_n)) \longrightarrow \dots$. Applying the naturality of ϵ to the inclusions $|\mathbb{C}|_n \hookrightarrow \mathbb{C}$ we get a commutative square

$$\begin{array}{ccc} F_\omega(N_\omega(|\mathbb{C}|_n)) & \xrightarrow{\epsilon_{|\mathbb{C}|_n}} & |\mathbb{C}|_n \\ F_\omega(N_\omega(i_n)) \downarrow & & \downarrow i_n \\ F_\omega(N_\omega(\mathbb{C})) & \xrightarrow{\epsilon_{\mathbb{C}}} & \mathbb{C} \end{array}$$

for each $n \in \mathbb{N}$, and this family displays $\epsilon_{\mathbb{C}}$ as the ω -functor induced between these colimits by the family of ω -functors $\{\epsilon_{|\mathbb{C}|_n}\}_{n \in \mathbb{N}}$. However, we've already shown that the component of ϵ on each n -category $|\mathbb{C}|_n$ is an isomorphism and so the induced ω -functor $\epsilon_{\mathbb{C}}$ is also an isomorphism. Applying a dual argument to the ascending chain of superstructures of a complicial set A we see that each η_A is also an isomorphism. It follows that $F_\omega \dashv N_\omega$ is an adjoint equivalence as required.

We now turn to completing our proof by establishing the isomorphism depicted in display (78), the construction of which relies primarily on the parity complex calculations laid out in the next few observations.

Notation 258 (intervals). In the sequel if $s \leq t$ are integers then we will use the following traditional interval notation for sequences of integers:

$$\begin{aligned} [s, t] &\stackrel{\text{def}}{=} \{i \in \mathbb{N} \mid s \leq i \leq t\} & (s, t) &\stackrel{\text{def}}{=} \{i \in \mathbb{N} \mid s < i < t\} \\ [s, t) &\stackrel{\text{def}}{=} \{i \in \mathbb{N} \mid s \leq i < t\} & (s, t] &\stackrel{\text{def}}{=} \{i \in \mathbb{N} \mid s < i \leq t\} \end{aligned}$$

Observation 259 (two point suspensions of parity complexes). If C is a parity complex and $n \in \mathbb{N}$ then we may define a parity complex $\Sigma_n C$ with underlying graded set

$$(\Sigma_n C)_0 = \{0, 1, \dots, n\} \quad (\Sigma_n C)_{r+1} = C_r \times \{1, \dots, n\}$$

and face operations:

$$\begin{aligned} (x, i)^- &= \{i - 1\} & \text{and} \\ (x, i)^+ &= \{i\} & \text{when } (x, i) \in (\Sigma_n C)_1, \\ (x, i)^\varphi &= x^\varphi \times \{i\} & \text{when } (x, i) \in (\Sigma_n C)_{r+1} \text{ with } r > 0. \end{aligned}$$

It is a routine matter to check that this is indeed a parity complex and that it satisfies all of the extra technical conditions discussed in definition 216. In outline, all of the basic parity complex axioms and the globularity condition involve relationships between 2 or 3 adjacent dimensions so they clearly hold for $\Sigma_n C$ above dimension 0, where it is no more than a disjoint union of copies of C which have all been shifted up a dimension. Instances of these conditions which involve faces at dimension 0 all admit simple direct verifications. Finally the remaining technical condition, which involves the dimension traversing ordering $\blacktriangleleft$ on $\Sigma_n C$ and its odd dual, follows directly from the simple observation that if $(c, i) \blacktriangleleft (x', i')$ in $\Sigma_n C$ then either $i < i'$ or we have $i = i'$ and $x \blacktriangleleft x'$ in C .

We can extend this construction to a functor $\Sigma_* C: \Delta \longrightarrow \underline{\text{Parity}}$ which maps each ordinal $[n] \in \Delta$ to $\Sigma_n C$ and each simplicial operator $\alpha: [n] \longrightarrow [m]$ to a parity complex morphism

$\Sigma_\alpha C: \Sigma_n C \longrightarrow \Sigma_m C$ given by:


$$\begin{aligned} (\Sigma_\alpha C)(i) &= \{\alpha(i)\} && \text{for } i \in (\Sigma_n C)_0, \\ (\Sigma_\alpha C)(x, i) &= \{x\} \times (\alpha(i-1), \alpha(i)] && \text{for } (x, i) \in (\Sigma_n C)_{r+1} \text{ with } r \in \mathbb{N}. \end{aligned}$$

Showing that this morphism of graded sets does satisfy the conditions given in lemma 231 and indeed that this construction is functorial in $\alpha \in \Delta$ are matters of routine verification which we leave to the reader. Notice also that if $f: C \longrightarrow D$ is a parity complex morphism then we may define a morphism of graded sets $\Sigma_n f: \Sigma_n C \longrightarrow \Sigma_n D$ by

$$\begin{aligned} (\Sigma_n f)(i) &= \{i\} && \text{for } i \in (\Sigma_n C)_0, \\ (\Sigma_n f)(x, i) &= f(x) \times \{i\} && \text{for } (x, i) \in (\Sigma_n C)_{r+1} \text{ with } r \in \mathbb{N}. \end{aligned}$$

which is again easily shown to be a parity complex morphism and functorial in $f \in \underline{\text{Parity}}$. Combining these observations it is clear that we've succeeded in naturally extending our construction to a functor $\Sigma: \Delta \times \underline{\text{Parity}} \longrightarrow \underline{\text{Parity}}$.

To aid the intuition it is appropriate to think of $\Sigma_1 C$ as being the two point suspension of C and to view $\Sigma_n C$ as an oriented path of n such suspensions abutting at dimension 0, for instance we might picture $\Sigma_3 \tilde{\Delta}[1]$ as:



Indeed, in line with this intuition it is possible to show that the functor $\mathcal{O}(\Sigma_* C): \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$ is actually a cocategory ($\mathbb{T}_{\underline{\text{Cat}}}\text{-coalgebra}$). To do so we apply the alternate, finite sketch based, presentation of $\mathbb{T}_{\underline{\text{Cat}}}\text{-(co)algebras}$ given in observation 70 and consider the commutative square

$$\begin{array}{ccc} \mathcal{O}(\Sigma_0 C) & \xrightarrow{\mathcal{O}(\Sigma_{\varepsilon_p^p C})} & \mathcal{O}(\Sigma_p C) \\ \mathcal{O}(\Sigma_{\varepsilon_0^q C}) \downarrow & \searrow & \downarrow \mathcal{O}(\Sigma_{\varepsilon_1^{p,q} C}) \\ \mathcal{O}(\Sigma_q C) & \xrightarrow[\mathcal{O}(\Sigma_{\varepsilon_2^{p,q} C})]{} & \mathcal{O}(\Sigma_{p+q} C) \end{array} \quad (80)$$

obtained by applying the functor $\mathcal{O}(\Sigma_* C)$ to the square in display (9). It is clear, from the definition of $\Sigma_\alpha C$ for a simplicial operator α , that the lower horizontal in this square restricts to an isomorphism between $\mathcal{O}(\Sigma_q C)$ and the sub- ω -category of $\mathcal{O}(\Sigma_{p+q} C)$ freely generated by the sub-parity complex consisting of those 0-elements i with $i \leq p$ and $(r+1)$ -elements (x, i) with $i \leq p$. Dually, its right hand vertical restricts to an isomorphism between $\mathcal{O}(\Sigma_p C)$ and the sub- ω -category of $\mathcal{O}(\Sigma_{p+q} C)$ freely generated by the sub-parity complex of 0-elements i with $i \geq p$ and $(r+1)$ -elements (x, i) with $i > p$. Furthermore, the union of these sub-parity complexes is $\Sigma_{p+q} C$ itself and the diagonal map in our square restricts to an isomorphism with the sub- ω -category freely generated by the intersection of these sub-parity complexes, which consists of the solitary 0-cell $\langle p \rangle$. Finally, it follows that we may apply lemma 220 to show that the square in display (80) is a pushout in $\omega\text{-}\underline{\text{Cat}}$ as required.

To identify the functor $P_C: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cat}}$ obtained by applying Kan's construction (observation 63) to this cocategory we start by studying $\mathcal{O}(\Sigma_1 C)$. It is immediate, from the definition of $\Sigma_1 C$, that each cell (N, P) in the freely generated ω -category $\mathcal{O}(C)$ gives rise

to a corresponding cell $(N, P)^* \stackrel{\text{def}}{=} (\{0\} \cup (N \times \{1\}), \{1\} \cup (P \times \{1\}))$ in $\mathcal{O}(\Sigma_1 C)$ and, indeed, that the only cells of $\mathcal{O}(\Sigma_1 C)$ that are not of this form are the 0-cells $\langle 0 \rangle$ and $\langle 1 \rangle$. Furthermore it is also clear that cells (N, P) and (M, Q) are r -composable in $\mathcal{O}(C)$ iff $(N, P)^*$ and $(M, Q)^*$ are $(r+1)$ -composable in $\mathcal{O}(\Sigma_1 C)$ and that in this case we have $(M, Q)^* *_{r+1} (N, P)^* = ((M, Q) *_{r+1} (N, P))^*$. However, if $\mathbb{C}$ is any ω -category then we may combine this analysis with the definition of the enriched category $\mathbf{P}(\mathbb{C}) \in (\omega\text{-}\underline{\mathbf{Cat}})\text{-}\underline{\mathbf{Cat}}$ given in observation 85 to show that ω -functors $f: \mathcal{O}(\Sigma_1 C) \longrightarrow \mathbb{C}$ with $u = f(\langle 0 \rangle)$ and $v = f(\langle 1 \rangle)$ correspond bijectively to ω -functors $\hat{f}: \mathcal{O}(C) \longrightarrow \mathbf{P}(\mathbb{C})(u, v)$ where these are related by the equality $\hat{f}(N, P) = f(N, P)^*$ for each cell $(N, P) \in \mathcal{O}(C)$.

In order to describe the composition of $\mathbf{P}_C(\mathbb{C})$ in these terms we need to study the ω -functor $w: \mathcal{O}(\Sigma_2 C) \longrightarrow \mathbb{C}$ that witnesses the composite of a pair of arrows $f, g: \mathcal{O}(\Sigma_1 C) \longrightarrow \mathbb{C}$ and which, by the definition given in observation 70, is the unique such ω -functor with $w \circ (\Sigma_{\delta_2^2} C) = f$, $w \circ (\Sigma_{\delta_0^2} C) = g$ and $w \circ (\Sigma_{\delta_1^2} C) = g * f$. Now, a simple calculation demonstrates that if (N, P) is a cell in $\mathcal{O}(C)$ then we have the equality

$$(\Sigma_{\delta_2^2} C)(N, P)^* *_0 (\Sigma_{\delta_2^2} C)(N, P)^* = (\Sigma_{\delta_1^2} C)(N, P)^*$$

in $\mathcal{O}(\Sigma_2 C)$, to which we may apply the ω -functor w , the witnessing equalities of the last sentence and the defining relationship of the last paragraph to show that:

$$\begin{aligned} \widehat{g * f}(N, P) &= (g * f)(N, P)^* \\ &= w((\Sigma_{\delta_1^2} C)(N, P)^*) = w((\Sigma_{\delta_2^2} C)(N, P)^* *_0 w((\Sigma_{\delta_2^2} C)(N, P)^*)) \\ &= g(N, P)^* *_0 f(N, P)^* = \hat{g}(N, P) *_0 \hat{f}(N, P) \end{aligned}$$

In summary, our analysis shows that the objects of $\mathbf{P}_C(\mathbb{C})$ correspond to 0-cells of $\mathbb{C}$, its homset $\mathbf{P}_C(\mathbb{C})(u, v)$ is isomorphic to the set $\omega\text{-}\underline{\mathbf{Cat}}(\mathcal{O}(C), \mathbf{P}(\mathbb{C})(u, v))$ and under these isomorphisms its composition corresponds to the point-wise composition of functors.

In other words, we have established a canonical isomorphism, natural in the ω -category $\mathbb{C} \in \omega\text{-}\underline{\mathbf{Cat}}$, between the category $\mathbf{P}_C(\mathbb{C})$ and the one obtained by applying the finite limit preserving functor $\omega\text{-}\underline{\mathbf{Cat}}(\mathcal{O}(C), -)$ point-wise to the $\omega\text{-}\underline{\mathbf{Cat}}$ -enriched category $\mathbf{P}(\mathbb{C})$.

Observation 260 (generalising Σ to stratified parity complexes). We may extend Σ once more to make it into a functor $\Sigma: \Delta \times \underline{\mathbf{SParity}} \longrightarrow \underline{\mathbf{SParity}}$. If $C \in \underline{\mathbf{SParity}}$ then we define $\Sigma_n C$ to be the stratified parity complex whose underlying parity complex is $\Sigma_n U(C) \in \underline{\mathbf{Parity}}$, and in which an element (x, i) is defined to be thin iff x is thin in C . Under this choice of stratification, it is clear from the definitions given in the last observation that $\Sigma_\alpha C$ preserves thinness for each simplicial operator $\alpha \in \Delta$ and that $\Sigma_m f$ preserves thinness whenever f is a stratified morphism.

Notice that if x is an element of C then the atoms $\langle x \rangle$ of $\mathcal{O}(U(C))$ and $\langle x, i \rangle$ of $\mathcal{O}(\Sigma_1 U(C))$ are related by $\langle x, i \rangle = \langle x \rangle^*$, where $(-)^*$ is the operation on cells which we defined in the last observation. It follows, therefore, that an ω -functor $g: \mathcal{O}(\Sigma_1 U(C)) \longrightarrow \mathbb{C}$ collapses the atoms on thin elements in $\Sigma_1 C$ iff the corresponding ω -functor $\hat{g}: \mathcal{O}(U(C)) \longrightarrow \mathbf{P}(\mathbb{C})(u, v)$ collapses the atoms on thin elements in C . However, we know that $\mathcal{O}(\Sigma_1 C)$ and $\mathcal{O}(C)$ are obtained by taking collapsers of these sets of cells, and thus that the bijection of the last observation extends to the stratified context.

Furthermore, we may extend the work of the last observation to show that the functor $\mathcal{O}(\Sigma_* C): \Delta \longrightarrow \omega\text{-}\underline{\mathbf{Cat}}$ is a cocategory for each stratified parity complex C . To be precise, we argue that in this generalised context the square depicted in display (80) is constructed by

taking collapsers of the nodes in the corresponding square for the underlying parity complex $U(C)$. However we already know, from the last observation, that this latter square is a pushout and so it easily follows, from the fact that as colimits the pushout and collapser constructions commute, that the square for C is also a pushout as required. As before, we adopt the notation P_C to denote the Kan functor associated with this cocategory.

By definition $\Sigma_n U(C)$ is the underlying complex of $\Sigma_n C$ and so we have a natural family of collapsers $q_t: \mathcal{O}(\Sigma_n U(C)) \twoheadrightarrow \mathcal{O}(\Sigma_n C)$ and we may apply Kan's construction (observation 63) to this epimorphic coalgebra map to construct a monomorphic natural transformation $P_C \hookrightarrow P_{U(C)}$. For each ω -category $\mathbb{C}$, this identifies $P_C(\mathbb{C})$ with a subcategory of $P_{U(C)}(\mathbb{C})$ consisting of those arrows $p: \mathcal{O}(\Sigma_1 U(C)) \longrightarrow \mathbb{C}$ which collapse the atoms on thin elements in $\Sigma_1 C$.

Observation 261 (a coalgebraic description of the composite $N_\omega\text{-}\underline{\text{Cat}} \circ P$). Consider the functor $\mathcal{O}(\Sigma_* \tilde{\Delta}(-)): t\Delta \times \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$ and recall that the last two observations demonstrate that $\mathcal{O}(\Sigma_* \tilde{\Delta}[n]?): \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$ is a cocategory for each $[n]_? \in t\Delta$. They also demonstrate that for each $\mathbb{C} \in \omega\text{-}\underline{\text{Cat}}$ the presheaf $\omega\text{-}\underline{\text{Cat}}(\mathcal{O}(\Sigma_1 \tilde{\Delta}(-)), \mathbb{C}): t\Delta^{\text{op}} \longrightarrow \underline{\text{Set}}$ is naturally isomorphic to the complicial set $\coprod_{u,v \in |\mathbb{C}|_0} N_\omega(P(\mathbb{C}))(u, v)$. To be precise, this latter fact follows from our observation that ω -functors $g: \mathcal{O}(\Sigma_1 \tilde{\Delta}[n]?) \longrightarrow (C)$ with $g(\langle 0 \rangle) = u$ and $g(\langle 1 \rangle) = v$ naturally correspond to ω -functors $\hat{g}: \mathcal{O}(\tilde{\Delta}[n]?) \longrightarrow P(\mathbb{C})(u, v)$ which themselves correspond to (thin) n -simplices in $N_\omega(P(\mathbb{C}))(u, v)$ since N_ω is the Kan functor on the $\mathbb{T}_{\underline{\text{Cs}}}$ -coalgebra $\mathcal{O} \circ \tilde{\Delta}: t\Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$.

It follows therefore that the functor $\mathcal{O}(\Sigma_* \tilde{\Delta}(-)): t\Delta \times \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$ is a $\mathbb{T}_{\underline{\text{Cs}}} \otimes \mathbb{T}_{\underline{\text{Cat}}}$ -coalgebra and, as usual, we use the notation $K_{\mathcal{O}(\Sigma_* \tilde{\Delta}(-))}: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ for the corresponding Kan functor. Applying the analysis of observation 259 again, we know that for each $\mathbb{C} \in \omega\text{-}\underline{\text{Cat}}$ the category of (thin) n -arrows in the complicial category $K_{\mathcal{O}(\Sigma_* \tilde{\Delta}(-))}(\mathbb{C})$ is isomorphic to the category obtained by applying the representable $\omega\text{-}\underline{\text{Cat}}(\mathcal{O}(\tilde{\Delta}[n]?), -)$ point-wise to the ω -category enriched category $P(\mathbb{C})$. Abstracting over $[n]_? \in t\Delta$, we therefore see that $K_{\mathcal{O}(\Sigma_* \tilde{\Delta}(-))}(\mathbb{C})$ is isomorphic to the complicially enriched category constructed by applying N_ω point-wise to $P(\mathbb{C})$. In other words, the composite functor $N_\omega\text{-}\underline{\text{Cat}} \circ P: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}\text{-}\underline{\text{Cat}}$ may be represented as the Kan functor on our coalgebra $\mathcal{O}(\Sigma_* \tilde{\Delta}(-))$.

Observation 262 (a coalgebraic description of the composite $\mathbb{P}^r \circ N_\omega$). As discussed in observation 177, the prism functor $\mathbb{P}^r: \underline{\text{Cs}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ bears an external description as the functor obtained by applying Kan's construction (observation 63) to the f-almost $\mathbb{T}_{\underline{\text{Cs}}} \otimes \mathbb{T}_{\underline{\text{Cat}}}$ -coalgebra $\Delta \otimes \Delta_{\mathbb{P}}: t\Delta \times \Delta \longrightarrow \underline{\text{Strat}}$. Of course the left adjoint functor $F_\omega: \underline{\text{Strat}} \longrightarrow \omega\text{-}\underline{\text{Cat}}$ preserves colimits and, by Street's theorem 249, it also carries f-extensions to isomorphisms. So it follows that it carries f-almost colimits to colimits and consequently that we may apply it point-wise to the f-almost coalgebra $\Delta \otimes \Delta_{\mathbb{P}}$ to give a genuine $\mathbb{T}_{\underline{\text{Cs}}} \otimes \mathbb{T}_{\underline{\text{Cat}}}$ -coalgebra $F_\omega \circ (\Delta \otimes \Delta_{\mathbb{P}}): t\Delta \times \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$. Furthermore, the adjunction $F_\omega \dashv N_\omega$ provides us with isomorphisms

$$\begin{aligned} K_{F_\omega \circ (\Delta \otimes \Delta_{\mathbb{P}})}(\mathbb{C}) &\stackrel{\text{def}}{=} \omega\text{-}\underline{\text{Cat}}(F_\omega(\Delta(-) \otimes \Delta_{\mathbb{P}}(*)), \mathbb{C}) \\ &\cong \underline{\text{Strat}}(\Delta(-) \otimes \Delta_{\mathbb{P}}(*), N_\omega(\mathbb{C})) \stackrel{\text{def}}{=} \mathbb{P}^r(N_\omega(\mathbb{C})) \end{aligned}$$

in $\mathbb{T}_{\underline{\text{Cs}}} \otimes \mathbb{T}_{\underline{\text{Cat}}}\text{-}\underline{\text{Alg}} \cong \underline{\text{Cat}}(\underline{\text{Cs}})$ which are natural in $\mathbb{C} \in \omega\text{-}\underline{\text{Cat}}$. In other words, we've shown that we may represent the composite $\mathbb{P}^r \circ N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ as the Kan functor on the coalgebra $F_\omega \circ (\Delta \otimes \Delta_{\mathbb{P}})$.

Now applying corollary 256 it is easy to see that each ω -category $F_\omega(\Delta[n]_? \otimes \Delta_{\mathbb{P}}[m])$ is canonically isomorphic to $\mathcal{O}(\tilde{\Delta}[n]_? \times \text{Th}_1(\tilde{\Delta}[m]))$ (see observation 242 for a discussion of the functor Th_n on SParity) and that these isomorphisms are natural in $[n]_? \in t\Delta$ and $[m] \in \Delta$. In other words, we could equally well describe $\mathbb{P}^r \circ N_\omega$ as the Kan functor on the coalgebra $\mathcal{O}(\tilde{\Delta}(-) \times \text{Th}_1(\tilde{\Delta}(*)))$: $t\Delta \times \Delta \longrightarrow \omega\text{-Cat}$.

Observation 263 (and its sub-functor $\mathbb{P}^r \circ N_\omega$: $\omega\text{-Cat} \longrightarrow \text{Cs-Cat}$). Let $(\Delta[n]_? \otimes \Delta[1])'$ denote the stratified set constructed from $\Delta[n]_? \otimes \Delta[1]$ by making thin all non-degenerate r -simplices ($r = 1, 2, \dots$) of the form $(\alpha, \varepsilon_i^1 \circ \eta^r)$. Then we know, from observation 180, that if A is a complicial set then a (thin) r -arrow of $p: \Delta[n]_? \otimes \Delta[1] \longrightarrow A$ of $\mathbb{P}^r(A)$ is in the subcategory $\mathbb{P}^r(A)$ iff it may be lifted to a stratified map with domain $(\Delta[n]_? \otimes \Delta[1])'$.

We may define a corresponding stratified parity complex $(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1])'$ by making thin all elements of $\tilde{\Delta}[n]_? \times \tilde{\Delta}[1]$ of the form $(v_0 \dots v_p, w_0)$ with $p > 0$. Arguing as in corollary 256 we have a square

$$\begin{array}{ccc} F_\omega(\Delta[n]_? \otimes \Delta[1]) & \xrightarrow{\cong} & \mathcal{O}(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1]) \\ F_\omega(\subseteq_e) \downarrow & & \downarrow \mathcal{O}(\subseteq_e) \\ F_\omega(\Delta[n]_? \otimes \Delta[1])' & \xrightarrow[\cong]{\subseteq_e} & \mathcal{O}(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1])' \end{array} \quad (81)$$

in which the upper horizontal is the isomorphism of that corollary and the verticals are collapsers of the sets of cells associated with the extra thin elements in the dashed versions of these stratified structures (as discussed in observations 239 and 246). From the description of the upper horizontal given in corollary 256 it is clear that it restricts to a bijection between the sets of cells collapsed by these vertical ω -functors and thus that it induces the (dashed) isomorphism at the bottom of this square.

Returning to the last observation, we know that (thin) n -arrows of $\mathbb{P}^r(N_\omega(\mathbb{C}))$ correspond naturally to ω -functors $p: \mathcal{O}(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1]) \longrightarrow \mathbb{C}$. It follows therefore, from the comment in the first paragraph of this observation and the square in display (81), that this ω -functor represents a (thin) n -arrow in the subcategory $\mathbb{P}^r(N_\omega(\mathbb{C}))$ iff it factors through the collapser $\mathcal{O}(\subseteq_e): \mathcal{O}(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1]) \longrightarrow \mathcal{O}(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1])'$.

Observation 264 (comparing $C \times \text{Th}_1(\tilde{\Delta}[m])$ and $\Sigma_m C$). If C is a parity complex, we may define a parity complex morphism $k^{C,m}: C \times \tilde{\Delta}[m] \longrightarrow \Sigma_m C$ by:

$$k^{C,m}(x, w_0 \dots w_q) = \begin{cases} \{w_0\} & \text{when } q = 0 \text{ and } x \in C_0, \\ \{x\} \times (w_0, w_1] & \text{when } q = 1, \\ \emptyset & \text{otherwise.} \end{cases}$$

This clearly respects dimensions and it is a matter of simple case-wise verification to check that it satisfies the movement condition given in lemma 231. In a similar fashion it is also easily demonstrated that this family of parity complex morphisms is natural in $C \in \text{Parity}$ and $[m] \in \Delta$. We leave the completely routine verification of these facts for the delectation of the reader.

Having defined this parity complex morphism, it is worth spending a few moments studying the action of the associated ω -functor $\mathcal{O}(k^{C,m}): \mathcal{O}(C \times \tilde{\Delta}[m]) \longrightarrow \mathcal{O}(\Sigma_m C)$ on the atoms of its domain. As we may easily establish directly, or verify by arguing as in observation 243 using the result of corollary 234, we have

- $\mathcal{O}(k^{C,m})(\langle x, w_0 \rangle) = \langle w_0 \rangle$, and
- $\mathcal{O}(k^{C,m})(\langle x, w_0 \dots w_q \rangle) = \langle x, w_q \rangle * \langle x, w_q - 1 \rangle * \dots * \langle x, w_0 + 1 \rangle$ whenever $q > 0$ and in particular if $q = 1$ and $w_1 = w_0 + 1$ this is equal to the atom $\langle x, w_1 \rangle$.

Now suppose that C is a stratified parity complex and consider the product $C \times \text{Th}_1(\tilde{\Delta}[m])$ in $\underline{\text{SParity}}$. By definition, the element $(x, w_0 \dots w_q)$ is thin in $C \times \text{Th}_1(\tilde{\Delta}[m])$ iff either $w_0 \dots w_q$ is thin in $\text{Th}_1(\tilde{\Delta}[n])$, which happens when $q > 1$ in which case $k^{\text{U}(C),m}(x, w_0 \dots w_q)$ is empty, or x is thin in C , in which case its dimension is > 0 and so either $k^{\text{U}(C),m}(x, w_0 \dots w_q)$ is empty or $q = 1$ and it is equal to the set $\{x\} \times (w_0, w_1]$ of thin elements in $\Sigma_m C$. It follows that $k^{\text{U}(C),m}$ satisfies the thinness preservation condition making it into a stratified morphism $k^{C,m}: C \times \text{Th}_1(\tilde{\Delta}[m]) \longrightarrow \Sigma_m C$. Forgetting stratifications, we see that the naturality of this stratified morphism in $C \in \underline{\text{SParity}}$ and $[n] \in \Delta$ follows directly from its established naturality on underlying parity complexes.

Lemma 265. *The stratified morphism $k: \tilde{\Delta}[n]_? \times \tilde{\Delta}[1] \longrightarrow \Sigma_1 \tilde{\Delta}[n]_?$ of the last observation lifts to one with domain $(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1])'$ and the ω -functor obtained by applying $\mathcal{O}$ to this lifted morphism provides an isomorphism between $\mathcal{O}(\tilde{\Delta}[n]_? \times \tilde{\Delta}[1])'$ and $\mathcal{O}(\Sigma_1 \tilde{\Delta}[n]_?)$.*

Proof. The first part simply follows from the definition of k which tells us that $k(v_0 \dots v_p, w_0) = \emptyset$ whenever $p > 0$.

To prove the remainder, first consider the case $[n]_? = [n]$ and recall, from the last observation, that $\mathcal{O}(k): \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])' \longrightarrow \mathcal{O}(\Sigma_1 \tilde{\Delta}[n])$ maps $\langle 0, w_0 \rangle \in \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])'$ to $\langle w_0 \rangle \in \mathcal{O}(\Sigma_1 \tilde{\Delta}[n])$ and $\langle \underline{v}, 01 \rangle \in \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])'$ to $\langle \underline{v}, 1 \rangle \in \mathcal{O}(\Sigma_1 \tilde{\Delta}[n])$. It follows that this ω -functor restricts to a bijection between the subset $S = \{\langle 0, w_0 \rangle \mid w_0 = 0, 1\} \cup \{\langle \underline{v}, 01 \rangle \mid \underline{v} \in \tilde{\Delta}[n]\}$ of cells in $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])'$ and the set of atoms of the freely generated category $\mathcal{O}(\Sigma_1 \tilde{\Delta}[n])$. Notice however that any element $(v_0 v_1, w_0)$ in $(\tilde{\Delta}[n] \times \tilde{\Delta}[1])'$ is thin and so it follows that $\langle v_0 v_1, w_0 \rangle$ is a 0-cell in $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])'$ and therefore that its 0-source $\langle v_0, w_0 \rangle$ and 0-target $\langle v_1, w_0 \rangle$ are equal. From this it follows that any cell $\langle v_0, w_0 \rangle$ is in S and thus that this subset is a weak generator for $\mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])'$ since it contains the weak generator $\langle (\tilde{\Delta}[n] \times \tilde{\Delta}[1])' \rangle$ discussed in observation 239. Now we may apply lemma 223 to show that $\mathcal{O}(k): \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])' \longrightarrow \mathcal{O}(\Sigma_1 \tilde{\Delta}[n])$ is an isomorphism as required.

To extend this to the case $[n]_? = [n]_t$ consider the commutative square

$$\begin{array}{ccc} \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])' & \xrightarrow[\cong]{\mathcal{O}(k)} & \mathcal{O}(\Sigma_1 \tilde{\Delta}[n]) \\ \mathcal{O}(\subseteq_e) \downarrow & & \downarrow \mathcal{O}(\subseteq_e) \\ \mathcal{O}(\tilde{\Delta}[n]_t \times \tilde{\Delta}[1])' & \xrightarrow{\mathcal{O}(k)} & \mathcal{O}(\Sigma_1 \tilde{\Delta}[n]_t) \end{array}$$

and again, as discussed in observation 239, the verticals here are collapsers of the cells $\langle 01 \dots n, 01 \rangle \in \mathcal{O}(\tilde{\Delta}[n] \times \tilde{\Delta}[1])'$ and $\langle 01 \dots n, 1 \rangle \in \mathcal{O}(\Sigma_1 \tilde{\Delta}[n])$ respectively. The horizontal isomorphism at the top of this square maps these cells to each other and it follows that we may apply the collapse property of the right hand vertical to induce an inverse to the lower horizontal as required. $\square$

Finally we arrive at our ultimate destination.

Theorem 266 (the Street-Roberts conjecture). *We may construct the essentially commutative square illustrated in display (78) and thus apply the argument described in observation 257*

to show that Street's nerve functor $N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}$ provides an equivalence between the categories of ω -categories and complicial sets.

Proof. Observation 261 showed that we may represent $N_\omega\text{-}\underline{\text{Cat}} \circ P: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cs}}\text{-}\underline{\text{Cat}}$ as the Kan functor associated with a coalgebra $\Sigma_*\tilde{\Delta}(-): t\Delta \times \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$. Similarly observation 262 demonstrated that the composite $\mathbb{P}^r \circ N_\omega: \omega\text{-}\underline{\text{Cat}} \longrightarrow \underline{\text{Cat}}(\underline{\text{Cs}})$ may be represented by a coalgebra $\tilde{\Delta}(-) \times \text{Th}_1(\tilde{\Delta}(*)): t\Delta \times \Delta \longrightarrow \omega\text{-}\underline{\text{Cat}}$. Furthermore, observation 264 provides us with a coalgebra map from the latter to the former which we can use to construct a comparison 2-cell

$$\begin{array}{ccc}
 \omega\text{-}\underline{\text{Cat}} & \xrightarrow{P} & (\omega\text{-}\underline{\text{Cat}})\text{-}\underline{\text{Cat}} \\
 \downarrow N_\omega & \Longleftarrow & \downarrow N_\omega\text{-}\underline{\text{Cat}} \\
 & & \underline{\text{Cs}}\text{-}\underline{\text{Cat}} \\
 \underline{\text{Cs}} & \xrightarrow{\mathbb{P}^r} & \underline{\text{Cat}}(\underline{\text{Cs}})
 \end{array} \tag{82}$$

by pre-composition, as discussed in observation 63. Now applying observation 263 and lemma 265, we see that for each ω -category $\mathbb{C}$ the component $N_\omega\text{-}\underline{\text{Cat}}(P(\mathbb{C})) \longrightarrow \mathbb{P}^r(N_\omega(\mathbb{C}))$ of this acts as an isomorphism between the complicial set of arrows of $N_\omega\text{-}\underline{\text{Cat}}(P(\mathbb{C}))$ and that of the subcategory $\mathbb{P}^r(N_\omega(\mathbb{C}))$ of $\mathbb{P}^r(N_\omega(\mathbb{C}))$, and thus restricts to an isomorphism of those categories. In other words, the natural transformation in display (82) restricts to provide the isomorphism of display (78) as required. $\square$

REFERENCES

- [1] F. Al-Agl, R. Brown, and R. Steiner. Multiple categories: the equivalence between a globular and cubical approach. *Advances in Mathematics*, 170:71–118, 2002.
- [2] J. Baez and J. Dolan. Higher-dimensional algebra III: n -categories and the algebra of opetopes. *Advances in Mathematics*, 135:145–206, 1998.
- [3] M. Barr and C. Wells. *Toposes, Triples, and Theories*, volume 278 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, New York, 1985.
- [4] M. A. Batanin. Monoidal globular categories as natural environment for the theory of weak n -categories. *Advances in Mathematics*, 136:39–103, 1998.
- [5] R. Brown. Crossed complexes and homotopy groupoids as non-commutative tools for higher dimensional local-to-global problems. *Contemporary Mathematics*, to appear, 2004.
- [6] R. Brown and P. J. Higgins. The equivalence of crossed complexes and ∞ -groupoids. *Cahiers topologie et géométrie différentielle catégoriques*, 22:371–386, 1981.
- [7] R. Brown and G. H. Mosa. Double categories, 2-categories, thin structures and connections. *Theory and Applications of Categories*, 5:163–175, 1999.
- [8] S. Crans. *On Combinatorial Models for Higher Dimensional Homotopies*. PhD thesis, Universiteit Utrecht, 1995.
- [9] M. K. Dakin. *Kan Complexes and Multiple Groupoid Structures*. PhD thesis, University of Wales, 1977.
- [10] B. Day. On closed categories of functors. In *Reports of the Midwest Category Seminar IV*, volume 137 of *Lecture Notes in Mathematics*, pages 1–38. Springer-Verlag, 1970.
- [11] P. J. Freyd and G. M. Kelly. Categories of continuous functors I. *Journal of Pure and Applied Algebra*, 2:169–191, 1972.
- [12] P. Gabriel and F. Ulmer. *Lokal Präsentierbare Kategorien*, volume 221 of *Lecture Notes in Mathematics*. Springer-Verlag, 1971.
- [13] P. Gabriel and M. Zisman. *Calculus of Fractions and Homotopy Theory*. Number 35 in *Ergebnisse der Math.* Springer-Verlag, Berlin, 1967.
- [14] R. Gordon, A. J. Power, and R. H. Street. Coherence for tricategories. *Memoirs of the American Math Society*, 117(558), 1995.
- [15] J. W. Gray. *Formal Category Theory: Adjointness for 2-Categories*. Number 391 in *Lecture Notes in Mathematics*. Springer-Verlag, Berlin and New York, 1974.
- [16] A. Joyal. Disks, duality and θ -categories. *Preprint and Talk at AMS Meeting in Montréal*, 1997.
- [17] D. M. Kan. Adjoint functors. *Transactions of the American Mathematical Society*, 87:294–329, 1958.
- [18] G. M. Kelly. A unified treatment of transfinite constructions for free algebras, free monoids, colimits, associated sheaves and so on. *Bulletin of the Australian Mathematical Society*, 22:1–83, 1980.
- [19] G. M. Kelly. *Basic Concepts of Enriched Category Theory*. Number 64 in *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1982.
- [20] G. M. Kelly and R. H. Street. Review of the elements of 2-categories. In *Category Seminar*, pages 75–103. Springer-Verlag, 1974.
- [21] J. F. Kennison. On limit-preserving functors. *Illinois Journal of Mathematics*, pages 616–619, 1968.
- [22] S. M. Lane. *Categories for the Working Mathematician*. Springer-Verlag, New York, 1971.
- [23] T. Leinster. A survey of definitions of n -category. *Theory and Applications of Categories*, 10(1):1–70, 2002.
- [24] J. P. May. *Simplicial Objects in Algebraic Topology*. D. Van Nostrand, 1967.
- [25] A. J. Power. A 2-categorical pasting theorem. *Journal of Algebra*, 129:439–445, 1988.
- [26] J. E. Roberts. Mathematical aspects of local cohomology. In *Proceedings of the Colloquium on Operator Algebras and their Application to Mathematical Physics*, Marseille, 1977.
- [27] J. E. Roberts. Complicial sets. *handwritten manuscript*, 1978.
- [28] R. Steiner. Tensor products of infinity-categories. Technical report, University of Glasgow, 1991.
- [29] R. H. Street. The formal theory of monads. *Journal of Pure and Applied Algebra*, 2:149–168, 1972.
- [30] R. H. Street. Fibrations in bicategories. *Cahiers topologie et géométrie différentielle catégoriques*, 21:111–120, 1980.
- [31] R. H. Street. The algebra of oriented simplexes. *Journal of Pure and Applied Algebra*, 49:283–335, 1987.
- [32] R. H. Street. Fillers for nerves. *Lecture Notes in Mathematics*, 1348:337–341, 1988.
- [33] R. H. Street. Parity complexes. *Cahiers topologie et géométrie différentielle catégoriques*, 32:315–343, 1991.

- [34] R. H. Street. Parity complexes: Corrigenda. *Cahiers topologie et géométrie différentielle catégoriques*, 35:359–361, 1994.
- [35] R. H. Street. Categorical structures. In M. Hazewinkel, editor, *Handbook of Algebra*, volume 1, pages 529–577. Elsevier Science, Amsterdam, 1996.
- [36] R. H. Street. Weak omega-categories. In *Diagrammatic Morphisms and Applications*, volume 318 of *Contemporary Mathematics*, pages 207–213. American Mathematical Society, 2003.

CENTRE OF AUSTRALIAN CATEGORY THEORY, MACQUARIE UNIVERSITY, NSW 2109, AUSTRALIA
E-mail address: domv@ics.mq.edu.au